

$\Lambda_b(5920)^0$ $J^P = \frac{3}{2}^-$

Status: ***

Quantum numbers are based on quark model expectations.

 $\Lambda_b(5920)^0$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-------------------------|-----------|----------------------------|
| 5920.09±0.17 OUR AVERAGE | | | |
| 5920.09±0.02±0.17 | ¹ AAIJ | 20Q LHCb | $p\bar{p}$ at 7, 8, 13 TeV |
| 5920.16±0.07±0.17 | ² SIRUNYAN | 20K CMS | $p\bar{p}$ at 13 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |
| 5919.4 ± 0.5 ± 0.2 | ^{3,4} AALTONEN | 13V CDF | $p\bar{p}$ at 1.96 TeV |
| 5920.00±0.09±0.17 | ^{5,6} AAIJ | 12AL LHCb | Repl. by AAIJ 20Q |
| ¹ AAIJ 20Q measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) = 300.492 \pm 0.019 \pm 0.010$ MeV. We have adjusted the measurement to our best value of $m(\Lambda_b^0) = 5619.60 \pm 0.17$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values. | | | |
| ² SIRUNYAN 20K measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) = 300.56 \pm 0.07 \pm 0.01$ MeV. We have adjusted the measurement to our best value of $m(\Lambda_b^0) = 5619.60 \pm 0.17$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values. | | | |
| ³ Measured in $\Lambda_b(5920)^0 \rightarrow \Lambda_b^0 \pi^+ \pi^-$ decays with $17.3^{+5.3}_{-4.6}$ events, with a significance of 3.5 sigma. | | | |
| ⁴ AALTONEN 13V measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) - 2m(\pi) = 20.68 \pm 0.35 \pm 0.30$ MeV. We have adjusted the measurement to our best values of $m(\Lambda_b^0) = 5619.60 \pm 0.17$ MeV and $m(\pi) = 139.57039 \pm 0.00018$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values. | | | |
| ⁵ Observed in $\Lambda_b(5920)^0 \rightarrow \Lambda_b^0 \pi^+ \pi^-$ decays with 52.5 ± 8.1 candidates with a significance of 10.2 sigma. | | | |
| ⁶ AAIJ 12AL measures $m(\Lambda_b(5920)^0) - m(\Lambda_b^0) = 300.40 \pm 0.08 \pm 0.04$ MeV. We have adjusted the measurement to our best value of $m(\Lambda_b^0) = 5619.60 \pm 0.17$ MeV. Our first error is their experiment's error and our second error is the systematic error from using our best values. | | | |

 $\Lambda_b(5920)^0$ WIDTH

| VALUE (MeV) | CL% | DOCUMENT ID | TECN | COMMENT |
|---|-----|-------------|-----------|----------------------------|
| <0.19 | 90 | AAIJ | 20Q LHCb | $p\bar{p}$ at 7, 8, 13 TeV |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| <0.63 | 90 | AAIJ | 12AL LHCb | Repl. by AAIJ 20Q |

 $\Lambda_b(5920)^0$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|--|--------------------------------|
| $\Gamma_1 \quad \Lambda_b^0 \pi^+ \pi^-$ | seen |

$\Lambda_b(5920)^0$ BRANCHING RATIOS

$\Gamma(\Lambda_b^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$

| VALUE | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|----------------------------|
| seen | AAIJ | 20Q | LHCb $p p$ at 7, 8, 13 TeV |
| seen | SIRUNYAN | 20K | LHCb $p p$ at 13 TeV |
| seen | AALTOMEN | 13V | CDF $p\bar{p}$ at 1.96 TeV |
| seen | AAIJ | 12AL | LHCb $p p$ at 7 TeV |

Γ_1/Γ

$\Lambda_b(5920)^0$ REFERENCES

| | | | | |
|----------|------|----------------|-----------------------------|----------------|
| AAIJ | 20Q | JHEP 2006 136 | R. Aaij <i>et al.</i> | (LHCb Collab.) |
| SIRUNYAN | 20K | PL B803 135345 | A.M. Sirunyan <i>et al.</i> | (CMS Collab.) |
| AALTOMEN | 13V | PR D88 071101 | T. Aaltonen <i>et al.</i> | (CDF Collab.) |
| AAIJ | 12AL | PRL 109 172003 | R. Aaij <i>et al.</i> | (LHCb Collab.) |