

STRANGE MESONS ($S = \pm 1$, $C = B = 0$)

$K^+ = u\bar{s}$, $K^0 = d\bar{s}$, $\bar{K}^0 = \bar{d}s$, $K^- = \bar{u}s$, similarly for K^* 's

K^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 493.677 \pm 0.016$ MeV ^[a] ($S = 2.8$)

Mean life $\tau = (1.2380 \pm 0.0020) \times 10^{-8}$ s ($S = 1.8$)

$$c\tau = 3.711 \text{ m}$$

***CPT* violation parameters ($\Delta = \text{rate difference/sum}$)**

$$\Delta(K^\pm \rightarrow \mu^\pm \nu_\mu) = (-0.27 \pm 0.21)\%$$

$$\Delta(K^\pm \rightarrow \pi^\pm \pi^0) = (0.4 \pm 0.6)\% \quad [b]$$

***CP* violation parameters ($\Delta = \text{rate difference/sum}$)**

$$\Delta(K^\pm \rightarrow \pi^\pm e^+ e^-) = (-2.2 \pm 1.6) \times 10^{-2}$$

$$\Delta(K^\pm \rightarrow \pi^\pm \mu^+ \mu^-) = 0.010 \pm 0.023$$

$$\Delta(K^\pm \rightarrow \pi^\pm \pi^0 \gamma) = (0.0 \pm 1.2) \times 10^{-3}$$

$$\Delta(K^\pm \rightarrow \pi^\pm \pi^+ \pi^-) = (0.04 \pm 0.06)\%$$

$$\Delta(K^\pm \rightarrow \pi^\pm \pi^0 \pi^0) = (-0.02 \pm 0.28)\%$$

***T* violation parameters**

$$K^+ \rightarrow \pi^0 \mu^+ \nu_\mu \quad P_T = (-1.7 \pm 2.5) \times 10^{-3}$$

$$K^+ \rightarrow \mu^+ \nu_\mu \gamma \quad P_T = (-0.6 \pm 1.9) \times 10^{-2}$$

$$K^+ \rightarrow \pi^0 \mu^+ \nu_\mu \quad \text{Im}(\xi) = -0.006 \pm 0.008$$

Slope parameter g ^[c]

(See Particle Listings for quadratic coefficients and alternative parametrization related to $\pi\pi$ scattering)

$$K^\pm \rightarrow \pi^\pm \pi^+ \pi^- \quad g = -0.21134 \pm 0.00017$$

$$(g_+ - g_-) / (g_+ + g_-) = (-1.5 \pm 2.2) \times 10^{-4}$$

$$K^\pm \rightarrow \pi^\pm \pi^0 \pi^0 \quad g = 0.626 \pm 0.007$$

$$(g_+ - g_-) / (g_+ + g_-) = (1.8 \pm 1.8) \times 10^{-4}$$

K^\pm decay form factors ^[d,e]

Assuming μ -e universality

$$\lambda_+(K_{\mu 3}^+) = \lambda_+(K_{e 3}^+) = (2.959 \pm 0.025) \times 10^{-2}$$

$$\lambda_0(K_{\mu 3}^+) = (1.76 \pm 0.25) \times 10^{-2} \quad (S = 2.7)$$

Not assuming μ -e universality

$$\lambda_+(K_{e3}^+) = (2.956 \pm 0.025) \times 10^{-2}$$

$$\lambda_+(K_{\mu 3}^+) = (3.09 \pm 0.25) \times 10^{-2} \quad (S = 1.5)$$

$$\lambda_0(K_{\mu 3}^+) = (1.73 \pm 0.27) \times 10^{-2} \quad (S = 2.6)$$

K_{e3} form factor quadratic fit

$$\lambda'_+ (K_{e3}^\pm) \text{ linear coeff.} = (2.59 \pm 0.04) \times 10^{-2}$$

$$\lambda''_+ (K_{e3}^\pm) \text{ quadratic coeff.} = (0.186 \pm 0.021) \times 10^{-2}$$

$$\lambda'_+ (\text{LINEAR } K_{\mu 3}^\pm \text{ FORM FACTOR FROM QUADRATIC FIT}) = (24 \pm 4) \times 10^{-3}$$

$$\lambda''_+ (\text{QUADRATIC } K_{\mu 3}^\pm \text{ FORM FACTOR}) = (1.8 \pm 1.5) \times 10^{-3}$$

$$M_V (\text{VECTOR POLE MASS FOR } K_{e3}^\pm \text{ DECAY}) = 890.3 \pm 2.8 \text{ MeV}$$

$$M_V (\text{VECTOR POLE MASS FOR } K_{\mu 3}^\pm \text{ DECAY}) = 878 \pm 12 \text{ MeV}$$

$$M_S (\text{SCALAR POLE MASS FOR } K_{\mu 3}^\pm \text{ DECAY}) = 1215 \pm 50 \text{ MeV}$$

$$\Lambda_+ (\text{DISPERSIVE VECTOR FORM FACTOR IN } K_{e3}^\pm \text{ DECAY}) = (2.460 \pm 0.017) \times 10^{-2}$$

$$\Lambda_+ (\text{DISPERSIVE VECTOR FORM FACTOR IN } K_{\mu 3}^\pm \text{ DECAY}) = (25.4 \pm 0.9) \times 10^{-3}$$

$$\ln(C) (\text{DISPERSIVE SCALAR FORM FACTOR in } K_{\mu 3}^\pm \text{ decays }) = (182 \pm 16) \times 10^{-3}$$

$$K_{e3}^+ \quad |f_S/f_+| = (-0.08^{+0.34}_{-0.40}) \times 10^{-2}$$

$$K_{e3}^+ \quad |f_T/f_+| = (-1.2^{+1.3}_{-1.1}) \times 10^{-2}$$

$$K_{\mu 3}^+ \quad |f_S/f_+| = (0.2 \pm 0.6) \times 10^{-2}$$

$$K_{\mu 3}^+ \quad |f_T/f_+| = (-0.1 \pm 0.7) \times 10^{-2}$$

$$K^+ \rightarrow e^+ \nu_e \gamma \quad |F_A + F_V| = 0.133 \pm 0.008 \quad (S = 1.3)$$

$$K^+ \rightarrow \mu^+ \nu_\mu \gamma \quad |F_A + F_V| = 0.165 \pm 0.013$$

$$K^+ \rightarrow e^+ \nu_e \gamma \quad |F_A - F_V| < 0.49, \text{ CL} = 90\%$$

$$K^+ \rightarrow \mu^+ \nu_\mu \gamma \quad |F_A - F_V| = -0.153 \pm 0.033 \quad (S = 1.1)$$

Charge radius

$$\langle r \rangle = 0.560 \pm 0.031 \text{ fm}$$

Forward-backward asymmetry

$$A_{FB}(K_{\pi \mu \mu}^\pm) = \frac{\Gamma(\cos(\theta_{K\mu}) > 0) - \Gamma(\cos(\theta_{K\mu}) < 0)}{\Gamma(\cos(\theta_{K\mu}) > 0) + \Gamma(\cos(\theta_{K\mu}) < 0)} < 0.9 \times 10^{-2}, \text{ CL} = 90\%$$

K^- modes are charge conjugates of the modes below.

| K^+ DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ p | |
|---|--|-------------------------|-----|
| | | Confidence level(MeV/c) | |
| Leptonic and semileptonic modes | | | |
| $e^+ \nu_e$ | (1.582 ± 0.007) $\times 10^{-5}$ | | 247 |
| $\mu^+ \nu_\mu$ | (63.56 ± 0.11) % | S=1.2 | 236 |
| $\pi^0 e^+ \nu_e$ Called K_{e3}^+ . | (5.07 ± 0.04) % | S=2.1 | 228 |
| $\pi^0 \mu^+ \nu_\mu$ Called $K_{\mu 3}^+$. | (3.352 ± 0.033) % | S=1.9 | 215 |
| $\pi^0 \pi^0 e^+ \nu_e$ | (2.55 ± 0.04) $\times 10^{-5}$ | S=1.1 | 206 |
| $\pi^+ \pi^- e^+ \nu_e$ | (4.247 ± 0.024) $\times 10^{-5}$ | | 203 |
| $\pi^+ \pi^- \mu^+ \nu_\mu$ | (1.4 ± 0.9) $\times 10^{-5}$ | | 151 |
| $\pi^0 \pi^0 \pi^0 e^+ \nu_e$ | < 3.5×10^{-6} | CL=90% | 135 |
| Hadronic modes | | | |
| $\pi^+ \pi^0$ | (20.67 ± 0.08) % | S=1.2 | 205 |
| $\pi^+ \pi^0 \pi^0$ | (1.760 ± 0.023) % | S=1.1 | 133 |
| $\pi^+ \pi^+ \pi^-$ | (5.583 ± 0.024) % | | 125 |
| Leptonic and semileptonic modes with photons | | | |
| $\mu^+ \nu_\mu \gamma$ | [f,g] (6.2 ± 0.8) $\times 10^{-3}$ | | 236 |
| $\mu^+ \nu_\mu \gamma (\text{SD}^+)$ | [d,h] (1.33 ± 0.22) $\times 10^{-5}$ | | - |
| $\mu^+ \nu_\mu \gamma (\text{SD}^+ \text{INT})$ | [d,h] < 2.7×10^{-5} | CL=90% | - |
| $\mu^+ \nu_\mu \gamma (\text{SD}^- + \text{SD}^- \text{INT})$ | [d,h] < 2.6×10^{-4} | CL=90% | - |
| $e^+ \nu_e \gamma$ | (9.9 ± 1.0) $\times 10^{-6}$ | | 247 |
| $\pi^0 e^+ \nu_e \gamma$ | [f,g] (2.66 ± 0.09) $\times 10^{-4}$ | | 228 |
| $\pi^0 e^+ \nu_e \gamma (\text{SD})$ | [d,h] < 5.3×10^{-5} | CL=90% | 228 |
| $\pi^0 \mu^+ \nu_\mu \gamma$ | [f,g] (1.25 ± 0.25) $\times 10^{-5}$ | | 215 |
| $\pi^0 \pi^0 e^+ \nu_e \gamma$ | < 5×10^{-6} | CL=90% | 206 |
| Hadronic modes with photons or $\ell\bar{\ell}$ pairs | | | |
| $\pi^+ \pi^0 \gamma (\text{INT})$ | (- 4.2 ± 0.9) $\times 10^{-6}$ | | - |
| $\pi^+ \pi^0 \gamma (\text{DE})$ | [f,i] (6.0 ± 0.4) $\times 10^{-6}$ | | 205 |
| $\pi^+ \pi^0 e^+ e^-$ | (4.24 ± 0.14) $\times 10^{-6}$ | | 205 |
| $\pi^+ \pi^0 \pi^0 \gamma$ | [f,g] ($7.6 \begin{array}{l} +6.0 \\ -3.0 \end{array} \pm 0.5$) $\times 10^{-6}$ | | 133 |
| $\pi^+ \pi^+ \pi^- \gamma$ | [f,g] (7.1 ± 0.5) $\times 10^{-6}$ | | 125 |
| $\pi^+ \gamma \gamma$ | [f] (1.01 ± 0.06) $\times 10^{-6}$ | | 227 |
| $\pi^+ 3\gamma$ | [f] < 1.0×10^{-4} | CL=90% | 227 |
| $\pi^+ e^+ e^- \gamma$ | (1.19 ± 0.13) $\times 10^{-8}$ | | 227 |
| Leptonic modes with $\ell\bar{\ell}$ pairs | | | |
| $e^+ \nu_e \nu \bar{\nu}$ | < 6×10^{-5} | CL=90% | 247 |
| $\mu^+ \nu_\mu \nu \bar{\nu}$ | < 1.0×10^{-6} | CL=90% | 236 |

| | | | | |
|-----------------------------|---|-----------------|--------------------|------------|
| $e^+ \nu_e e^+ e^-$ | (| 2.48 ± 0.20 |) $\times 10^{-8}$ | 247 |
| $\mu^+ \nu_\mu e^+ e^-$ | (| 7.06 ± 0.31 |) $\times 10^{-8}$ | 236 |
| $e^+ \nu_e \mu^+ \mu^-$ | (| 1.7 ± 0.5 |) $\times 10^{-8}$ | 223 |
| $\mu^+ \nu_\mu \mu^+ \mu^-$ | < | 4.1 | $\times 10^{-7}$ | CL=90% 185 |

**Lepton family number (*LF*), Lepton number (*L*), $\Delta S = \Delta Q$ (*SQ*)
violating modes, or $\Delta S = 1$ weak neutral current (*S1*) modes**

| | | | | | |
|-----------------------------------|----------------|----------------|------------------------|---------------------|------------|
| $\pi^+ \pi^+ e^- \bar{\nu}_e$ | <i>SQ</i> | < | 1.3 | $\times 10^{-8}$ | CL=90% 203 |
| $\pi^+ \pi^+ \mu^- \bar{\nu}_\mu$ | <i>SQ</i> | < | 3.0 | $\times 10^{-6}$ | CL=95% 151 |
| $\pi^+ e^+ e^-$ | <i>S1</i> | (| 3.00 ± 0.09 |) $\times 10^{-7}$ | 227 |
| $\pi^+ \mu^+ \mu^-$ | <i>S1</i> | (| 9.17 ± 0.14 |) $\times 10^{-8}$ | S=1.8 172 |
| $\pi^+ \nu \bar{\nu}$ | <i>S1</i> | (| $1.14^{+0.40}_{-0.33}$ |) $\times 10^{-10}$ | 227 |
| $\pi^+ \pi^0 \nu \bar{\nu}$ | <i>S1</i> | < | 4.3 | $\times 10^{-5}$ | 205 |
| $\mu^- \nu e^+ e^+$ | <i>LF</i> | < | 2.1 | $\times 10^{-8}$ | CL=90% 236 |
| $\mu^+ \nu_e$ | <i>LF</i> | [<i>j</i>] < | 4 | $\times 10^{-3}$ | CL=90% 236 |
| $\pi^+ \mu^+ e^-$ | <i>LF</i> | < | 1.3 | $\times 10^{-11}$ | CL=90% 214 |
| $\pi^+ \mu^- e^+$ | <i>LF</i> | < | 6.6 | $\times 10^{-11}$ | CL=90% 214 |
| $\pi^- \mu^+ e^+$ | <i>L</i> | < | 4.2 | $\times 10^{-11}$ | CL=90% 214 |
| $\pi^- e^+ e^+$ | <i>L</i> | < | 5.3 | $\times 10^{-11}$ | CL=90% 227 |
| $\pi^- \mu^+ \mu^+$ | <i>L</i> | < | 4.2 | $\times 10^{-11}$ | CL=90% 172 |
| $\pi^- \pi^0 e^+ e^+$ | <i>L</i> | < | 8.5 | $\times 10^{-10}$ | CL=90% 205 |
| $\mu^+ \bar{\nu}_e$ | <i>L</i> | [<i>j</i>] < | 3.3 | $\times 10^{-3}$ | CL=90% 236 |
| $\pi^0 e^+ \bar{\nu}_e$ | <i>L</i> | < | 3 | $\times 10^{-3}$ | CL=90% 228 |
| $\pi^+ \gamma$ | [<i>k</i>] < | 2.3 | $\times 10^{-9}$ | CL=90% | 227 |

K^0

$$I(J^P) = \frac{1}{2}(0^-)$$

50% K_S , 50% K_L

Mass $m = 497.611 \pm 0.013$ MeV (S = 1.2)

$m_{K^0} - m_{K^\pm} = 3.934 \pm 0.020$ MeV (S = 1.6)

Mean square charge radius

$$\langle r^2 \rangle = -0.077 \pm 0.010 \text{ fm}^2$$

T-violation parameters in K^0 - \bar{K}^0 mixing [e]

Asymmetry A_T in K^0 - \bar{K}^0 mixing = $(6.6 \pm 1.6) \times 10^{-3}$

CP-violation parameters

$$\text{Re}(\epsilon) = (1.596 \pm 0.013) \times 10^{-3}$$

CPT-violation parameters [e]

$$\text{Re } \delta = (2.5 \pm 2.3) \times 10^{-4}$$

$$\text{Im } \delta = (-1.5 \pm 1.6) \times 10^{-5}$$

$$\text{Re}(y), K_{e3} \text{ parameter} = (0.4 \pm 2.5) \times 10^{-3}$$

$$\begin{aligned} \text{Re}(x_-), K_{e3} \text{ parameter} &= (-2.9 \pm 2.0) \times 10^{-3} \\ |m_{K^0} - m_{\bar{K}^0}| / m_{\text{average}} &< 6 \times 10^{-19}, \text{ CL} = 90\% \quad [l] \\ (\Gamma_{K^0} - \Gamma_{\bar{K}^0})/m_{\text{average}} &= (8 \pm 8) \times 10^{-18} \end{aligned}$$

Tests of $\Delta S = \Delta Q$

$$\text{Re}(x_+), K_{e3} \text{ parameter} = (-0.9 \pm 3.0) \times 10^{-3}$$

K_S^0

$$I(J^P) = \frac{1}{2}(0^-)$$

Mean life $\tau = (0.8954 \pm 0.0004) \times 10^{-10} \text{ s}$ ($S = 1.1$) Assuming *CPT*

Mean life $\tau = (0.89564 \pm 0.00033) \times 10^{-10} \text{ s}$ Not assuming *CPT*

$c\tau = 2.6844 \text{ cm}$ Assuming *CPT*

CP-violation parameters [n]

$$\text{Im}(\eta_{+-0}) = -0.002 \pm 0.009$$

$$\text{Im}(\eta_{000}) = -0.001 \pm 0.016$$

$$|\eta_{000}| = |A(K_S^0 \rightarrow 3\pi^0)/A(K_L^0 \rightarrow 3\pi^0)| < 0.0088, \text{ CL} = 90\%$$

$$CP \text{ asymmetry } A \text{ in } \pi^+ \pi^- e^+ e^- = (-0.4 \pm 0.8)\%$$

| K_S^0 DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level | p (MeV/c) |
|---|---|-----------------------------------|--------------|
| Hadronic modes | | | |
| $\pi^0 \pi^0$ | $(30.69 \pm 0.05) \%$ | | 209 |
| $\pi^+ \pi^-$ | $(69.20 \pm 0.05) \%$ | | 206 |
| $\pi^+ \pi^- \pi^0$ | $(3.5 \pm 1.1) \times 10^{-7}$ | | 133 |
| Modes with photons or $\ell\bar{\ell}$ pairs | | | |
| $\pi^+ \pi^- \gamma$ | $[g.o] \quad (1.79 \pm 0.05) \times 10^{-3}$ | | 206 |
| $\pi^+ \pi^- e^+ e^-$ | $(4.79 \pm 0.15) \times 10^{-5}$ | | 206 |
| $\pi^0 \gamma \gamma$ | $[o] \quad (4.9 \pm 1.8) \times 10^{-8}$ | | 230 |
| $\gamma \gamma$ | $(2.63 \pm 0.17) \times 10^{-6}$ | S=3.1 | 249 |
| Semileptonic modes | | | |
| $\pi^\pm e^\mp \nu_e$ | $[p] \quad (7.04 \pm 0.08) \times 10^{-4}$ | | 229 |
| <i>CP</i> violating (<i>CP</i>) and $\Delta S = 1$ weak neutral current (<i>S1</i>) modes | | | |
| $3\pi^0$ | $CP \quad < 2.6 \times 10^{-8}$ | CL=90% | 139 |
| $\mu^+ \mu^-$ | $S1 \quad < 2.1 \times 10^{-10}$ | CL=90% | 225 |
| $e^+ e^-$ | $S1 \quad < 9 \times 10^{-9}$ | CL=90% | 249 |
| $\pi^0 e^+ e^-$ | $S1 \quad [o] \quad (3.0 \pm 1.5) \times 10^{-9}$ | | 230 |

| | | | |
|---------------------|-----------|--------------------------------|-----|
| $\pi^0 \mu^+ \mu^-$ | <i>s1</i> | $(2.9 \pm 1.5) \times 10^{-9}$ | 177 |
|---------------------|-----------|--------------------------------|-----|



$$I(J^P) = \frac{1}{2}(0^-)$$

$$\begin{aligned} m_{K_L} - m_{K_S} &= (0.5293 \pm 0.0009) \times 10^{10} \text{ } \hbar \text{ s}^{-1} \quad (S = 1.3) \quad \text{Assuming CPT} \\ &= (3.484 \pm 0.006) \times 10^{-12} \text{ MeV} \quad \text{Assuming CPT} \\ &= (0.5289 \pm 0.0010) \times 10^{10} \text{ } \hbar \text{ s}^{-1} \quad \text{Not assuming CPT} \\ \text{Mean life } \tau &= (5.116 \pm 0.021) \times 10^{-8} \text{ s} \quad (S = 1.1) \\ c\tau &= 15.34 \text{ m} \end{aligned}$$

Slope parameters [c]

(See Particle Listings for other linear and quadratic coefficients)

$$\begin{aligned} K_L^0 \rightarrow \pi^+ \pi^- \pi^0: g &= 0.678 \pm 0.008 \quad (S = 1.5) \\ K_L^0 \rightarrow \pi^+ \pi^- \pi^0: h &= 0.076 \pm 0.006 \\ K_L^0 \rightarrow \pi^+ \pi^- \pi^0: k &= 0.0099 \pm 0.0015 \\ K_L^0 \rightarrow \pi^0 \pi^0 \pi^0: h &= (0.6 \pm 1.2) \times 10^{-3} \end{aligned}$$

K_L decay form factors [e]

Linear parametrization assuming μ -e universality

$$\begin{aligned} \lambda_+(K_{\mu 3}^0) &= \lambda_+(K_{e3}^0) = (2.82 \pm 0.04) \times 10^{-2} \quad (S = 1.1) \\ \lambda_0(K_{\mu 3}^0) &= (1.38 \pm 0.18) \times 10^{-2} \quad (S = 2.2) \end{aligned}$$

Quadratic parametrization assuming μ -e universality

$$\begin{aligned} \lambda'_+(K_{\mu 3}^0) &= \lambda'_+(K_{e3}^0) = (2.40 \pm 0.12) \times 10^{-2} \quad (S = 1.2) \\ \lambda''_+(K_{\mu 3}^0) &= \lambda''_+(K_{e3}^0) = (0.20 \pm 0.05) \times 10^{-2} \quad (S = 1.2) \\ \lambda_0(K_{\mu 3}^0) &= (1.16 \pm 0.09) \times 10^{-2} \quad (S = 1.2) \end{aligned}$$

Pole parametrization assuming μ -e universality

$$\begin{aligned} M_V^\mu (K_{\mu 3}^0) &= M_V^e (K_{e3}^0) = 878 \pm 6 \text{ MeV} \quad (S = 1.1) \\ M_S^\mu (K_{\mu 3}^0) &= 1252 \pm 90 \text{ MeV} \quad (S = 2.6) \end{aligned}$$

Dispersive parametrization assuming μ -e universality

$$\begin{aligned} \Lambda_+ &= (2.51 \pm 0.06) \times 10^{-2} \quad (S = 1.5) \\ \ln(C) &= (1.75 \pm 0.18) \times 10^{-1} \quad (S = 2.0) \\ K_{e3}^0 \quad |f_S/f_+| &= (1.5^{+1.4}_{-1.6}) \times 10^{-2} \\ K_{e3}^0 \quad |f_T/f_+| &= (5^{+4}_{-5}) \times 10^{-2} \\ K_{\mu 3}^0 \quad |f_T/f_+| &= (12 \pm 12) \times 10^{-2} \end{aligned}$$

$$\begin{aligned}
 K_L &\rightarrow \ell^+ \ell^- \gamma, K_L \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-: \alpha_{K^*} = -0.205 \pm 0.022 \quad (S = 1.8) \\
 K_L^0 &\rightarrow \ell^+ \ell^- \gamma, K_L^0 \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-: \alpha_{DIP} = -1.69 \pm 0.08 \quad (S = 1.7) \\
 K_L &\rightarrow \pi^+ \pi^- e^+ e^-: a_1/a_2 = -0.737 \pm 0.014 \text{ GeV}^2 \\
 K_L &\rightarrow \pi^0 2\gamma: a_V = -0.43 \pm 0.06 \quad (S = 1.5)
 \end{aligned}$$

***CP*-violation parameters [n]**

$$\begin{aligned}
 A_L &= (0.332 \pm 0.006)\% \\
 |\eta_{00}| &= (2.220 \pm 0.011) \times 10^{-3} \quad (S = 1.8) \\
 |\eta_{+-}| &= (2.232 \pm 0.011) \times 10^{-3} \quad (S = 1.8) \\
 |\epsilon| &= (2.228 \pm 0.011) \times 10^{-3} \quad (S = 1.8) \\
 |\eta_{00}/\eta_{+-}| &= 0.9950 \pm 0.0007 [q] \quad (S = 1.6) \\
 \text{Re}(\epsilon'/\epsilon) &= (1.66 \pm 0.23) \times 10^{-3} [q] \quad (S = 1.6)
 \end{aligned}$$

Assuming *CPT*

$$\begin{aligned}
 \phi_{+-} &= (43.51 \pm 0.05)^\circ \quad (S = 1.2) \\
 \phi_{00} &= (43.52 \pm 0.05)^\circ \quad (S = 1.3) \\
 \phi_\epsilon = \phi_{SW} &= (43.52 \pm 0.05)^\circ \quad (S = 1.2) \\
 \text{Im}(\epsilon'/\epsilon) &= -(\phi_{00} - \phi_{+-})/3 = (-0.002 \pm 0.005)^\circ \quad (S = 1.7)
 \end{aligned}$$

Not assuming *CPT*

$$\begin{aligned}
 \phi_{+-} &= (43.4 \pm 0.5)^\circ \quad (S = 1.2) \\
 \phi_{00} &= (43.7 \pm 0.6)^\circ \quad (S = 1.2) \\
 \phi_\epsilon &= (43.5 \pm 0.5)^\circ \quad (S = 1.3)
 \end{aligned}$$

CP asymmetry A in $K_L^0 \rightarrow \pi^+ \pi^- e^+ e^- = (13.7 \pm 1.5)\%$

β_{CP} from $K_L^0 \rightarrow e^+ e^- e^+ e^- = -0.19 \pm 0.07$

γ_{CP} from $K_L^0 \rightarrow e^+ e^- e^+ e^- = 0.01 \pm 0.11 \quad (S = 1.6)$

j for $K_L^0 \rightarrow \pi^+ \pi^- \pi^0 = 0.0012 \pm 0.0008$

f for $K_L^0 \rightarrow \pi^+ \pi^- \pi^0 = 0.004 \pm 0.006$

$|\eta_{+-\gamma}| = (2.35 \pm 0.07) \times 10^{-3}$

$\phi_{+-\gamma} = (44 \pm 4)^\circ$

$|\epsilon'_{+-\gamma}|/\epsilon < 0.3$, CL = 90%

$|g_{E1}|$ for $K_L^0 \rightarrow \pi^+ \pi^- \gamma < 0.21$, CL = 90%

***T*-violation parameters**

$$\text{Im}(\xi) \text{ in } K_{\mu 3}^0 = -0.007 \pm 0.026$$

***CPT* invariance tests**

$$\phi_{00} - \phi_{+-} = (0.34 \pm 0.32)^\circ$$

$$\text{Re}(\frac{2}{3}\eta_{+-} + \frac{1}{3}\eta_{00}) - \frac{A_L}{2} = (-3 \pm 35) \times 10^{-6}$$

$\Delta S = -\Delta Q$ in K_{e3}^0 decay

$$\text{Re } x = -0.002 \pm 0.006$$

$$\text{Im } x = 0.0012 \pm 0.0021$$

| K_L^0 DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level (MeV/c) | <i>p</i> |
|---|--|---|----------|
| Semileptonic modes | | | |
| $\pi^\pm e^\mp \nu_e$ Called K_{e3}^0 . | [<i>p</i>] $(40.55 \pm 0.11) \%$ | S=1.7 | 229 |
| $\pi^\pm \mu^\mp \nu_\mu$ Called $K_{\mu 3}^0$. | [<i>p</i>] $(27.04 \pm 0.07) \%$ | S=1.1 | 216 |
| $(\pi \mu \text{atom}) \nu$ | $(1.05 \pm 0.11) \times 10^{-7}$ | | 188 |
| $\pi^0 \pi^\pm e^\mp \nu$ | [<i>p</i>] $(5.20 \pm 0.11) \times 10^{-5}$ | | 207 |
| $\pi^\pm e^\mp \nu e^+ e^-$ | [<i>p</i>] $(1.26 \pm 0.04) \times 10^{-5}$ | | 229 |
| Hadronic modes, including Charge conjugation×Parity Violating (<i>CPV</i>) modes | | | |
| $3\pi^0$ | $(19.52 \pm 0.12) \%$ | S=1.6 | 139 |
| $\pi^+ \pi^- \pi^0$ | $(12.54 \pm 0.05) \%$ | | 133 |
| $\pi^+ \pi^-$ | <i>CPV</i> [<i>r</i>] $(1.967 \pm 0.010) \times 10^{-3}$ | S=1.5 | 206 |
| $\pi^0 \pi^0$ | <i>CPV</i> $(8.64 \pm 0.06) \times 10^{-4}$ | S=1.8 | 209 |
| Semileptonic modes with photons | | | |
| $\pi^\pm e^\mp \nu_e \gamma$ | [<i>g,p,s</i>] $(3.79 \pm 0.06) \times 10^{-3}$ | | 229 |
| $\pi^\pm \mu^\mp \nu_\mu \gamma$ | $(5.65 \pm 0.23) \times 10^{-4}$ | | 216 |
| Hadronic modes with photons or $\ell\bar{\ell}$ pairs | | | |
| $\pi^0 \pi^0 \gamma$ | $< 2.43 \times 10^{-7}$ | CL=90% | 209 |
| $\pi^+ \pi^- \gamma$ | [<i>g,s</i>] $(4.15 \pm 0.15) \times 10^{-5}$ | S=2.8 | 206 |
| $\pi^+ \pi^- \gamma$ (DE) | $(2.84 \pm 0.11) \times 10^{-5}$ | S=2.0 | 206 |
| $\pi^0 2\gamma$ | [<i>s</i>] $(1.273 \pm 0.033) \times 10^{-6}$ | | 230 |
| $\pi^0 \gamma e^+ e^-$ | $(1.62 \pm 0.17) \times 10^{-8}$ | | 230 |
| Other modes with photons or $\ell\bar{\ell}$ pairs | | | |
| 2γ | $(5.47 \pm 0.04) \times 10^{-4}$ | S=1.1 | 249 |
| 3γ | $< 7.4 \times 10^{-8}$ | CL=90% | 249 |
| $e^+ e^- \gamma$ | $(9.4 \pm 0.4) \times 10^{-6}$ | S=2.0 | 249 |
| $\mu^+ \mu^- \gamma$ | $(3.59 \pm 0.11) \times 10^{-7}$ | S=1.3 | 225 |
| $e^+ e^- \gamma\gamma$ | [<i>s</i>] $(5.95 \pm 0.33) \times 10^{-7}$ | | 249 |
| $\mu^+ \mu^- \gamma\gamma$ | [<i>s</i>] $(1.0 \pm 0.8) \times 10^{-8}$ | | 225 |

**Charge conjugation \times Parity (CP) or Lepton Family number (LF)
violating modes, or $\Delta S = 1$ weak neutral current ($S1$) modes**

| | | | |
|-------------------------------|----------|--|------------|
| $\mu^+ \mu^-$ | $S1$ | $(6.84 \pm 0.11) \times 10^{-9}$ | 225 |
| $e^+ e^-$ | $S1$ | $(9 \quad {}^{+6}_{-4}) \times 10^{-12}$ | 249 |
| $\pi^+ \pi^- e^+ e^-$ | $S1$ | $[s] \quad (3.11 \pm 0.19) \times 10^{-7}$ | 206 |
| $\pi^0 \pi^0 e^+ e^-$ | $S1$ | $< 6.6 \times 10^{-9}$ | CL=90% 209 |
| $\pi^0 \pi^0 \mu^+ \mu^-$ | $S1$ | $< 9.2 \times 10^{-11}$ | CL=90% 57 |
| $\mu^+ \mu^- e^+ e^-$ | $S1$ | $(2.69 \pm 0.27) \times 10^{-9}$ | 225 |
| $e^+ e^- e^+ e^-$ | $S1$ | $(3.56 \pm 0.21) \times 10^{-8}$ | 249 |
| $\pi^0 \mu^+ \mu^-$ | $CP, S1$ | $[t] < 3.8 \times 10^{-10}$ | CL=90% 177 |
| $\pi^0 e^+ e^-$ | $CP, S1$ | $[t] < 2.8 \times 10^{-10}$ | CL=90% 230 |
| $\pi^0 \nu \bar{\nu}$ | $CP, S1$ | $[u] < 3.0 \times 10^{-9}$ | CL=90% 230 |
| $\pi^0 \pi^0 \nu \bar{\nu}$ | $S1$ | $< 8.1 \times 10^{-7}$ | CL=90% 209 |
| $e^\pm \mu^\mp$ | LF | $[p] < 4.7 \times 10^{-12}$ | CL=90% 238 |
| $e^\pm e^\pm \mu^\mp \mu^\mp$ | LF | $[p] < 4.12 \times 10^{-11}$ | CL=90% 225 |
| $\pi^0 \mu^\pm e^\mp$ | LF | $[p] < 7.6 \times 10^{-11}$ | CL=90% 217 |
| $\pi^0 \pi^0 \mu^\pm e^\mp$ | LF | $< 1.7 \times 10^{-10}$ | CL=90% 159 |

Lorentz invariance violating modes

| | | | |
|----------------|------------------------|--------|-----|
| $\pi^0 \gamma$ | $< 1.7 \times 10^{-7}$ | CL=90% | 230 |
|----------------|------------------------|--------|-----|

$K_0^*(700)$

$$I(J^P) = \frac{1}{2}(0^+)$$

also known as κ ; was $K_0^*(800)$

See the review on "Scalar Mesons below 1 GeV."

Mass (T-Matrix Pole \sqrt{s}) = $(630\text{--}730) - i(260\text{--}340)$ MeV

Mass (Breit-Wigner) = 845 ± 17 MeV

Full width (Breit-Wigner) = 468 ± 30 MeV

| $K_0^*(700)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|--|--------------------------------|-------------|
| $K\pi$ | 100 % | 256 |

$K^*(892)$

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass (T-Matrix Pole \sqrt{s}) = $(890 \pm 14) - i(26 \pm 6)$ MeV

$K^*(892)^\pm$ hadroproduced mass $m = 891.67 \pm 0.26$ MeV

$K^*(892)^\pm$ in τ decays mass $m = 895.5 \pm 0.8$ MeV

$K^*(892)^0$ mass $m = 895.55 \pm 0.20$ MeV (S = 1.7)

$K^*(892)^\pm$ hadroproduced full width $\Gamma = 51.4 \pm 0.8$ MeV

$K^*(892)^\pm$ in τ decays full width $\Gamma = 46.2 \pm 1.3$ MeV

$K^*(892)^0$ full width $\Gamma = 47.3 \pm 0.5$ MeV (S = 1.9)

| $K^*(892)$ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | (MeV/c) p |
|--|----------------------------------|------------------|----------------|
| $K\pi$ | ~ 100 | % | 289 |
| $K^0\gamma$ | $(2.46 \pm 0.21) \times 10^{-3}$ | | 307 |
| $K^\pm\gamma$ | $(9.8 \pm 0.9) \times 10^{-4}$ | | 309 |
| $K\pi\pi$ | $< 7 \times 10^{-4}$ | 95% | 223 |

$K_1(1270)$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 1253 \pm 7$ MeV ($S = 2.2$)

Full width $\Gamma = 90 \pm 20$ MeV [ν]

| $K_1(1270)$ DECAY MODES | Fraction (Γ_i/Γ) | Scale factor | (MeV/c) p |
|---|--------------------------------|--------------|----------------|
| $K\rho$ | $(38 \pm 13) \%$ | 2.2 | † |
| $K_0^*(1430)\pi$ | $(28 \pm 4) \%$ | | † |
| $K^*(892)\pi$ | $(21 \pm 10) \%$ | 2.2 | 286 |
| $K\omega$ | $(11.0 \pm 2.0) \%$ | | † |
| $Kf_0(1370)$ | $(3.0 \pm 2.0) \%$ | | † |
| γK^0 | seen | | 528 |

$K_1(1400)$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 1403 \pm 7$ MeV

Full width $\Gamma = 174 \pm 13$ MeV ($S = 1.6$)

| $K_1(1400)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $K^*(892)\pi$ | $(94 \pm 6) \%$ | 402 |
| $K\rho$ | $(3.0 \pm 3.0) \%$ | 293 |
| $Kf_0(1370)$ | $(2.0 \pm 2.0) \%$ | † |
| $K\omega$ | $(1.0 \pm 1.0) \%$ | 284 |
| $K_0^*(1430)\pi$ | not seen | † |
| γK^0 | seen | 613 |
| $K\phi$ | seen | † |

$K^*(1410)$

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass $m = 1414 \pm 15$ MeV ($S = 1.3$)

Full width $\Gamma = 232 \pm 21$ MeV ($S = 1.1$)

| $K^*(1410)$ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | p (MeV/c) |
|---|--------------------------------|------------------|-------------|
| $K^*(892)\pi$ | > 40 % | 95% | 410 |
| $K\pi$ | (6.6 ± 1.3) % | | 612 |
| $K\rho$ | < 7 % | 95% | 305 |
| γK^0 | $< 2.3 \times 10^{-4}$ | 90% | 619 |
| $K\phi$ | seen | | † |

$K_0^*(1430)$

$$I(J^P) = \frac{1}{2}(0^+)$$

Mass $m = 1425 \pm 50$ MeV [v]

Full width $\Gamma = 270 \pm 80$ MeV [v]

| $K_0^*(1430)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $K\pi$ | (93 ± 10) % | 619 |
| $K\eta$ | ($8.6^{+2.7}_{-3.4}$) % | 486 |
| $K\eta'(958)$ | seen | † |

$K_2^*(1430)$

$$I(J^P) = \frac{1}{2}(2^+)$$

$K_2^*(1430)^{\pm}$ mass $m = 1427.3 \pm 1.5$ MeV (S = 1.3)

$K_2^*(1430)^0$ mass $m = 1432.4 \pm 1.3$ MeV

$K_2^*(1430)^{\pm}$ full width $\Gamma = 100.0 \pm 2.1$ MeV

$K_2^*(1430)^0$ full width $\Gamma = 109 \pm 5$ MeV (S = 1.9)

| $K_2^*(1430)$ DECAY MODES | Fraction (Γ_i/Γ) | Scale factor/ Confidence level | p (MeV/c) |
|---|--|-----------------------------------|-------------|
| $K\pi$ | (49.9 ± 1.2) % | | 620 |
| $K^*(892)\pi$ | (24.7 ± 1.5) % | | 420 |
| $K^*(892)\pi\pi$ | (13.4 ± 2.2) % | | 373 |
| $K\rho$ | (8.7 ± 0.8) % | S=1.2 | 320 |
| $K\omega$ | (2.9 ± 0.8) % | | 313 |
| $K^+\gamma$ | (2.4 ± 0.5) $\times 10^{-3}$ | S=1.1 | 628 |
| $K\eta$ | ($1.5^{+3.4}_{-1.0}$) $\times 10^{-3}$ | S=1.3 | 488 |
| $K\omega\pi$ | $< 7.2 \times 10^{-4}$ | CL=95% | 106 |
| $K^0\gamma$ | $< 9 \times 10^{-4}$ | CL=90% | 627 |

K(1460)

$$I(J^P) = \frac{1}{2}(0^-)$$

| K(1460) DECAY MODES | Fraction (Γ_i/Γ) | <i>p</i> (MeV/c) |
|----------------------------|--------------------------------|------------------|
| $K^*(892)\pi$ | seen | — |
| $K\rho$ | seen | — |
| $K_0^*(1430)\pi$ | seen | — |
| $K\phi$ | seen | — |

K₁(1650)

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 1650 \pm 50$ MeV
 Full width $\Gamma = 150 \pm 50$ MeV

K^{*}(1680)

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass $m = 1718 \pm 18$ MeV
 Full width $\Gamma = 322 \pm 110$ MeV ($S = 4.2$)

| K[*](1680) DECAY MODES | Fraction (Γ_i/Γ) | <i>p</i> (MeV/c) |
|--|--------------------------------|------------------|
| $K\pi$ | (38.7 \pm 2.5) % | 782 |
| $K\rho$ | (31.4 \pm 5.0) % | 571 |
| $K^*(892)\pi$ | (29.9 \pm 2.2) % | 618 |
| $K\phi$ | seen | 387 |
| $K\eta$ | (1.4 \pm 1.0) % | 683 |

K₂(1770) [x]

$$I(J^P) = \frac{1}{2}(2^-)$$

Mass $m = 1773 \pm 8$ MeV
 Full width $\Gamma = 186 \pm 14$ MeV

| K₂(1770) DECAY MODES | Fraction (Γ_i/Γ) | <i>p</i> (MeV/c) |
|--|--------------------------------|------------------|
| $K\pi\pi$ | | 794 |
| $K_2^*(1430)\pi$ | seen | 287 |
| $K^*(892)\pi$ | seen | 654 |
| $Kf_2(1270)$ | seen | 53 |
| $K\phi$ | seen | 441 |
| $K\omega$ | seen | 607 |

$K_3^*(1780)$

$$I(J^P) = \frac{1}{2}(3^-)$$

Mass $m = 1779 \pm 8$ MeV ($S = 1.2$)
 Full width $\Gamma = 161 \pm 17$ MeV ($S = 1.1$)

| $K_3^*(1780)$ DECAY MODES | Fraction (Γ_i/Γ) | Confidence level | p (MeV/c) |
|---|--------------------------------|------------------|-------------|
| $K\rho$ | (31 \pm 9) % | | 616 |
| $K^*(892)\pi$ | (20 \pm 5) % | | 657 |
| $K\pi$ | (18.8 \pm 1.0) % | | 815 |
| $K\eta$ | (30 \pm 13) % | | 721 |
| $K_2^*(1430)\pi$ | < 16 % | 95% | 292 |

$K_2(1820)$ [x]

$$I(J^P) = \frac{1}{2}(2^-)$$

Mass $m = 1819 \pm 12$ MeV
 Full width $\Gamma = 264 \pm 34$ MeV

| $K_2(1820)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $K\pi\pi$ | seen | 819 |
| $K_2^*(1430)\pi$ | seen | 328 |
| $K^*(892)\pi$ | seen | 683 |
| $Kf_2(1270)$ | seen | 191 |
| $K\omega$ | seen | 640 |
| $K\phi$ | seen | 483 |

$K_2^*(1980)$

$$I(J^P) = \frac{1}{2}(2^+)$$

Mass $m = 1994^{+60}_{-50}$ MeV ($S = 2.8$)
 Full width $\Gamma = 348^{+50}_{-30}$ MeV ($S = 1.3$)

| $K_2^*(1980)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $K^*(892)\pi$ | possibly seen | 791 |
| $K\rho$ | possibly seen | 762 |
| $Kf_2(1270)$ | possibly seen | 424 |
| $K\phi$ | seen | 627 |
| $K\eta$ | seen | 850 |

$K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

Mass $m = 2048^{+8}_{-9}$ MeV ($S = 1.1$)Full width $\Gamma = 199^{+27}_{-19}$ MeV

| $K_4^*(2045)$ DECAY MODES | Fraction (Γ_i/Γ) | p (MeV/c) |
|---|--------------------------------|-------------|
| $K\pi$ | (9.9±1.2) % | 960 |
| $K^*(892)\pi\pi$ | (9 ± 5) % | 804 |
| $K^*(892)\pi\pi\pi$ | (7 ± 5) % | 770 |
| $\rho K\pi$ | (5.7±3.2) % | 744 |
| $\omega K\pi$ | (5.0±3.0) % | 740 |
| $\phi K\pi$ | (2.8±1.4) % | 597 |
| $\phi K^*(892)$ | (1.4±0.7) % | 368 |

NOTES

[a] See the note in the K^\pm Particle Listings.[b] Neglecting photon channels. See, e.g., A. Pais and S.B. Treiman, Phys. Rev. **D12**, 2744 (1975).[c] The definition of the slope parameters of the $K \rightarrow 3\pi$ Dalitz plot is as follows (see also “Note on Dalitz Plot Parameters for $K \rightarrow 3\pi$ Decays” in the K^\pm Particle Listings):

$$|M|^2 = 1 + g(s_3 - s_0)/m_{\pi^+}^2 + \dots$$

[d] See the review on “Form Factors for Radiative Pion and Kaon Decays” for definitions and details.

[e] For more details and definitions of parameters see the Particle Listings.

[f] See the K^\pm Particle Listings for the energy limits used in this measurement.[g] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ ’s.

[h] Structure-dependent part.

[i] Direct-emission branching fraction.

[j] Derived from an analysis of neutrino-oscillation experiments.

[k] Violates angular-momentum conservation.

[l] Derived from measured values of ϕ_{+-} , ϕ_{00} , $|\eta|$, $|m_{K_L^0} - m_{K_S^0}|$, and $\tau_{K_S^0}$, as described in the introduction to “Tests of Conservation Laws.”[n] The CP -violation parameters are defined as follows (see also “Note on CP Violation in $K_S \rightarrow 3\pi$ ” and “Note on CP Violation in K_L^0 Decay” in the Particle Listings):

$$\eta_{+-} = |\eta_{+-}| e^{i\phi_{+-}} = \frac{A(K_L^0 \rightarrow \pi^+ \pi^-)}{A(K_S^0 \rightarrow \pi^+ \pi^-)} = \epsilon + \epsilon'$$

$$\eta_{00} = |\eta_{00}| e^{i\phi_{00}} = \frac{A(K_L^0 \rightarrow \pi^0 \pi^0)}{A(K_S^0 \rightarrow \pi^0 \pi^0)} = \epsilon - 2\epsilon'$$

$$\delta = \frac{\Gamma(K_L^0 \rightarrow \pi^- \ell^+ \nu) - \Gamma(K_L^0 \rightarrow \pi^+ \ell^- \nu)}{\Gamma(K_L^0 \rightarrow \pi^- \ell^+ \nu) + \Gamma(K_L^0 \rightarrow \pi^+ \ell^- \nu)},$$

$$\text{Im}(\eta_{+-0})^2 = \frac{\Gamma(K_S^0 \rightarrow \pi^+ \pi^- \pi^0)^{\text{CP viol.}}}{\Gamma(K_L^0 \rightarrow \pi^+ \pi^- \pi^0)},$$

$$\text{Im}(\eta_{000})^2 = \frac{\Gamma(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)}{\Gamma(K_L^0 \rightarrow \pi^0 \pi^0 \pi^0)}.$$

where for the last two relations *CPT* is assumed valid, *i.e.*, $\text{Re}(\eta_{+-0}) \simeq 0$ and $\text{Re}(\eta_{000}) \simeq 0$.

- [o] See the K_S^0 Particle Listings for the energy limits used in this measurement.
- [p] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [q] $\text{Re}(\epsilon'/\epsilon) = \epsilon'/\epsilon$ to a very good approximation provided the phases satisfy *CPT* invariance.
- [r] This mode includes gammas from inner bremsstrahlung but not the direct emission mode $K_L^0 \rightarrow \pi^+ \pi^- \gamma$ (DE).
- [s] See the K_L^0 Particle Listings for the energy limits used in this measurement.
- [t] Allowed by higher-order electroweak interactions.
- [u] Violates *CP* in leading order. Test of direct *CP* violation since the indirect *CP*-violating and *CP*-conserving contributions are expected to be suppressed.
- [v] Our estimate. See the Particle Listings for details.
- [x] See our minireview under the $K_2(1770)$ in the 2004 edition of this *Review*.