

86. W' -Boson Searches

Revised September 10, 2023 by B.A. Dobrescu (FNAL) and S. Willocq (U. Massachusetts).

The W' boson is a massive hypothetical particle of spin 1 and electric charge ± 1 , which is a color singlet and is predicted in various extensions of the Standard Model (SM).

86.1 W' couplings to quarks and leptons

The Lagrangian terms describing the couplings of a W'^+ boson to fermions are given by

$$\frac{W'_\mu^{i+}}{\sqrt{2}} \left[\bar{u}_i \left(C_{q_{ij}}^R P_R + C_{q_{ij}}^L P_L \right) \gamma^\mu d_j + \bar{\nu}_i \left(C_{\ell_{ij}}^R P_R + C_{\ell_{ij}}^L P_L \right) \gamma^\mu e_j \right]. \quad (86.1)$$

Here, u, d, ν , and e are the SM fermions in the mass eigenstate basis, $i, j = 1, 2, 3$ label the fermion generation, and $P_{R,L} = (1 \pm \gamma_5)/2$. The coefficients $C_{q_{ij}}^L$, $C_{q_{ij}}^R$, $C_{\ell_{ij}}^L$, and $C_{\ell_{ij}}^R$ are complex dimensionless parameters. If $C_{\ell_{ij}}^R \neq 0$, then the i th generation includes a right-handed neutrino. Using this notation, the SM W couplings are $C_q^L = g V_{CKM}$, $C_\ell^L = g \approx 0.63$ and $C_q^R = C_\ell^R = 0$.

Unitarity considerations imply that the W' boson is associated with a spontaneously-broken gauge symmetry. This is true even when it is a composite particle (*e.g.* ρ^\pm -like bound states [1]) if its mass is much smaller than the compositeness scale, or a Kaluza-Klein mode in theories where the W boson propagates in extra dimensions [2]. The simplest extension of the electroweak gauge group that includes a W' boson is $SU(2)_1 \times SU(2)_2 \times U(1)$, but larger groups are encountered in some theories. A generic property of these gauge theories is that they also include a Z' boson [3]; the W' -to- Z' mass ratio is often a free parameter.

A tree-level mass mixing may be induced between the electrically-charged gauge bosons. Upon diagonalization of their mass matrix, the W -to- Z mass ratio and the couplings of the observed W boson are shifted from the SM values. Their measurements imply that the mixing angle, θ_+ , between the gauge eigenstates must be smaller than about 10^{-2} [4]. In certain theories the mixing is negligible (*e.g.*, due to a new parity [5]), even when the W' mass is near the electroweak scale. Note that $SU(2)$ gauge invariance suppresses the kinetic mixing between the W and W' bosons (in contrast to the case of a Z' boson [3]).

The W' coupling to WZ is fixed by Lorentz and gauge invariances, and to leading order in θ_+ is given by [6]

$$\frac{g \theta_+ i}{\cos \theta_W} \left[W_\mu'^+ (W_\nu^- Z^{\nu\mu} + Z_\nu W^{-\mu\nu}) + Z^\nu W^{-\mu} W_{\nu\mu}'^+ \right] + \text{H.c.}, \quad (86.2)$$

where $W^{\mu\nu} \equiv \partial^\mu W^\nu - \partial^\nu W^\mu$, etc. The θ_W dependence shown here corrects the one given in Ref. [7], which has been referred to as the Extended Gauge Model by the experimental collaborations. The W' coupling to Wh^0 , where h^0 is the SM Higgs boson, is

$$-\xi_h g_{W'} M_W W_\mu'^+ W^{\mu-} h^0 + \text{H.c.}, \quad (86.3)$$

where $g_{W'}$ is the gauge coupling of the W' boson, and the coefficient ξ_h satisfies $\xi_h \leq 1$ in simple Higgs sectors [6].

In models based on the “left-right symmetric” gauge group [8], $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$, the SM fermions that couple to the W boson transform as doublets under $SU(2)_L$ while the other fermions transform as doublets under $SU(2)_R$. Consequently, the W' boson couples primarily to right-handed fermions; its coupling to left-handed fermions arises due to the θ_+ mixing, so that C_q^L is proportional to the CKM matrix and its elements are much smaller than the diagonal elements of C_q^R . Generically, C_q^R does not need to be proportional to V_{CKM} .

There are many other models based on the $SU(2)_1 \times SU(2)_2 \times U(1)$ gauge symmetry. In the “alternate left-right” model [9], all the couplings shown in Eq. (86.1) vanish, but there are some new fermions such that the W' boson couples to pairs involving a SM fermion and a new fermion. In the “ununified SM” [10], the left-handed quarks are doublets under one $SU(2)$, and the left-handed leptons are doublets under a different $SU(2)$, leading to a mostly leptophobic W' boson: $C_{\ell_{ij}}^L \ll C_{q_{ij}}^L$ and $C_{\ell_{ij}}^R = C_{q_{ij}}^R = 0$. Fermions of different generations may also transform as doublets under different $SU(2)$ gauge groups [11]. In particular, the couplings to third generation quarks may be enhanced [12].

It is also possible that the W' couplings to SM fermions are highly suppressed. For example, if the quarks and leptons are singlets under one $SU(2)$ [13], then the couplings are proportional to the tiny mixing angle θ_+ . Similar suppressions may arise if some vectorlike fermions mix with the SM fermions [14].

Gauge groups that embed the electroweak symmetry, such as $SU(3)_W \times U(1)$ or $SU(4)_W \times U(1)$, also include one or more W' bosons [15].

86.2 Collider searches

At hadron colliders, W' bosons can be detected through resonant pair production of fermions (f and f') or electroweak bosons with a net electric charge equal to ± 1 . When W' has a width much smaller than its mass ($\Gamma_{W'}/M_{W'} \lesssim 7\%$), the contribution of the s -channel W' exchange to the total rate for $pp \rightarrow f\bar{f}'X$, where X is any final state, may be approximated by the branching fraction $B(W' \rightarrow f\bar{f}')$ times the production cross section [16], which may be written as

$$\sigma(pp \rightarrow W'X) \simeq \frac{\pi}{6s} \sum_{i,j} [(C_{q_{ij}}^L)^2 + (C_{q_{ij}}^R)^2] w_{ij}(M_{W'}^2/s, M_{W'}) . \quad (86.4)$$

The functions w_{ij} include the information about proton structure, and are given to leading order in α_s by

$$w_{ij}(z, \mu) = \int_z^1 \frac{dx}{x} \left[u_i(x, \mu) \bar{d}_j \left(\frac{z}{x}, \mu \right) + \bar{u}_i(x, \mu) d_j \left(\frac{z}{x}, \mu \right) \right], \quad (86.5)$$

where $u_i(x, \mu)$ and $d_i(x, \mu)$ are the parton distributions inside the proton at the factorization scale μ and parton momentum fraction x for the up- and down-type quarks of the i th generation, respectively. QCD corrections to W' production are sizable (they also include quark-gluon initial states), but preserve the above factorization of couplings at next-to-leading order [17].

The most commonly studied W' signal consists of a high-momentum electron or muon and large missing transverse momentum. The signal transverse mass distribution forms a Jacobian peak with its endpoint at $M_{W'}$ (see Fig. 1 (top) of Ref. [18]). Given that the branching fractions for $W' \rightarrow e\nu$ and $W' \rightarrow \mu\nu$ could be very different, the results in these channels should be presented separately. Searches in these channels often implicitly assume that the left-handed couplings vanish (no interference between W and W'), and that the right-handed neutrino is light compared to the W' boson and escapes the detector. An example of parameter values that satisfy these assumptions is $C_q^R = gV_{CKM}$, $C_\ell^R = g$, $C_q^L = C_\ell^L = 0$, which define a model that preserves lepton universality and predicts the same total cross section as the Sequential SM used in many W' searches. However, if a W' boson were discovered and the final state fermions have left-handed helicity, then the effects of $W - W'$ interference could be observed [19], providing information about the W' couplings. The effects of the W' width on interference are discussed in Ref. [20].

In the $e\nu$ channel, the ATLAS and CMS collaborations set limits on the W' production cross section times branching fraction $\sigma \times B$ (and thus indirectly on the W' couplings). These limits are set for $M_{W'}$ in the $0.15 - 7$ TeV range and are based on approximately 140 fb^{-1} at $\sqrt{s} = 13$ TeV [18, 21], with the most stringent limits reproduced in Fig. 86.1. ATLAS sets the strongest mass

limit $M_{W'} > 6.0$ TeV in the Sequential SM (all limits in this mini-review are at the 95% CL). The coupling limits are much weaker for $M_{W'} < 150$ GeV, a range last explored with the Tevatron at $\sqrt{s} = 1.8$ TeV [22].

In the $\mu\nu$ channel, ATLAS and CMS set rate limits for $M_{W'}$ in the 0.15 – 7 TeV range [18, 21], with the strongest mass lower limit of 5.6 TeV in the Sequential SM set by CMS [21] using 138 fb^{-1} of $\sqrt{s} = 13$ TeV data, as shown in Fig. 86.1. When combined with the $e\nu$ channel assuming lepton universality, the upper limit on the $\sqrt{s} = 13$ TeV cross section times branching fraction to $\ell\nu$ varies between 0.05 and 2.1 fb for $M_{W'}$ values between 1 and 6 TeV [18]. Only weak limits on $W' \rightarrow \mu\nu$ exist for $M_{W'} < 150$ GeV [23]. Note that masses of the order of the electroweak scale are interesting from a theoretical point of view, while lepton universality does not necessarily apply to a W' boson.




Figure 86.1: Upper limit on $\sigma(pp \rightarrow W'X) \times B(W' \rightarrow \ell\nu)$ in the $e\nu$ channel from ATLAS [18] (left) and the $\mu\nu$ channel from CMS [21] (right). The red (black) line shows the theoretical prediction in the Sequential SM in the $e\nu$ ($\mu\nu$) channel.

Searches for $W' \rightarrow \tau\nu$ have been performed at 13 TeV by CMS with 138 fb^{-1} [24], and by ATLAS with 139 fb^{-1} [25]. Limits are set on $\sigma \times B$ for $M_{W'}$ between 0.5 and 6 TeV. A mass lower limit of 5.0 TeV is set in the Sequential SM, and the upper limit on the cross section times branching fraction to $\tau\nu$ at 13 TeV varies between 0.4 and 9 fb for $M_{W'}$ values between 1 and 5 TeV [25].

The W' decay into a charged lepton and a right-handed neutrino, ν_R , may also be followed by the ν_R decay through a virtual W' boson into a charged lepton and two quark jets. The CMS [26] and ATLAS [27] searches in the $eejj$ and $\mu\mu jj$ channels have set limits at $\sqrt{s} = 13$ TeV on the cross section times branching fractions as a function of the ν_R mass and of $M_{W'}$. No requirement is placed on the charge of the lepton pair. A related W' search in the $\tau\tau jj$ channel with hadronic τ decays was also performed by CMS [28].

The $t\bar{b}$ channel is particularly important because a W' boson that couples only to right-handed fermions cannot decay to leptons when the right-handed neutrinos are heavier than $M_{W'}$. Additional motivations are provided by a W' boson with enhanced couplings to the third generation [12], and by a leptophobic W' boson. The usual signature for $W' \rightarrow t\bar{b}$ consists of a leptonically-decaying W boson and two b -jets. Recent studies have also incorporated the fully hadronic decay channel for $M_{W'} \gg m_t$ with the use of jet substructure techniques to tag highly boosted top-jets. For a detailed discussion of this channel, see Ref. [29].

Searches for dijet resonances may be used to set limits on $W' \rightarrow q\bar{q}'$. ATLAS [30] and CMS [31] provide similar coverage in the ~ 1.5 – 8 TeV mass range with 139 and 137 fb^{-1} of data, respectively, collected at $\sqrt{s} = 13$ TeV. Interpretation in terms of W' decays with 139 fb^{-1} of 13 TeV data yields

86. W' -Boson Searches

a W' mass lower limit of 4.0 TeV in the Sequential SM [30]. For masses in the range $\sim 0.5\text{--}1.5$ TeV, analyses based on jets reconstructed online provide the best sensitivity because they circumvent trigger bandwidth limitations [32, 33]. For W' masses below ~ 0.5 TeV, the best limits are set in novel analyses exploiting boosted technologies and initial state radiation [34–37]. Cross-section limits for W' masses below ~ 1.5 TeV can be derived from the dijet limits on Z' bosons summarized in Ref. [3].

In some theories [5] the W' couplings to SM fermions are suppressed by discrete symmetries. W' production then occurs in pairs, through a photon or Z boson. The decay modes are model-dependent and often involve other new particles. The ensuing collider signals arise from cascade decays and often include missing transverse momentum or boosted multi-jet topologies. Even when single production of W' is large enough, it is possible that its main decay mode is into a new particle and a SM one. An example is a search performed by CMS [38] for W' decays into a vector-like quark and a top or a bottom quark. The final state studied in this analysis involves a boosted SM Higgs or Z boson, as well as a $t\bar{t}$ or $b\bar{b}$ pair, with all heavy particles decaying into jets. Another example is a search performed by ATLAS [39] for W' decays into a W boson and a Z' boson, the latter decaying into a jet pair.

Searches for WZ resonances at the LHC have focused on the process $pp \rightarrow W' \rightarrow WZ$ with the production mainly from $u\bar{d} \rightarrow W'$, assuming SM-like couplings to quarks. ATLAS and CMS have set upper limits on the $W'WZ$ coupling for $M_{W'}$ in the $0.2\text{--}5.0$ TeV range with a combination of fully leptonic, semi-leptonic and fully hadronic channels with $\sim 36\text{--}139\text{ fb}^{-1}$ at 13 TeV [40–42] (see also Ref. [29]). Recent constraints with $138\text{--}139\text{ fb}^{-1}$ at 13 TeV were also set in individual channels by ATLAS [43–45] and CMS [46–49]. The strongest lower limit on the W' mass is set by ATLAS [44] in the semi-leptonic channel with a lower limit on $M_{W'}$ of 3.9 TeV [44] in the context of the Heavy Vector Triplet (HVT) weakly-coupled scenario A [50]. A similar result is obtained by CMS in the fully-hadronic channel [46]. A fermiophobic W' boson that couples to WZ may be produced at hadron colliders in association with a Z boson, or via WZ fusion. This would give rise to $(WZ)Z$ and $(WZ)jj$ final states [51]. Results of the search for the latter are reported in Refs. [40, 44, 46, 47, 49].




Figure 86.2: Upper limits on W' production cross section times branching fraction into a W and a SM Higgs boson decaying into heavy-flavor quarks, from ATLAS [52] (left) and CMS [46] (right).

W' bosons have also been searched for in final states with a W boson and a SM Higgs boson in the channels $W \rightarrow \ell\nu$ or $W \rightarrow q\bar{q}'$ and $h^0 \rightarrow b\bar{b}$ by ATLAS [52, 53] and CMS [46, 49] with $138\text{--}139\text{ fb}^{-1}$ at $\sqrt{s} = 13$ TeV. CMS also searched for W' bosons in the $q\bar{q}'\tau\tau$ final state [54]. Cross-section limits are set for W' masses between 0.4 and 6.0 TeV. The most stringent upper limit on the cross

section is set by the ATLAS analysis with $W \rightarrow \ell\nu$ at low $M_{W'}$ and the CMS analysis with $W \rightarrow q\bar{q}'$ at high $M_{W'}$, as shown in Fig. 86.2.

At lepton colliders, W' bosons may be produced in pairs via their photon coupling, which is model independent. At LEP-II, although dedicated searches for W' bosons have not been performed, the large cross section for $e^+e^- \rightarrow \gamma^* \rightarrow W'^+W'^-$ and small backgrounds suggest that any W' is ruled out up to the kinematic limit, $M_{W'} < \sqrt{s}/2 \approx 105$ GeV, for most decay modes. Sensitivity to $M_{W'}$ above \sqrt{s} could be achieved [55] using the $e^+e^- \rightarrow \gamma\nu\bar{\nu}$ process via a t -channel W' exchange, if the W' coupling to $e\nu$ is large enough.

86.3 Low-energy constraints

The properties of W' bosons are also constrained by measurements of processes at energies much below $M_{W'}$. The bounds on $W - W'$ mixing [4] are mostly due to the change in W properties compared to the SM. Limits on deviations in the ZWW couplings provide a leading constraint for fermiophobic W' bosons [14].

Constraints arising from low-energy effects of W' exchange are strongly model-dependent. If the W' couplings to quarks are not suppressed, then box diagrams involving a W and a W' boson contribute to neutral meson-mixing. In the case of W' couplings to right-handed quarks as in the left-right symmetric model, the limit from $K_L - K_S$ mixing is severe: $M_{W'} > 2.9$ TeV for $C_q^R = gV_{CKM}$ [56]. However, if no correlation between the W' and W couplings is assumed, then the limit on $M_{W'}$ may be significantly relaxed [57].

W' exchange also contributes at tree level to various low-energy processes. In particular, it would impact the measurement of the Fermi constant G_F in muon decay, which in turn would change the predictions of many other electroweak processes. A recent test of parity violation in polarized muon decay [58] has set limits of about 600 GeV on $M_{W'}$, assuming W' couplings to right-handed leptons as in left-right symmetric models and a light ν_R . There are also W' contributions to the neutron electric dipole moment, β decays, and other processes [4].

If right-handed neutrinos have Majorana masses, then there are tree-level contributions to neutrinoless double-beta decay, and a limit on $M_{W'}$ versus the ν_R mass may be derived [59]. For ν_R masses below a few GeV, the W' boson contributes to leptonic and semileptonic B meson decays, so that limits may be placed on various combinations of W' parameters [57]. For ν_R masses below ~ 30 MeV, the most stringent constraints on $M_{W'}$ are due to the limits on ν_R emission from supernovae [60].

References

- [1] M. Bando, T. Kugo and K. Yamawaki, *Phys. Rept.* **164**, 217 (1988).
- [2] H.-C. Cheng *et al.*, *Phys. Rev. D* **64**, 065007 (2001), [[hep-th/0104179](#)].
- [3] See the Section on “ Z' -boson searches” in this *Review*.
- [4] See the particle listings for W' in this *Review*.
- [5] H.-C. Cheng and I. Low, *JHEP* **09**, 051 (2003), [[hep-ph/0308199](#)].
- [6] B. A. Dobrescu and Z. Liu, *JHEP* **10**, 118 (2015), [[arXiv:1507.01923](#)].
- [7] G. Altarelli, B. Mele and M. Ruiz-Altaba, *Z. Phys. C* **45**, 109 (1989), [Erratum: *Z. Phys. C* **47**, 676 (1990)].
- [8] R. N. Mohapatra and J. C. Pati, *Phys. Rev. D* **11**, 566 (1975).
- [9] K. S. Babu, X.-G. He and E. Ma, *Phys. Rev. D* **36**, 878 (1987).
- [10] H. Georgi, E. E. Jenkins and E. H. Simmons, *Nucl. Phys. B* **331**, 541 (1990).
- [11] X.-y. Li and E. Ma, *J. Phys. G* **19**, 1265 (1993), [[hep-ph/9208210](#)].
- [12] D. J. Muller and S. Nandi, *Phys. Lett. B* **383**, 345 (1996), [[hep-ph/9602390](#)].

- [13] A. Donini *et al.*, Nucl. Phys. B **507**, 51 (1997), [[hep-ph/9705450](#)].
- [14] R. S. Chivukula *et al.*, Phys. Rev. D **74**, 075011 (2006), [[hep-ph/0607124](#)].
- [15] F. Pisano and V. Pleitez, Phys. Rev. D **46**, 410 (1992), [[hep-ph/9206242](#)].
- [16] V. D. Barger and R. J. N. Phillips, *Collider Physics* (1996), ISBN 978-0201149456.
- [17] Z. Sullivan, Phys. Rev. D **66**, 075011 (2002), [[hep-ph/0207290](#)].
- [18] G. Aad *et al.* (ATLAS), Phys. Rev. D **100**, 052013 (2019), [[arXiv:1906.05609](#)].
- [19] T. G. Rizzo, JHEP **05**, 037 (2007), [[arXiv:0704.0235](#)].
- [20] E. Accomando *et al.*, Phys. Rev. D **85**, 115017 (2012), [[arXiv:1110.0713](#)].
- [21] A. Tumasyan *et al.* (CMS), JHEP **07**, 067 (2022), [[arXiv:2202.06075](#)].
- [22] F. Abe *et al.* (CDF), Phys. Rev. Lett. **74**, 2900 (1995).
- [23] F. Abe *et al.* (CDF), Phys. Rev. Lett. **67**, 2609 (1991).
- [24] A. Tumasyan *et al.* (CMS), JHEP **09**, 051 (2023), [[arXiv:2212.12604](#)].
- [25] ATLAS Collab., ATLAS-CONF-2021-025, Jun. 2021.
- [26] A. Tumasyan *et al.* (CMS), JHEP **04**, 047 (2022), [[arXiv:2112.03949](#)].
- [27] G. Aad *et al.* (ATLAS) (2023), [[arXiv:2304.09553](#)].
- [28] A. M. Sirunyan *et al.* (CMS), JHEP **03**, 170 (2019), [[arXiv:1811.00806](#)].
- [29] K.M. Black *et al.*, “Dynamical electroweak symmetry breaking” in this *Review*.
- [30] G. Aad *et al.* (ATLAS), JHEP **03**, 145 (2020), [[arXiv:1910.08447](#)].
- [31] A. M. Sirunyan *et al.* (CMS), JHEP **05**, 033 (2020), [[arXiv:1911.03947](#)].
- [32] M. Aaboud *et al.* (ATLAS), Phys. Rev. Lett. **121**, 081801 (2018), [[arXiv:1804.03496](#)].
- [33] A. M. Sirunyan *et al.* (CMS), JHEP **08**, 130 (2018), [[arXiv:1806.00843](#)].
- [34] M. Aaboud *et al.* (ATLAS), Phys. Lett. B **788**, 316 (2019), [[arXiv:1801.08769](#)].
- [35] M. Aaboud *et al.* (ATLAS), Phys. Lett. B **795**, 56 (2019), [[arXiv:1901.10917](#)].
- [36] A. M. Sirunyan *et al.* (CMS), Phys. Rev. D **100**, 112007 (2019), [[arXiv:1909.04114](#)].
- [37] A. M. Sirunyan *et al.* (CMS), Phys. Rev. Lett. **123**, 231803 (2019), [[arXiv:1905.10331](#)].
- [38] A. Tumasyan *et al.* (CMS), JHEP **09**, 088 (2022), [[arXiv:2202.12988](#)].
- [39] G. Aad *et al.* (ATLAS), JHEP **07**, 202 (2023), [[arXiv:2211.08945](#)].
- [40] M. Aaboud *et al.* (ATLAS), Phys. Rev. D **98**, 052008 (2018), [[arXiv:1808.02380](#)].
- [41] ATLAS Collab., ATLAS-CONF-2022-028, May 2022.
- [42] A. M. Sirunyan *et al.* (CMS), Phys. Lett. B **798**, 134952 (2019), [[arXiv:1906.00057](#)].
- [43] G. Aad *et al.* (ATLAS), JHEP **09**, 091 (2019), [Erratum: JHEP **06** (2020) 042], [[arXiv:1906.08589](#)].
- [44] G. Aad *et al.* (ATLAS), Eur. Phys. J. C **80**, 1165 (2020), [[arXiv:2004.14636](#)].
- [45] G. Aad *et al.* (ATLAS), Eur. Phys. J. C **83**, 633 (2023), [[arXiv:2207.03925](#)].
- [46] A. Tumasyan *et al.* (CMS), Phys. Lett. B **844**, 137813 (2023), [[arXiv:2210.00043](#)].
- [47] A. Tumasyan *et al.* (CMS), Phys. Rev. D **106**, 012004 (2022), [[arXiv:2109.08268](#)].
- [48] A. Tumasyan *et al.* (CMS), JHEP **04**, 087 (2022), [[arXiv:2111.13669](#)].
- [49] A. Tumasyan *et al.* (CMS), Phys. Rev. D **105**, 032008 (2022), [[arXiv:2109.06055](#)].
- [50] D. Pappadopulo *et al.*, JHEP **09**, 060 (2014), [[arXiv:1402.4431](#)].

- [51] H.-J. He *et al.*, Phys. Rev. D **78**, 031701 (2008), [[arXiv:0708.2588](#)].
- [52] G. Aad *et al.* (ATLAS), JHEP **06**, 016 (2023), [[arXiv:2207.00230](#)].
- [53] G. Aad *et al.* (ATLAS), Phys. Rev. D **102**, 112008 (2020), [[arXiv:2007.05293](#)].
- [54] A. M. Sirunyan *et al.* (CMS), JHEP **01**, 051 (2019), [[arXiv:1808.01365](#)].
- [55] S. Godfrey *et al.*, Phys. Rev. D **61**, 113009 (2000), [[hep-ph/0001074](#)].
- [56] Y. Zhang *et al.*, Phys. Rev. D **76**, 091301 (2007), [[arXiv:0704.1662](#)].
- [57] P. Langacker and S. U. Sankar, Phys. Rev. D **40**, 1569 (1989).
- [58] J. F. Bueno *et al.* (TWIST), Phys. Rev. D **84**, 032005 (2011), [[arXiv:1104.3632](#)].
- [59] G. Prezeau, M. Ramsey-Musolf and P. Vogel, Phys. Rev. D **68**, 034016 (2003), [[hep-ph/0303205](#)].
- [60] G. G. Raffelt, Ann. Rev. Nucl. Part. Sci. **49**, 163 (1999), [[hep-ph/9903472](#)].