



$I(J^P) = 0(\frac{1}{2}^+)$  Status: \*\*\*

In the quark model, a  $\Lambda_b^0$  is an isospin-0  $ud\bar{b}$  state. The lowest  $\Lambda_b^0$  ought to have  $J^P = 1/2^+$ . None of  $I$ ,  $J$ , or  $P$  have actually been measured.

## $\Lambda_b^0$ MASS

$m_{\Lambda_b^0}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>5619.60 ± 0.17 OUR AVERAGE</b>				
5619.62 ± 0.16 ± 0.13		1 AAIJ	17AM LHCb	$p\bar{p}$ at 7, 8 TeV
5619.30 ± 0.34		2 AAIJ	14AA LHCb	$p\bar{p}$ at 7 TeV
5620.15 ± 0.31 ± 0.47		3 AALTONEN	14B CDF	$p\bar{p}$ at 1.96 TeV
5619.7 ± 0.7 ± 1.1		3 AAD	13U ATLAS	$p\bar{p}$ at 7 TeV
5621 ± 4 ± 3		4 ABE	97B CDF	$p\bar{p}$ at 1.8 TeV
5668 ± 16 ± 8	4	5 ABREU	96N DLPH	$e^+e^- \rightarrow Z$
5614 ± 21 ± 4	4	5 BUSKULIC	96L ALEP	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5619.65 ± 0.17 ± 0.17		6 AAIJ	16Y LHCb	Repl. by AAIJ 17AM
5619.44 ± 0.13 ± 0.38		3 AAIJ	13AV LHCb	Repl. by AAIJ 13AV
5619.19 ± 0.70 ± 0.30		3 AAIJ	12E LHCb	Repl. by AAIJ 12E
5619.7 ± 1.2 ± 1.2		7 ACOSTA	06 CDF	Repl. by AALTO-NEN 14B
not seen		8 ABE	93B CDF	Repl. by ABE 97B
5640 ± 50 ± 30	16	9 ALBAJAR	91E UA1	$p\bar{p}$ 630 GeV
5640 +100 -210	52	BARI	91 SFM	$\Lambda_b^0 \rightarrow p D^0 \pi^-$
5650 +150 -200	90	BARI	91 SFM	$\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-$

<sup>1</sup> Uses  $\Lambda_b^0 \rightarrow \chi_{c1} p K^-$ ,  $\Lambda_b^0 \rightarrow \chi_{c2} p K^-$ ,  $\Lambda_b^0 \rightarrow J/\psi \Lambda$ ,  $\Lambda_b^0 \rightarrow p \psi(2S) K^-$ ,  $\Lambda_b^0 \rightarrow p J/\psi \pi^+ \pi^- K^-$ , and  $\Lambda_b^0 \rightarrow p J/\psi K^-$  decays.

<sup>2</sup> Uses exclusively reconstructed final states  $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ ,  $\Lambda_c^+ D^-$  and  $\bar{B}^0 \rightarrow D^+ D_s^-$  decays. The uncertainty includes both statistical and systematic contributions.

<sup>3</sup> Uses  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  fully reconstructed decays.

<sup>4</sup> ABE 97B observed 38 events with a background of  $18 \pm 1.6$  events in the mass range 5.60–5.65 GeV/ $c^2$ , a significance of  $> 3.4$  standard deviations.

<sup>5</sup> Uses 4 fully reconstructed  $\Lambda_b$  events.

<sup>6</sup> Uses  $\Lambda_b^0 \rightarrow p \psi(2S) K^-$ ,  $\Lambda_b^0 \rightarrow p J/\psi \pi^+ \pi^- K^-$ , and  $\Lambda_b^0 \rightarrow p J/\psi K^-$  decays.

<sup>7</sup> Uses exclusively reconstructed final states containing a  $J/\psi \rightarrow \mu^+ \mu^-$  decays.

<sup>8</sup> ABE 93B states that, based on the signal claimed by ALBAJAR 91E, CDF should have found  $30 \pm 23$   $\Lambda_b^0 \rightarrow J/\psi(1S) \Lambda$  events. Instead, CDF found not more than 2 events.

<sup>9</sup> ALBAJAR 91E claims  $16 \pm 5$  events above a background of  $9 \pm 1$  events, a significance of about 5 standard deviations.

**$m_{\Lambda_b^0} - m_{B^0}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>339.2±1.4±0.1</b>	<sup>1</sup> ACOSTA	06	CDF $p\bar{p}$ at 1.96 TeV

<sup>1</sup> Uses exclusively reconstructed final states containing  $J/\psi \rightarrow \mu^+ \mu^-$  decays.

 **$m_{\Lambda_b^0} - m_{B^+}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>339.72±0.28 OUR AVERAGE</b>			
339.72±0.24±0.18	<sup>1</sup> AAIJ	14AA LHCb	$p\bar{p}$ at 7 TeV
339.71±0.71±0.09	<sup>2</sup> AAIJ	12E LHCb	$p\bar{p}$ at 7 TeV

<sup>1</sup> Uses exclusively reconstructed final states  $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ ,  $\Lambda_c^+ D^-$  and  $\bar{B}^0 \rightarrow D^+ D_s^-$  decays.

<sup>2</sup> Uses exclusively reconstructed final states containing  $J/\psi \rightarrow \mu^+ \mu^-$  decays.

 **$\Lambda_b^0$  MEAN LIFE**

See *b*-baryon Admixture section for data on *b*-baryon mean life average over species of *b*-baryon particles.

“OUR EVALUATION” is an average using rescaled values of the data listed below. The average and rescaling were performed by the Heavy Flavor Averaging Group (HFLAV) and are described at <https://hflav.web.cern.ch/>. The averaging/rescaling procedure takes into account correlations between the measurements and asymmetric lifetime errors.

VALUE ( $10^{-12}$ s)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1.471±0.009 OUR EVALUATION</b>				
1.477±0.027±0.009	<sup>1</sup> SIRUNYAN	18BY CMS	$p\bar{p}$ at 8 TeV	
1.415±0.027±0.006	<sup>2</sup> AAIJ	14E LHCb	$p\bar{p}$ at 7 TeV	
1.479±0.009±0.010	<sup>3</sup> AAIJ	14U LHCb	$p\bar{p}$ at 7, 8 TeV	
1.565±0.035±0.020	<sup>2</sup> AALTONEN	14B CDF	$p\bar{p}$ at 1.96 TeV	
1.449±0.036±0.017	<sup>2</sup> AAD	13U ATLAS	$p\bar{p}$ at 7 TeV	
1.503±0.052±0.031	<sup>2</sup> CHATRCHYAN	13AC CMS	$p\bar{p}$ at 7 TeV	
1.303±0.075±0.035	<sup>2</sup> ABAZOV	12U D0	$p\bar{p}$ at 1.96 TeV	
1.401±0.046±0.035	<sup>4</sup> AALTONEN	10B CDF	$p\bar{p}$ at 1.96 TeV	
1.27 $^{+0.35}_{-0.29}$ $\pm 0.09$	ABREU	95S DLPH	Excess $p\mu^-$ , decay lengths	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.482±0.018±0.012	<sup>5</sup> AAIJ	13BB LHCb	Repl. by AAIJ 14U	
1.537±0.045±0.014	<sup>2</sup> AALTONEN	11 CDF	Repl. by AALTONEN 14B	
1.218 $^{+0.130}_{-0.115}$ $\pm 0.042$	<sup>2</sup> ABAZOV	07S D0	Repl. by ABAZOV 12U	
1.290 $^{+0.119}_{-0.110}$ $^{+0.087}_{-0.091}$	<sup>6</sup> ABAZOV	07U D0	$p\bar{p}$ at 1.96 TeV	
1.593 $^{+0.083}_{-0.078}$ $\pm 0.033$	<sup>2</sup> ABULENCIA	07A CDF	Repl. by AALTONEN 11	
1.22 $^{+0.22}_{-0.18}$ $\pm 0.04$	<sup>2</sup> ABAZOV	05C D0	Repl. by ABAZOV 07S	
1.11 $^{+0.19}_{-0.18}$ $\pm 0.05$	<sup>7</sup> ABREU	99W DLPH	$e^+ e^- \rightarrow Z$	

1.29	$\begin{array}{l} +0.24 \\ -0.22 \end{array}$	$\pm 0.06$	7 ACKERSTAFF	98G OPAL	$e^+ e^- \rightarrow Z$
1.21	$\pm 0.11$		7 BARATE	98D ALEP	$e^+ e^- \rightarrow Z$
1.32	$\pm 0.15$	$\pm 0.07$	8 ABE	96M CDF	$p\bar{p}$ at 1.8 TeV
1.19	$\begin{array}{l} +0.21 \\ -0.18 \end{array}$	$\begin{array}{l} +0.07 \\ -0.08 \end{array}$	ABREU	96D DLPH	Repl. by ABREU 99W
1.14	$\begin{array}{l} +0.22 \\ -0.19 \end{array}$	$\pm 0.07$	69 AKERS	95K OPAL	Repl. by ACKERSTAFF 98G
1.02	$\begin{array}{l} +0.23 \\ -0.18 \end{array}$	$\pm 0.06$	44 BUSKULIC	95L ALEP	Repl. by BARATE 98D

<sup>1</sup> Measured using  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  decays.

<sup>2</sup> Measured mean life using fully reconstructed  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  decays.

<sup>3</sup> Used  $\Lambda_b^0 \rightarrow J/\psi p K^-$  decays.

<sup>4</sup> Measured mean life using fully reconstructed  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$  decays.

<sup>5</sup> Measured the lifetime ratio of decays  $\Lambda_b^0 \rightarrow J/\psi p K^-$  to  $B^0 \rightarrow J/\psi \pi^+ K^-$  to be  $0.976 \pm 0.012 \pm 0.006$  with  $\tau_{B^0} = 1.519 \pm 0.007$  ps.

<sup>6</sup> Measured using semileptonic decays  $\Lambda_b^0 \rightarrow \Lambda_c^+ \mu\nu X$  and  $\Lambda_c^+ \rightarrow K_S^0 \mu$ .

<sup>7</sup> Measured using  $\Lambda_c \ell^-$  and  $\Lambda \ell^+ \ell^-$ .

<sup>8</sup> Excess  $\Lambda_c \ell^-$ , decay lengths.

## $\tau_{\Lambda_b^0}/\tau_{\Lambda_b^0}$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.940 <math>\pm 0.035 \pm 0.006</math></b>	<sup>1</sup> AAIJ	14E LHCb	$p\bar{p}$ at 7 TeV

<sup>1</sup> Measured using  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  decays.

## $\tau_{\Lambda_b^0}/\tau_{B^0}$ MEAN LIFE RATIO

### $\tau_{\Lambda_b^0}/\tau_{B^0}$ (direct measurements)

"OUR EVALUATION" has been obtained by the Heavy Flavor Averaging Group (HFLAV) by including both  $B^0$  and  $B^+$  decays.

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.964 <math>\pm 0.007</math> OUR EVALUATION</b>			

**0.970  $\pm 0.009$  OUR AVERAGE** Error includes scale factor of 1.4. See the ideogram below.

0.978 $\pm 0.018 \pm 0.006$	<sup>1</sup> SIRUNYAN	18BY CMS	$p\bar{p}$ at 8 TeV
0.929 $\pm 0.018 \pm 0.004$	<sup>1</sup> AAIJ	14E LHCb	$p\bar{p}$ at 7 TeV
0.974 $\pm 0.006 \pm 0.004$	<sup>2</sup> AAIJ	14U LHCb	$p\bar{p}$ at 7, 8 TeV
0.960 $\pm 0.025 \pm 0.016$	<sup>3</sup> AAD	13U ATLAS	$p\bar{p}$ at 7 TeV
0.864 $\pm 0.052 \pm 0.033$	<sup>4,5</sup> ABAZOV	12U D0	$p\bar{p}$ at 1.96 TeV
1.020 $\pm 0.030 \pm 0.008$	<sup>4</sup> AALTONEN	11 CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.976 $\pm 0.012 \pm 0.006$	<sup>6</sup> AAIJ	13BB LHCb	Repl. by AAIJ 14U
$0.811^{+0.096}_{-0.087} \pm 0.034$	<sup>4,5</sup> ABAZOV	07S D0	Repl. by ABAZOV 12U
1.041 $\pm 0.057$	<sup>7</sup> ABULENCIA	07A CDF	Repl. by AALTONEN 11
$0.87^{+0.17}_{-0.14} \pm 0.03$	<sup>7</sup> ABAZOV	05C D0	Repl. by ABAZOV 07S

<sup>1</sup> Measured using  $\Lambda_b^0 \rightarrow J/\psi \Lambda$  and  $B^0 \rightarrow J/\psi K^*(892)^0$  decays.

<sup>2</sup> Used  $\Lambda_b^0 \rightarrow J/\psi p K^-$  and  $B^0 \rightarrow J/\psi K^*(892)^0$  decays.


<sup>3</sup> Measured with  $\Lambda_b^0 \rightarrow J/\psi(\mu^+ \mu^-) \Lambda^0(p\pi^-)$  decays.

<sup>4</sup> Uses fully reconstructed  $\Lambda_b \rightarrow J/\psi \Lambda$  decays.

<sup>5</sup> Uses  $B^0 \rightarrow J/\psi K_S^0$  decays for denominator.

<sup>6</sup> Measures  $1/\tau_{\Lambda_b^0} - 1/\tau_{B^0}$  and uses  $\tau_{B^0} = 1.519 \pm 0.007$  ps to extract lifetime ratio.

<sup>7</sup> Measured mean life ratio using fully reconstructed decays.



## $\Lambda_b^0$ DECAY MODES

The branching fractions  $B(b\text{-baryon} \rightarrow \Lambda \ell^- \bar{\nu}_\ell \text{anything})$  and  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})$  are not pure measurements because the underlying measured products of these with  $B(b \rightarrow b\text{-baryon})$  were used to determine  $B(b \rightarrow b\text{-baryon})$ , as described in the note “Production and Decay of  $b$ -Flavored Hadrons.”

For inclusive branching fractions, e.g.,  $\Lambda_b \rightarrow \bar{\Lambda}_c$  anything, the values usually are multiplicities, not branching fractions. They can be greater than one.

Mode	Fraction ( $\Gamma_i/\Gamma$ )	Scale factor/ Confidence level
$\Gamma_1$ $J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)$	$(5.8 \pm 0.8) \times 10^{-5}$	
$\Gamma_2$ $J/\psi(1S)\Lambda$		
$\Gamma_3$ $J/\psi(1S)\Lambda\phi$		
$\Gamma_4$ $\psi(2S)\Lambda$		

$\Gamma_5$	$p D^0 \pi^-$	$( 6.2 \pm 0.6 ) \times 10^{-4}$
$\Gamma_6$	$p D^+ \pi^- \pi^-$	$( 2.7 \pm 0.4 ) \times 10^{-4}$
$\Gamma_7$	$p D^*(2010)^+ \pi^- \pi^-$	$( 5.2 \pm 1.0 ) \times 10^{-4}$
$\Gamma_8$	$\Lambda_c(2860)^+ \pi^-, \Lambda_c^+ \rightarrow D^0 p$	
$\Gamma_9$	$\Lambda_c(2880)^+ \pi^-, \Lambda_c^+ \rightarrow D^0 p$	
$\Gamma_{10}$	$\Lambda_c(2940)^+ \pi^-, \Lambda_c^+ \rightarrow D^0 p$	
$\Gamma_{11}$	$p D^0 K^-$	$( 4.5 \pm 0.8 ) \times 10^{-5}$
$\Gamma_{12}$	$p D K^-, D \rightarrow K^- \pi^+$	
$\Gamma_{13}$	$p D K^-, D \rightarrow K^+ \pi^-$	
$\Gamma_{14}$	$p J/\psi \pi^-$	$( 2.6 \begin{array}{l} +0.5 \\ -0.4 \end{array} ) \times 10^{-5}$
$\Gamma_{15}$	$p \pi^- J/\psi, J/\psi \rightarrow \mu^+ \mu^-$	$( 1.6 \pm 0.8 ) \times 10^{-6}$
$\Gamma_{16}$	$p J/\psi K^-$	$( 3.2 \begin{array}{l} +0.6 \\ -0.5 \end{array} ) \times 10^{-4}$
$\Gamma_{17}$	$p \eta_c(1S) K^-$	$( 1.06 \pm 0.26 ) \times 10^{-4}$
$\Gamma_{18}$	$P_c(4312)^+ K^-, P_c(4312)^+ \rightarrow p \eta_c(1S)$	$< 2.5 \times 10^{-5}$ CL=95%
$\Gamma_{19}$	$P_c(4380)^+ K^-, P_c \rightarrow p J/\psi$	[a] $( 2.7 \pm 1.4 ) \times 10^{-5}$
$\Gamma_{20}$	$P_c(4450)^+ K^-, P_c \rightarrow p J/\psi$	[a] $( 1.3 \pm 0.4 ) \times 10^{-5}$
$\Gamma_{21}$	$\chi_{c1}(1P) p K^-$	$( 7.6 \begin{array}{l} +1.5 \\ -1.3 \end{array} ) \times 10^{-5}$
$\Gamma_{22}$	$\chi_{c1}(1P) p \pi^-$	$( 5.0 \begin{array}{l} +1.3 \\ -1.1 \end{array} ) \times 10^{-6}$
$\Gamma_{23}$	$\chi_{c2}(1P) p K^-$	$( 7.9 \begin{array}{l} +1.6 \\ -1.4 \end{array} ) \times 10^{-5}$
$\Gamma_{24}$	$\chi_{c2}(1P) p \pi^-$	$( 4.8 \pm 1.9 ) \times 10^{-6}$
$\Gamma_{25}$	$p J/\psi(1S) \pi^+ \pi^- K^-$	$( 6.6 \begin{array}{l} +1.3 \\ -1.1 \end{array} ) \times 10^{-5}$
$\Gamma_{26}$	$p \psi(2S) K^-$	$( 6.6 \begin{array}{l} +1.2 \\ -1.0 \end{array} ) \times 10^{-5}$
$\Gamma_{27}$	$\chi_{c1}(3872) p K^-$	$( 3.2 \pm 1.4 ) \times 10^{-5}$
$\Gamma_{28}$	$\chi_{c1}(3872) \Lambda(1520)$	$( 1.9 \pm 0.9 ) \times 10^{-5}$
$\Gamma_{29}$	$\psi(2S) p \pi^-$	$( 7.5 \begin{array}{l} +1.6 \\ -1.4 \end{array} ) \times 10^{-6}$
$\Gamma_{30}$	$p \bar{K}^0 \pi^-$	$( 1.3 \pm 0.4 ) \times 10^{-5}$
$\Gamma_{31}$	$p K^0 K^-$	$< 3.5 \times 10^{-6}$ CL=90%
$\Gamma_{32}$	$\Lambda_c^+ \pi^-$	$( 4.9 \pm 0.4 ) \times 10^{-3}$ S=1.2
$\Gamma_{33}$	$\Lambda_c^+ K^-$	$( 3.56 \pm 0.28 ) \times 10^{-4}$ S=1.2
$\Gamma_{34}$	$\Lambda_c^+ a_1(1260)^-$	seen
$\Gamma_{35}$	$\Lambda_c^+ D^-$	$( 4.6 \pm 0.6 ) \times 10^{-4}$
$\Gamma_{36}$	$\Lambda_c^+ D_s^-$	$( 1.10 \pm 0.10 ) \%$
$\Gamma_{37}$	$\Lambda_c^+ \pi^+ \pi^- \pi^-$	$( 7.6 \pm 1.1 ) \times 10^{-3}$ S=1.1
$\Gamma_{38}$	$\Lambda_c(2595)^+ \pi^-, \Lambda_c(2595)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-$	$( 3.4 \pm 1.4 ) \times 10^{-4}$
$\Gamma_{39}$	$\Lambda_c(2625)^+ \pi^-, \Lambda_c(2625)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-$	$( 3.3 \pm 1.3 ) \times 10^{-4}$

$\Gamma_{40}$	$\Sigma_c(2455)^0 \pi^+ \pi^-$ , $\Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-$	$(5.7 \pm 2.2) \times 10^{-4}$
$\Gamma_{41}$	$\Sigma_c(2455)^{++} \pi^- \pi^-$ , $\Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+$	$(3.2 \pm 1.5) \times 10^{-4}$
$\Gamma_{42}$	$\Lambda_c^+ K^+ K^- \pi^-$	$(1.02 \pm 0.11) \times 10^{-3}$
$\Gamma_{43}$	$\Lambda_c^+ p \bar{p} \pi^-$	$(2.63 \pm 0.27) \times 10^{-4}$
$\Gamma_{44}$	$\Sigma_c(2455)^0 p \bar{p}$ , $\Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-$	$(2.3 \pm 0.5) \times 10^{-5}$
$\Gamma_{45}$	$\Sigma_c(2520)^0 p \bar{p}$ , $\Sigma_c(2520)^0 \rightarrow \Lambda_c^+ \pi^-$	$(3.1 \pm 0.7) \times 10^{-5}$
$\Gamma_{46}$	$\Lambda K^0 2\pi^+ 2\pi^-$	
$\Gamma_{47}$	$\Lambda_c^+ \ell^- \bar{\nu}_\ell$ anything	[b] $(10.9 \pm 2.2) \%$
$\Gamma_{48}$	$\Lambda_c^+ \ell^- \bar{\nu}_\ell$	$(6.2 \pm 1.4) \%$
$\Gamma_{49}$	$\Lambda_c^+ \tau^- \bar{\nu}_\tau$	$(1.9 \pm 0.5) \%$
$\Gamma_{50}$	$\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell$	$(5.6 \pm 3.1) \%$
$\Gamma_{51}$	$\Lambda_c(2595)^+ \ell^- \bar{\nu}_\ell$	$(7.9 \pm 4.0) \times 10^{-3}$
$\Gamma_{52}$	$\Lambda_c(2625)^+ \ell^- \bar{\nu}_\ell$	$(1.3 \pm 0.6) \%$
$\Gamma_{53}$	$\Sigma_c(2455)^0 \pi^+ \ell^- \bar{\nu}_\ell$	
$\Gamma_{54}$	$\Sigma_c(2455)^{++} \pi^- \ell^- \bar{\nu}_\ell$	
$\Gamma_{55}$	$p h^-$	[c] $< 2.3 \times 10^{-5}$ CL=90%
$\Gamma_{56}$	$p \pi^-$	$(4.5 \pm 0.8) \times 10^{-6}$
$\Gamma_{57}$	$p K^-$	$(5.4 \pm 1.0) \times 10^{-6}$
$\Gamma_{58}$	$p D_s^-$	$< 4.8 \times 10^{-4}$ CL=90%
$\Gamma_{59}$	$p \mu^- \bar{\nu}_\mu$	$(4.1 \pm 1.0) \times 10^{-4}$
$\Gamma_{60}$	$\Lambda \mu^+ \mu^-$	$(1.08 \pm 0.28) \times 10^{-6}$
$\Gamma_{61}$	$p \pi^- \mu^+ \mu^-$	$(6.9 \pm 2.5) \times 10^{-8}$
$\Gamma_{62}$	$p K^- e^+ e^-$	$(3.1 \pm 0.6) \times 10^{-7}$
$\Gamma_{63}$	$p K^- \mu^+ \mu^-$	$(2.6 \pm 0.5) \times 10^{-7}$
$\Gamma_{64}$	$\Lambda \gamma$	$(7.1 \pm 1.7) \times 10^{-6}$
$\Gamma_{65}$	$\Lambda \eta$	$(9 \pm 7) \times 10^{-6}$
$\Gamma_{66}$	$\Lambda \eta'(958)$	$< 3.1 \times 10^{-6}$ CL=90%
$\Gamma_{67}$	$\Lambda \pi^+ \pi^-$	$(4.6 \pm 1.9) \times 10^{-6}$
$\Gamma_{68}$	$\Lambda K^+ \pi^-$	$(5.6 \pm 1.2) \times 10^{-6}$
$\Gamma_{69}$	$\Lambda K^+ K^-$	$(1.60 \pm 0.22) \times 10^{-5}$
$\Gamma_{70}$	$\Lambda \phi$	$(9.8 \pm 2.6) \times 10^{-6}$
$\Gamma_{71}$	$p \pi^- \pi^+ \pi^-$	$(2.09 \pm 0.21) \times 10^{-5}$
$\Gamma_{72}$	$p K^- K^+ \pi^-$	$(4.0 \pm 0.6) \times 10^{-6}$
$\Gamma_{73}$	$p K^- \pi^+ \pi^-$	$(5.0 \pm 0.5) \times 10^{-5}$
$\Gamma_{74}$	$p K^- K^+ K^-$	$(1.25 \pm 0.13) \times 10^{-5}$

[a]  $P_c^+$  is a pentaquark-charmonium state.

[b] Not a pure measurement. See note at head of  $\Lambda_b^0$  Decay Modes.

[c] Here  $h^-$  means  $\pi^-$  or  $K^-$ .

## CONSTRAINED FIT INFORMATION

An overall fit to 10 branching ratios uses 12 measurements and one constraint to determine 7 parameters. The overall fit has a  $\chi^2 = 10.8$  for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients  $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$ , in percent, from the fit to the branching fractions,  $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$ . The fit constrains the  $x_i$  whose labels appear in this array to sum to one.

x33	92				
x37	46	43			
x48	13	12	6		
x56	0	0	0	0	
x57	0	0	0	0	82
	x32	x33	x37	x48	x56

## $\Lambda_b^0$ BRANCHING RATIOS

$$\Gamma(J/\psi(1S)\Lambda \times B(b \rightarrow \Lambda_b^0)) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$

VALUE (units $10^{-5}$ )	EVTS	DOCUMENT ID	TECN	COMMENT
--------------------------	------	-------------	------	---------

**5.8 ± 0.8 OUR AVERAGE**

$6.01 \pm 0.60 \pm 0.58 \pm 0.28$  <sup>1</sup> ABAZOV 110 D0  $p\bar{p}$  at 1.96 TeV  
 $4.7 \pm 2.3 \pm 0.2$  <sup>2</sup> ABE 97B CDF  $p\bar{p}$  at 1.8 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

180  $\pm 60 \pm 90$  16 ALBAJAR 91E UA1  $p\bar{p}$  at 630 GeV

<sup>1</sup> ABAZOV 110 uses  $B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0) = (1.74 \pm 0.08) \times 10^{-4}$  to obtain the result. The  $(\pm 0.08) \times 10^{-4}$  uncertainty of this product is listed as the last uncertainty of the measurement,  $(\pm 0.28) \times 10^{-5}$ .

<sup>2</sup> ABE 97B reports  $[B(\Lambda_b^0 \rightarrow J/\psi \Lambda) \times B(b \rightarrow \Lambda_b^0)] / [B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0)] = 0.27 \pm 0.12 \pm 0.05$ . We multiply by our best value  $B(B^0 \rightarrow J/\psi K_S^0) \times B(b \rightarrow B^0) = (1.74 \pm 0.08) \times 10^{-4}$ . Our first error is their experiment error and our second error is the systematic error from using our best value.

$$\Gamma(\psi(2S)\Lambda) / \Gamma(J/\psi(1S)\Lambda)$$

$$\Gamma_4 / \Gamma_2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

**0.508 ± 0.023 OUR AVERAGE**

$0.513 \pm 0.023 \pm 0.019$  <sup>1</sup> AAIJ 19F LHCb  $p\bar{p}$  at 7, 8 TeV  
 $0.50 \pm 0.03 \pm 0.02$  <sup>2</sup> AAD 15CH ATLAS  $p\bar{p}$  at 8 TeV

<sup>1</sup> AAIJ 19F uses  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$  and  $B(\psi(2S) \rightarrow e^+ e^-) = (7.93 \pm 0.17) \times 10^{-3}$  from PDG 18 with assumption of lepton universality. AAIJ 19F

reports this result as  $0.513 \pm 0.023 \pm 0.016 \pm 0.011$ , where the last uncertainty is the contribution due to the external input of branching fractions used in the analysis.

<sup>2</sup> AAD 15CH uses  $B(J/\psi \rightarrow \mu^+ \mu^-) = (5.961 \pm 0.033) \times 10^{-2}$  and  $B(\psi(2S) \rightarrow \mu^+ \mu^-) = (7.89 \pm 0.17) \times 10^{-3}$  from PDG 14 with assumption of lepton universality.

### $\Gamma(J/\psi(1S)\Lambda\phi)/\Gamma(\psi(2S)\Lambda)$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.26 ± 0.90 ± 0.69</b>	SIRUNYAN 20H	CMS	$p p$ at 13 TeV

### $\Gamma_3/\Gamma_4$

### $\Gamma(pD^0\pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
seen	52	BARI	91	SFM $D^0 \rightarrow K^-\pi^+$
seen		BASILE	81	SFM $D^0 \rightarrow K^-\pi^+$

### $\Gamma_5/\Gamma$

### $\Gamma(pD^+\pi^-\pi^-)/\Gamma(\Lambda_c^+\pi^+\pi^-\pi^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>3.57 ± 0.18 <sup>+0.17</sup> <sub>-0.18</sub></b>	<sup>1</sup> AAIJ	22R	LHCb $p p$ at 7 and 8 TeV

### $\Gamma_6/\Gamma_{37}$

<sup>1</sup> AAIJ 22R reports  $[\Gamma(\Lambda_b^0 \rightarrow p D^+\pi^-\pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^+\pi^-\pi^-)] \times [B(D^+ \rightarrow K^-2\pi^+)/[B(\Lambda_c^+ \rightarrow p K^-\pi^+)] = 5.35 \pm 0.21 \pm 0.16$  % which we multiply or divide by our best values  $B(D^+ \rightarrow K^-2\pi^+) = (9.38 \pm 0.16) \times 10^{-2}$ ,  $B(\Lambda_c^+ \rightarrow p K^-\pi^+) = (6.26 \pm 0.29) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

### $\Gamma(pD^*(2010)^+\pi^-\pi^-)/\Gamma(pD^+\pi^-\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.90 ± 0.19</b>	<sup>1</sup> AAIJ	22R	LHCb $p p$ at 7 and 8 TeV

### $\Gamma_7/\Gamma_6$

<sup>1</sup> AAIJ 22R uses partial reconstruction of  $p D^+\pi^-\pi^-$  final state.

### $\Gamma(\Lambda_c(2860)^+\pi^-, \Lambda_c^+ \rightarrow D^0 p)/\Gamma(\Lambda_c(2880)^+\pi^-, \Lambda_c^+ \rightarrow D^0 p)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>4.54 <sup>+0.51</sup> <sub>-0.39</sub> <sup>+0.21</sup> <sub>-0.59</sub></b>	AAIJ	17S	LHCb $p p$ at 7, 8 TeV

### $\Gamma_8/\Gamma_9$

### $\Gamma(\Lambda_c(2940)^+\pi^-, \Lambda_c^+ \rightarrow D^0 p)/\Gamma(\Lambda_c(2880)^+\pi^-, \Lambda_c^+ \rightarrow D^0 p)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.83 <sup>+0.31</sup> <sub>-0.10</sub> <sup>+0.18</sup> <sub>-0.43</sub></b>	AAIJ	17S	LHCb $p p$ at 7, 8 TeV

### $\Gamma_{10}/\Gamma_9$

### $\Gamma(pD^0K^-)/\Gamma(pD^0\pi^-)$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>7.3 ± 0.8 <sup>+0.5</sup> <sub>-0.6</sub></b>	AAIJ	14H	LHCb $p p$ at 7 TeV

### $\Gamma_{11}/\Gamma_5$

$\Gamma(pDK^-, D \rightarrow K^-\pi^+)/\Gamma(pDK^-, D \rightarrow K^+\pi^-)$	$\Gamma_{12}/\Gamma_{13}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.1 \pm 0.8^{+0.4}_{-0.3}</math></b>	<sup>1</sup> AAIJ	21AD LHCb	$p\bar{p}$ at 7, 8, 13 TeV

<sup>1</sup> Measured in the full phase space.

$\Gamma(pJ/\psi\pi^-)/\Gamma(pJ/\psi K^-)$	$\Gamma_{14}/\Gamma_{16}$		
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>8.24 \pm 0.25 \pm 0.42</math></b>	AAIJ	14K	LHCb $p\bar{p}$ at 7, 8 TeV

$\Gamma(pJ/\psi K^-)/\Gamma_{\text{total}}$	$\Gamma_{16}/\Gamma$		
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>3.17 \pm 0.04^{+0.57}_{-0.45}</math></b>	<sup>1</sup> AAIJ	16A	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 16A reported the measurement of  $(3.17 \pm 0.04 \pm 0.07 \pm 0.34^{+0.45}_{-0.28}) \times 10^{-4}$  where the first uncertainty is statistical, the second is systematic, the third is due to the branching fraction of  $B^0 \rightarrow J/\psi K^*(892)^0$ , and the fourth is due to the knowledge of  $f_{\Lambda_b}/f_d$ . We combined in quadrature second to fourth uncertainties to a total systematic uncertainty.

$\Gamma(p\eta_c(1S)K^-)/\Gamma(pJ/\psi K^-)$	$\Gamma_{17}/\Gamma_{16}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.333 \pm 0.050 \pm 0.037</math></b>	<sup>1</sup> AAIJ	20AK	LHCb $p\bar{p}$ at 13 TeV

<sup>1</sup> AAIJ 20AK reported the measurement of  $0.333 \pm 0.050 \pm 0.019 \pm 0.032$ , where the last uncertainty is due uncertainties of the used branching fractions of  $J/\psi \rightarrow p\bar{p}$  and  $\eta_c \rightarrow p\bar{p}$  decays. We combined in quadrature the systematic uncertainties.

$\Gamma(P_c(4312)^+ K^-, P_c(4312)^+ \rightarrow p\eta_c(1S))/\Gamma(p\eta_c(1S)K^-)$	$\Gamma_{18}/\Gamma_{17}$			
<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;0.24</b>	95	AAIJ	20AK	LHCb $p\bar{p}$ at 13 TeV

$\Gamma(P_c(4380)^+ K^-, P_c \rightarrow pJ/\psi)/\Gamma_{\text{total}}$	$\Gamma_{19}/\Gamma$		
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.66 \pm 0.22^{+1.41}_{-1.38}</math></b>	<sup>1</sup> AAIJ	16A	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 16 total systematic includes the uncertainties on  $f(P_c^+)$  and  $B(\Lambda_b \rightarrow pJ/\psi K^-)$ .

$\Gamma(P_c(4450)^+ K^-, P_c \rightarrow pJ/\psi)/\Gamma_{\text{total}}$	$\Gamma_{20}/\Gamma$		
<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>1.30 \pm 0.16^{+0.42}_{-0.39}</math></b>	<sup>1</sup> AAIJ	16A	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 16 total systematic includes the uncertainties on  $f(P_c^+)$  and  $B(\Lambda_b \rightarrow pJ/\psi K^-)$ .

$\Gamma(\chi_{c1}(1P)pK^-)/\Gamma(pJ/\psi K^-)$   $\Gamma_{21}/\Gamma_{16}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.239±0.019±0.007</b>	<sup>1</sup> AAIJ	17AM LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> AAIJ 17AM reports  $0.242 \pm 0.014 \pm 0.016$  from a measurement of  $[\Gamma(\Lambda_b^0 \rightarrow \chi_{c1}(1P)pK^-)/\Gamma(\Lambda_b^0 \rightarrow pJ/\psi K^-)] \times [B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (33.9 \pm 1.2) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c1}(1P) \rightarrow \gamma J/\psi(1S)) = (34.3 \pm 1.0) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\chi_{c1}(1P)p\pi^-)/\Gamma(\chi_{c1}(1P)pK^-)$   $\Gamma_{22}/\Gamma_{21}$ 

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.59±1.01±0.22</b>	AAIJ	21R LHCb	$p p$ at 13 TeV

 $\Gamma(\chi_{c2}(1P)pK^-)/\Gamma(pJ/\psi K^-)$   $\Gamma_{23}/\Gamma_{16}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.250±0.025±0.007</b>	<sup>1</sup> AAIJ	17AM LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> AAIJ 17AM reports  $0.248 \pm 0.02 \pm 0.017$  from a measurement of  $[\Gamma(\Lambda_b^0 \rightarrow \chi_{c2}(1P)pK^-)/\Gamma(\Lambda_b^0 \rightarrow pJ/\psi K^-)] \times [B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S))]$  assuming  $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.2 \pm 0.7) \times 10^{-2}$ , which we rescale to our best value  $B(\chi_{c2}(1P) \rightarrow \gamma J/\psi(1S)) = (19.0 \pm 0.5) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(\chi_{c2}(1P)pK^-)/\Gamma(\chi_{c1}(1P)pK^-)$   $\Gamma_{23}/\Gamma_{21}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.06±0.05±0.04±0.04</b>	<sup>1</sup> AAIJ	21R LHCb	$p p$ at 13 TeV

<sup>1</sup> The first uncertainty is statistical, the second is systematic and the third is related to the uncertainties in the branching fractions of the  $\chi_{cJ} \rightarrow J/\psi \gamma$  decays.

 $\Gamma(\chi_{c2}(1P)p\pi^-)/\Gamma(\chi_{c1}(1P)p\pi^-)$   $\Gamma_{24}/\Gamma_{22}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.95±0.30±0.04±0.04</b>	<sup>1</sup> AAIJ	21R LHCb	$p p$ at 13 TeV

<sup>1</sup> Evidence for the  $\Lambda_b^0 \rightarrow \chi_{c2} p\pi^-$  decay is obtained with a significance of 3.5 standard deviations. The first uncertainty is statistical, the second is systematic and the third is related to the uncertainties in the branching fractions of the  $\chi_{cJ} \rightarrow J/\psi \gamma$  decays.

 $\Gamma(pJ/\psi(1S)\pi^+\pi^-K^-)/\Gamma(pJ/\psi K^-)$   $\Gamma_{25}/\Gamma_{16}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.2086±0.0096±0.0134</b>	<sup>1</sup> AAIJ	16Y LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> Excludes  $\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$ .

 $\Gamma(p\psi(2S)K^-)/\Gamma(pJ/\psi K^-)$   $\Gamma_{26}/\Gamma_{16}$ 

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.2070±0.0076±0.0059</b>	<sup>1</sup> AAIJ	16Y LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> AAIJ 16Y reports a measurement of  $0.2070 \pm 0.0076 \pm 0.0046 \pm 0.0037$  where the third uncertainty is due to the knowledge of  $J/\psi$  and  $\psi(2S)$  branching fractions. We have combined both systematic uncertainties in quadrature.

$\Gamma(\chi_{c1}(3872)\Lambda(1520))/\Gamma(\chi_{c1}(3872)pK^-)$   $\Gamma_{28}/\Gamma_{27}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.58±0.15</b>	AAIJ	19AN LHCb	$p p$ at 7, 8, 13 TeV

 $\Gamma(\chi_{c1}(3872)pK^-)/\Gamma(p\psi(2S)K^-)$   $\Gamma_{27}/\Gamma_{26}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.49±0.10±0.16</b>	<sup>1</sup> AAIJ	19AN LHCb	$p p$ at 7, 8, 13 TeV

<sup>1</sup> AAIJ 19AN reports  $[\Gamma(\Lambda_b^0 \rightarrow \chi_{c1}(3872)pK^-)/\Gamma(\Lambda_b^0 \rightarrow p\psi(2S)K^-)] \times [B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S))] / [B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-)] = (5.4 \pm 1.1 \pm 0.2) \times 10^{-2}$  which we multiply or divide by our best values  $B(\chi_{c1}(3872) \rightarrow \pi^+\pi^- J/\psi(1S)) = (3.8 \pm 1.2) \times 10^{-2}$ ,  $B(\psi(2S) \rightarrow J/\psi(1S)\pi^+\pi^-) = (34.68 \pm 0.30) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

 $\Gamma(p\psi(2S)p\pi^-)/\Gamma(p\psi(2S)K^-)$   $\Gamma_{29}/\Gamma_{26}$ 

<u>VALUE (%)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>11.4±1.3±0.2</b>	AAIJ	18AF LHCb	$p p$ at 7, 8, 13 TeV

 $\Gamma(p\bar{K}^0\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{30}/\Gamma$ 

<u>VALUE (units <math>10^{-5}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>1.26±0.19±0.36</b>	<sup>1</sup> AAIJ	14Q LHCb	$p p$ at 7 TeV

<sup>1</sup> Used the normalizing mode branching fraction value of  $B(B^0 \rightarrow K^0\pi^+\pi^-) = (4.96 \pm 0.20) \times 10^{-5}$ .

 $\Gamma(pK^0K^-)/\Gamma_{\text{total}}$   $\Gamma_{31}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;3.5 × 10<sup>-6</sup></b>	90	AAIJ	14Q LHCb	$p p$ at 7 TeV

 $\Gamma(\Lambda_c^+\pi^-)/\Gamma_{\text{total}}$   $\Gamma_{32}/\Gamma$ 

<u>VALUE (units <math>10^{-3}</math>)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.9 ± 0.4 OUR FIT</b>	Error includes scale factor of 1.2.			

**4.8 ± 0.5 OUR AVERAGE** Error includes scale factor of 1.5.

$4.60^{+0.31}_{-0.30} \pm 0.14$  <sup>1</sup> AAIJ 14I LHCb  $p p$  at 7 TeV

$5.97 \pm 0.28 \pm 0.81$  <sup>2</sup> AAIJ 14Q LHCb  $p p$  at 7 TeV

$8.8 \pm 2.8 \pm 1.5$  <sup>3</sup> ABULENCIA 07B CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 3 ABREU 96N DLPH  $\Lambda_c^+ \rightarrow pK^-\pi^+$

seen 4 BUSKULIC 96L ALEP  $\Lambda_c^+ \rightarrow pK^-\pi^+, p\bar{K}^0, \Lambda\pi^+\pi^+\pi^-$

<sup>1</sup> AAIJ 14I reports  $(4.30 \pm 0.03^{+0.12}_{-0.11} \pm 0.26 \pm 0.21) \times 10^{-3}$  from a measurement of  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)/\Gamma_{\text{total}}] \times [B(B^0 \rightarrow D^-\pi^+)]$  assuming  $B(B^0 \rightarrow D^-\pi^+) = (2.68 \pm 0.13) \times 10^{-3}$ , which we rescale to our best value  $B(B^0 \rightarrow D^-\pi^+) = (2.51 \pm 0.08) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses information on  $f_{\text{baryon}}/f_d$  from measurement in semileptonic decays by the same authors.

<sup>2</sup> Obtained using the branching fraction of  $\Lambda_c^+ \rightarrow pK^-\pi^+$  decay.

<sup>3</sup> The result is obtained from  $(f_{\text{baryon}}/f_d) \cdot (B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/B(\bar{B}^0 \rightarrow D^+ \pi^-)) = 0.82 \pm 0.08 \pm 0.11 \pm 0.22$ , assuming  $f_{\text{baryon}}/f_d = 0.25 \pm 0.04$  and  $B(\bar{B}^0 \rightarrow D^+ \pi^-) = (2.68 \pm 0.13) \times 10^{-3}$ .

$\Gamma(pD^0\pi^-)/\Gamma(\Lambda_c^+\pi^-)$	$\Gamma_5/\Gamma_{32}$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

**0.128±0.007±0.006** <sup>1</sup> AAIJ 14H LHCb  $p p$  at 7 TeV

<sup>1</sup> AAIJ 14H reports  $[\Gamma(\Lambda_b^0 \rightarrow pD^0\pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)] \times [B(D^0 \rightarrow K^-\pi^+)/[B(\Lambda_c^+ \rightarrow pK^-\pi^+)] = (8.06 \pm 0.23 \pm 0.35) \times 10^{-2}$  which we multiply or divide by our best values  $B(D^0 \rightarrow K^-\pi^+) = (3.947 \pm 0.030) \times 10^{-2}$ ,  $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (6.26 \pm 0.29) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(\Lambda_c^+ K^-)/\Gamma_{\text{total}}$	$\Gamma_{33}/\Gamma$
<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

**3.56±0.28 OUR FIT** Error includes scale factor of 1.2.

**3.55±0.44±0.50** <sup>1</sup> AAIJ 14Q LHCb  $p p$  at 7 TeV

<sup>1</sup> Obtained using the branching fraction of  $\Lambda_c^+ \rightarrow pK^-\pi^+$  decay.

$\Gamma(\Lambda_c^+ K^-)/\Gamma(\Lambda_c^+\pi^-)$	$\Gamma_{33}/\Gamma_{32}$
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

**7.31±0.22 OUR FIT**

**7.31±0.16±0.16** AAIJ 14H LHCb  $p p$  at 7 TeV

$\Gamma(\Lambda_c^+ a_1(1260)^-)/\Gamma_{\text{total}}$	$\Gamma_{34}/\Gamma$
<u>VALUE</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

**seen**

**1**

ABREU

96N DLPH

$\Lambda_c^+ \rightarrow pK^-\pi^+, a_1^- \rightarrow \rho^0\pi^- \rightarrow \pi^+\pi^-\pi^-$

$\Gamma(\Lambda_c^+ D_s^-)/\Gamma_{\text{total}}$	$\Gamma_{36}/\Gamma$
<u>VALUE (units <math>10^{-2}</math>)</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

**1.1±0.1**

<sup>1</sup> AAIJ

14AA

LHCb

$p p$  at 7 TeV

<sup>1</sup> Uses  $B(\bar{B}^0 \rightarrow D^+ D_s^-) = (7.2 \pm 0.8) \times 10^{-3}$  and their measured  $B(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)/B(\bar{B}^0 \rightarrow D^+\pi^-)$  values.

$\Gamma(\Lambda_c^+ D_s^-)/\Gamma(\Lambda_c^+ D_s^-)$	$\Gamma_{35}/\Gamma_{36}$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

**0.042±0.003±0.003** AAIJ 14AA LHCb  $p p$  at 7 TeV

$\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}$	$\Gamma_{37}/\Gamma$
<u>VALUE (units <math>10^{-3}</math>)</u> <u>EVTS</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>

**7.6±1.1 OUR FIT** Error includes scale factor of 1.1.

**14.8<sup>+3.8</sup><sub>-3.1</sub>±1.1** <sup>1</sup> AALTONEN 12A CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen                    90                    BARI                    91                    SFM                     $\Lambda_c^+ \rightarrow p K^- \pi^+$   
<sup>1</sup> AALTONEN 12A reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \pi^-)/\Gamma_{\text{total}}] / [B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] = 3.04 \pm 0.33^{+0.70}_{-0.55}$  which we multiply by our best value  $B(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (4.9 \pm 0.4) \times 10^{-3}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$$\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-) / \Gamma(\Lambda_c^+ \pi^-) \quad \Gamma_{37}/\Gamma_{32}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>1.57 ± 0.21 OUR FIT</b>			
<b>1.43 ± 0.16 ± 0.13</b>	AAIJ	11E LHCb	$p p$ at 7 TeV

$$\Gamma(\Lambda_c(2595)^+ \pi^-, \Lambda_c(2595)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-) / \Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-) \quad \Gamma_{38}/\Gamma_{37}$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.4 ± 1.7 ± 0.6</b>	AAIJ	11E LHCb	$p p$ at 7 TeV

$$\Gamma(\Lambda_c(2625)^+ \pi^-, \Lambda_c(2625)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-) / \Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-) \quad \Gamma_{39}/\Gamma_{37}$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.3 ± 1.5 ± 0.4</b>	AAIJ	11E LHCb	$p p$ at 7 TeV

$$\Gamma(\Sigma_c(2455)^0 \pi^+ \pi^-, \Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-) / \Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-) \quad \Gamma_{40}/\Gamma_{37}$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>7.4 ± 2.4 ± 1.2</b>	AAIJ	11E LHCb	$p p$ at 7 TeV

$$\Gamma(\Sigma_c(2455)^{++} \pi^- \pi^-, \Sigma_c^{++} \rightarrow \Lambda_c^+ \pi^+) / \Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-) \quad \Gamma_{41}/\Gamma_{37}$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.2 ± 1.8 ± 0.7</b>	AAIJ	11E LHCb	$p p$ at 7 TeV

$$\Gamma(\Lambda_c^+ K^+ K^- \pi^-) / \Gamma(\Lambda_c^+ D_s^-) \quad \Gamma_{42}/\Gamma_{36}$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>9.26 ± 0.29 ± 0.53</b>	<sup>1</sup> AAIJ	21B LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> AAIJ 21B systematic uncertainty includes the contribution from the  $D_s^- \rightarrow K^+ K^- \pi^-$  branching fraction.

$$\Gamma(\Lambda_c^+ p \bar{p} \pi^-) / \Gamma(\Lambda_c^+ \pi^-) \quad \Gamma_{43}/\Gamma_{32}$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.40 ± 0.23 ± 0.32</b>	AAIJ	18AW LHCb	$p p$ at 7 and 8 TeV

$$\Gamma(\Sigma_c(2455)^0 p \bar{p}, \Sigma_c^0 \rightarrow \Lambda_c^+ \pi^-) / \Gamma(\Lambda_c^+ p \bar{p} \pi^-) \quad \Gamma_{44}/\Gamma_{43}$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>8.9 ± 1.5 ± 0.6</b>	AAIJ	18AW LHCb	$p p$ at 7 and 8 TeV

$$\Gamma(\Sigma_c(2520)^0 p \bar{p}, \Sigma_c(2520)^0 \rightarrow \Lambda_c^+ \pi^-) / \Gamma(\Lambda_c^+ p \bar{p} \pi^-) \quad \Gamma_{45}/\Gamma_{43}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.119 ± 0.020 ± 0.014</b>	AAIJ	18AW LHCb	$p p$ at 7 and 8 TeV

**$\Gamma(\Lambda K^0 2\pi^+ 2\pi^-)/\Gamma_{\text{total}}$**   **$\Gamma_{46}/\Gamma$**

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
seen	4	<sup>1</sup> ARENTON	86	FMPS $\Lambda K_S^0 2\pi^+ 2\pi^-$

<sup>1</sup> See the footnote to the ARENTON 86 mass value.

**$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}$**   **$\Gamma_{47}/\Gamma$**

The values and averages in this section serve only to show what values result if one assumes our  $B(b \rightarrow b\text{-baryon})$ . They cannot be thought of as measurements since the underlying product branching fractions were also used to determine  $B(b \rightarrow b\text{-baryon})$  as described in the note on “Production and Decay of  $b$ -Flavored Hadrons.”

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.109 \pm 0.022</math> OUR AVERAGE</b>				
$0.102 \pm 0.019 \pm 0.013$		<sup>1</sup> BARATE	98D	ALEP $e^+ e^- \rightarrow Z$
$0.14^{+0.05}_{-0.04} \pm 0.02$	29	<sup>2</sup> ABREU	95S	DLPH $e^+ e^- \rightarrow Z$
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
$0.090 \pm 0.022 \pm 0.012$	55	<sup>3</sup> BUSKULIC	95L	ALEP Repl. by BARATE 98D
$0.18 \pm 0.07 \pm 0.02$	21	<sup>4</sup> BUSKULIC	92E	ALEP $\Lambda_c^+ \rightarrow p K^- \pi^+$

<sup>1</sup> BARATE 98D reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0086 \pm 0.0007 \pm 0.0014$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Measured using  $\Lambda_c \ell^-$  and  $\Lambda \ell^+ \ell^-$ .

<sup>2</sup> ABREU 95S reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.0118 \pm 0.0026^{+0.0031}_{-0.0021}$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>3</sup> BUSKULIC 95L reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.00755 \pm 0.0014 \pm 0.0012$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

<sup>4</sup> BUSKULIC 92E reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell \text{anything})/\Gamma_{\text{total}}] \times [B(\bar{b} \rightarrow b\text{-baryon})] = 0.015 \pm 0.0035 \pm 0.0045$  which we divide by our best value  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. Superseded by BUSKULIC 95L.

**$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/\Gamma_{\text{total}}$**   **$\Gamma_{48}/\Gamma$**

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.062^{+0.014}_{-0.013}</math> OUR FIT</b>			
$0.050^{+0.011+0.016}_{-0.008-0.012}$	<sup>1</sup> ABDALLAH	04A	DLPH $e^+ e^- \rightarrow Z^0$

<sup>1</sup> Derived from a combined likelihood and event rate fit to the distribution of the Isgur-Wise variable and using HQET. The slope of the form factor is measured to be  $\rho^2 = 2.03 \pm 0.46^{+0.72}_{-1.00}$ .

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/\Gamma(\Lambda_c^+ \pi^-)$	$\Gamma_{48}/\Gamma_{32}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>12.8^{+3.0}_{-2.7}</math> OUR FIT</b>			

<b><math>16.6 \pm 3.0^{+2.8}_{-3.6}</math></b>	AALTONEN	09E	CDF	$p\bar{p}$ at 1.96 TeV
--	----------	-----	-----	------------------------

$\Gamma(\Lambda_c^+ \tau^- \bar{\nu}_\tau)/\Gamma(\Lambda_c^+ \pi^+ \pi^- \pi^-)$	$\Gamma_{49}/\Gamma_{37}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.46 \pm 0.27 \pm 0.40</math></b>	1 AAIJ	22K	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Uses  $\tau^- \rightarrow \pi^- \pi^+ \pi^- (\pi^0) \nu_\tau$  decays.

$\Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)/\Gamma_{\text{total}}$	$\Gamma_{50}/\Gamma$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.056^{+0.031}_{-0.030}</math></b>	1 ABDALLAH	04A	DLPH $e^+ e^- \rightarrow Z^0$

<sup>1</sup> Derived from the fraction of  $\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) / (\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)) = 0.47^{+0.10+0.07}_{-0.08-0.06}$ .

$\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)/[\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell) + \Gamma(\Lambda_c^+ \pi^+ \pi^- \ell^- \bar{\nu}_\ell)]$	$\Gamma_{48}/(\Gamma_{48}+\Gamma_{50})$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.47^{+0.10+0.07}_{-0.08-0.06}</math></b>	ABDALLAH	04A	DLPH $e^+ e^- \rightarrow Z^0$

$\Gamma(\Lambda_c(2595)^+ \ell^- \bar{\nu}_\ell)/\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$	$\Gamma_{51}/\Gamma_{48}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.126 \pm 0.033^{+0.047}_{-0.038}</math></b>	1 AALTONEN	09E	CDF $p\bar{p}$ at 1.96 TeV

<sup>1</sup> AALTONEN 09E assumes isospin conservation for  $\Lambda_c(2595) \rightarrow \Lambda_c \pi^+ \pi^+$  and  $\Lambda_c(2595) \rightarrow \Lambda_c \pi^0 \pi^0$ . Significant isospin violation from thresholds in  $\Lambda_c(2595) \rightarrow \Sigma_c(2455) \pi \rightarrow \Lambda_c \pi \pi$  may alter the recovered ratio.

$\Gamma(\Lambda_c(2625)^+ \ell^- \bar{\nu}_\ell)/\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$	$\Gamma_{52}/\Gamma_{48}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.210 \pm 0.042^{+0.071}_{-0.050}</math></b>	AALTONEN	09E	CDF $p\bar{p}$ at 1.96 TeV

$[\frac{1}{2}\Gamma(\Sigma_c(2455)^0 \pi^+ \ell^- \bar{\nu}_\ell) + \frac{1}{2}\Gamma(\Sigma_c(2455)^{++} \pi^- \ell^- \bar{\nu}_\ell)]/\Gamma(\Lambda_c^+ \ell^- \bar{\nu}_\ell)$	$(\frac{1}{2}\Gamma_{53} + \frac{1}{2}\Gamma_{54})/\Gamma_{48}$		
<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.054 \pm 0.022^{+0.021}_{-0.018}</math></b>	AALTONEN	09E	CDF $p\bar{p}$ at 1.96 TeV

$\Gamma(ph^-)/\Gamma_{\text{total}}$	$\Gamma_{55}/\Gamma$		
<u>VALUE</u>	<u>CL %</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
<b><math>&lt;2.3 \times 10^{-5}</math></b>	90	1 ACOSTA	050 CDF $p\bar{p}$ at 1.96 TeV

<sup>1</sup> Assumes  $f_\Lambda / f_d = 0.25$ , and equal momentum distribution for  $\Lambda_b$  and  $B$  mesons.

$\Gamma(p\pi^-)/\Gamma_{\text{total}}$  $\Gamma_{56}/\Gamma$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.5±0.8 OUR FIT</b>				
<b>4.0±0.9±0.5</b>		<sup>1</sup> AALTONEN 09C	CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<50	90	<sup>2</sup> BUSKULIC 96V	ALEP	$e^+e^- \rightarrow Z$
<sup>1</sup> AALTONEN 09C reports $[\Gamma(\Lambda_b^0 \rightarrow p\pi^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.042 \pm 0.007 \pm 0.006$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ , $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ , $B(\bar{b} \rightarrow B^0) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.				
<sup>2</sup> BUSKULIC 96V assumes PDG 96 production fractions for $B^0$ , $B^+$ , $B_s$ , $b$ baryons.				

 $\Gamma(pK^-)/\Gamma_{\text{total}}$  $\Gamma_{57}/\Gamma$ 

<u>VALUE</u> (units $10^{-6}$ )	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>5.4±1.0 OUR FIT</b>				
<b>6.3±1.1±0.8</b>		<sup>1</sup> AALTONEN 09C	CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<360	90	<sup>2</sup> ADAM 96D	DLPH	$e^+e^- \rightarrow Z$
< 50	90	<sup>3</sup> BUSKULIC 96V	ALEP	$e^+e^- \rightarrow Z$
<sup>1</sup> AALTONEN 09C reports $[\Gamma(\Lambda_b^0 \rightarrow pK^-)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^+\pi^-)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.066 \pm 0.009 \pm 0.008$ which we multiply or divide by our best values $B(B^0 \rightarrow K^+\pi^-) = (1.96 \pm 0.05) \times 10^{-5}$ , $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ , $B(\bar{b} \rightarrow B^0) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.				
<sup>2</sup> ADAM 96D assumes $f_{B^0} = f_{B^-} = 0.39$ and $f_{B_s} = 0.12$ .				
<sup>3</sup> BUSKULIC 96V assumes PDG 96 production fractions for $B^0$ , $B^+$ , $B_s$ , $b$ baryons.				

 $\Gamma(p\pi^-)/\Gamma(pK^-)$  $\Gamma_{56}/\Gamma_{57}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.84±0.09 OUR FIT</b>			
<b>0.86±0.08±0.05</b>	AAIJ	12AR LHCb	$p\bar{p}$ at 7 TeV

 $\Gamma(pD_s^-)/\Gamma_{\text{total}}$  $\Gamma_{58}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>&lt;4.8 × 10<sup>-4</sup></b>	90	AAIJ	14Q LHCb	$p\bar{p}$ at 7 TeV

 $\Gamma(p\mu^-\bar{\nu}_\mu)/\Gamma_{\text{total}}$  $\Gamma_{59}/\Gamma$ 

<u>VALUE</u> (units $10^{-4}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.1±1.0</b>	<sup>1</sup> AAIJ	15BG LHCb	$p\bar{p}$ at 8 TeV

<sup>1</sup> The ratio of  $B(\Lambda_b^0 \rightarrow p\mu^-\bar{\nu}_\mu)$  to  $B(\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu)$  is measured within a restricted  $q^2$  region. Combined with theoretical calculations of the form factors and the previously measured value of  $|V_{cb}|$ , the first  $|V_{ub}| = (3.27 \pm 0.15 \pm 0.16 \pm 0.06) \times 10^{-3}$  measurement from the  $\Lambda_b$  decay is obtained, consistent with the exclusively measured world averages.

$\Gamma(p\mu^-\bar{\nu}_\mu)/\Gamma(\Lambda_c^+\ell^-\bar{\nu}_\ell)$   $\Gamma_{59}/\Gamma_{48}$

<u>VALUE</u> (units $10^{-2}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
$1.0 \pm 0.04 \pm 0.08$	<sup>1</sup> AAIJ	15BG LHCb	$p p$ at 8 TeV
<sup>1</sup> This measurement is a ratio of $\Gamma(\Lambda_b^0 \rightarrow p\mu^-\bar{\nu}_\mu)[q^2 > 15 \text{ GeV}/c^2]$ to $\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu)[q^2 > 7 \text{ GeV}/c^2]$ within a restricted $q^2$ region. Combined with theoretical calculations of the form factors and the previously measured value of $ V_{cb} $ , the first $ V_{ub}  = (3.27 \pm 0.15 \pm 0.16 \pm 0.06) \times 10^{-3}$ measurement from the $\Lambda_b$ decay is obtained, consistent with the exclusively measured world averages.			

$\Gamma(\Lambda\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_{60}/\Gamma$

<u>VALUE</u> (units $10^{-7}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>10.8 <math>\pm</math> 2.8 OUR AVERAGE</b>			
$9.6 \pm 1.6 \pm 2.5$	<sup>1</sup> AAIJ	13AJ LHCb	$p p$ at 7 TeV
$17.3 \pm 4.2 \pm 5.5$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
<sup>1</sup> Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$ . This measurement comes from the sum of the differential rates in $q^2$ regions excluding those corresponding to $J/\psi$ and $\psi(2S)$ ([8.68,10.09] and [12.86, 14.18] $\text{GeV}^2/c^4$ ).			

$\Gamma(p\pi^-\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_{61}/\Gamma$

<u>VALUE</u> (units $10^{-8}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>6.9 <math>\pm</math> 1.9 <math>^{+1.7}_{-1.5}</math></b>			
	<sup>1</sup> AAIJ	17P LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> Excludes  $J/\psi$  and  $\psi(2S)$  decays to  $\mu^+\mu^-$ .

$\Gamma(p\pi^-\mu^+\mu^-)/\Gamma(p\pi^-J/\psi, J/\psi \rightarrow \mu^+\mu^-)$   $\Gamma_{61}/\Gamma_{15}$

<u>VALUE</u> (units $10^{-2}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>4.4 <math>\pm</math> 1.2 <math>\pm</math> 0.7</b>			

<sup>1</sup> The  $p\pi^-\mu^+\mu^-$  mode excludes  $J/\psi$  and  $\psi(2S)$  decays to  $\mu^+\mu^-$ .

$\Gamma(pK^-\epsilon^+\epsilon^-)/\Gamma_{\text{total}}$   $\Gamma_{62}/\Gamma$

<u>VALUE</u> (units $10^{-6}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.310 <math>\pm</math> 0.040 <math>^{+0.054}_{-0.047}</math></b>			
	1,2 AAIJ	20M LHCb	$p p$ at 7, 8, 13 TeV

<sup>1</sup> Measured over  $0.1 < q^2 < 6.0 \text{ GeV}/c^2$ , and  $m_{pK} < 2.6 \text{ GeV}/c^2$ .

<sup>2</sup> The first uncertainty is the statistical uncertainty and the second is the combination of all systematic uncertainties including those related to the normalization of  $\Lambda_b^0 \rightarrow J/\psi pK^-$ .

$\Gamma(pK^-\mu^+\mu^-)/\Gamma_{\text{total}}$   $\Gamma_{63}/\Gamma$

<u>VALUE</u> (units $10^{-6}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>0.265 <math>\pm</math> 0.014 <math>^{+0.049}_{-0.039}</math></b>			
	1,2 AAIJ	20M LHCb	$p p$ at 7, 8, 13 TeV

<sup>1</sup> Measured over  $0.1 < q^2 < 6.0 \text{ GeV}/c^2$ , and  $m_{pK} < 2.6 \text{ GeV}/c^2$ .

<sup>2</sup> The first uncertainty is the statistical uncertainty and the second is the combination of all systematic uncertainties including those related to the normalization of  $\Lambda_b^0 \rightarrow J/\psi pK^-$ .

$\Gamma(pK^-\mu^+\mu^-)/\Gamma(pK^-\epsilon^+\epsilon^-)$   $\Gamma_{63}/\Gamma_{62}$ 

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.86^{+0.14}_{-0.11} \pm 0.05</math></b>	<sup>1</sup> AAIJ	20M LHCb	$p\bar{p}$ at 7, 8, 13 TeV

<sup>1</sup> Measured over  $0.1 < q^2 < 6.0 \text{ GeV}/c^2$ , and  $m_{pK} < 2.6 \text{ GeV}/c^2$ .

 $\Gamma(pK^-\epsilon^+\epsilon^-)/\Gamma(pJ/\psi K^-)$   $\Gamma_{62}/\Gamma_{16}$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>9.8^{+1.4}_{-1.3} \pm 0.8</math></b>	<sup>1</sup> AAIJ	20M LHCb	$p\bar{p}$ at 7, 8, 13 TeV

<sup>1</sup> Measured over  $0.1 < q^2 < 6.0 \text{ GeV}/c^2$ , and  $m_{pK} < 2.6 \text{ GeV}/c^2$ .

 $\Gamma(pK^-\mu^+\mu^-)/\Gamma(pJ/\psi K^-)$   $\Gamma_{63}/\Gamma_{16}$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>8.4 \pm 0.4 \pm 0.4</math></b>	<sup>1</sup> AAIJ	20M LHCb	$p\bar{p}$ at 7, 8, 13 TeV

<sup>1</sup> Measured over  $0.1 < q^2 < 6.0 \text{ GeV}/c^2$ , and  $m_{pK} < 2.6 \text{ GeV}/c^2$ .

 $\Gamma(\Lambda\gamma)/\Gamma_{\text{total}}$   $\Gamma_{64}/\Gamma$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>7.1 \pm 1.5 \pm 0.9</math></b>		<sup>1</sup> AAIJ	19Z LHCb	$p\bar{p}$ at 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1300 90 ACOSTA 02G CDF  $p\bar{p}$  at 1.8 TeV

<sup>1</sup> AAIJ 19Z normalized to  $B^0 \rightarrow K^{*0}\gamma$  and used an integrated luminosity of  $1.7 \text{ fb}^{-1}$ .

 $\Gamma(\Lambda\eta)/\Gamma_{\text{total}}$   $\Gamma_{65}/\Gamma$ 

<u>VALUE (units <math>10^{-6}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>9^{+7}_{-5} \pm 1</math></b>	<sup>1</sup> AAIJ	15AH LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 15AH reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda\eta)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \eta' K^0)] = 0.142^{+0.11}_{-0.08}$  which we multiply by our best value  $B(B^0 \rightarrow \eta' K^0) = (6.6 \pm 0.4) \times 10^{-5}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value. The single uncertainty quoted with the original measurement combines in quadrature statistical and systematic uncertainties.

 $\Gamma(\Lambda\eta'(958))/\Gamma_{\text{total}}$   $\Gamma_{66}/\Gamma$ 

<u>VALUE</u>	<u>CL%</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>&lt;3.1 \times 10^{-6}</math></b>	90	<sup>1</sup> AAIJ	15AH LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 15AH reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda\eta'(958))/\Gamma_{\text{total}}] / [B(B^0 \rightarrow \eta' K^0)] < 0.047$  which we multiply by our best value  $B(B^0 \rightarrow \eta' K^0) = 6.6 \times 10^{-5}$ .

 $\Gamma(\Lambda\pi^+\pi^-)/\Gamma(\Lambda_c^+\pi^-)$   $\Gamma_{67}/\Gamma_{32}$ 

<u>VALUE (units <math>10^{-4}</math>)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>9.4 \pm 3.8 \pm 0.4</math></b>	<sup>1</sup> AAIJ	16W LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 16W reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda\pi^+\pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)] / [B(\Lambda_c^+ \rightarrow \Lambda\pi^+)] = (7.3 \pm 1.9 \pm 2.2) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_c^+ \rightarrow \Lambda\pi^+) = (1.29 \pm 0.05) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda K^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$   $\Gamma_{68}/\Gamma_{32}$

VALUE (units $10^{-4}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**11.5±2.3<sup>+0.4</sup><sub>-0.5</sub>** <sup>1</sup> AAIJ 16W LHCb  $p p$  at 7, 8 TeV

<sup>1</sup> AAIJ 16W reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda K^+ \pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow \Lambda \pi^+)] = (8.9 \pm 1.2 \pm 1.3) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_c^+ \rightarrow \Lambda \pi^+) = (1.29 \pm 0.05) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda K^+ K^-)/\Gamma(\Lambda_c^+ \pi^-)$   $\Gamma_{69}/\Gamma_{32}$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**3.27±0.35±0.13** <sup>1</sup> AAIJ 16W LHCb  $p p$  at 7, 8 TeV

<sup>1</sup> AAIJ 16W reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda K^+ K^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow \Lambda \pi^+)] = (25.3 \pm 1.9 \pm 1.9) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_c^+ \rightarrow \Lambda \pi^+) = (1.29 \pm 0.05) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Lambda \phi)/\Gamma_{\text{total}}$   $\Gamma_{70}/\Gamma$

VALUE (units $10^{-6}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**9.8±2.1<sup>+1.6</sup><sub>-1.5</sub>** <sup>1</sup> AAIJ 16J LHCb  $p p$  at 7, 8 TeV

<sup>1</sup> AAIJ 16J reports  $[\Gamma(\Lambda_b^0 \rightarrow \Lambda \phi)/\Gamma_{\text{total}}] / [B(B^0 \rightarrow K^0 \phi)] \times [B(\bar{b} \rightarrow b\text{-baryon})] / [B(\bar{b} \rightarrow B^0)] = 0.275 \pm 0.055 \pm 0.020$  which we multiply or divide by our best values  $B(B^0 \rightarrow K^0 \phi) = (7.3 \pm 0.7) \times 10^{-6}$ ,  $B(\bar{b} \rightarrow b\text{-baryon}) = (8.4 \pm 1.1) \times 10^{-2}$ ,  $B(\bar{b} \rightarrow B^0) = (40.8 \pm 0.7) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best values.

$\Gamma(p \pi^- \pi^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$   $\Gamma_{71}/\Gamma_{32}$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**4.28±0.24<sup>+0.19</sup><sub>-0.20</sub>** <sup>1</sup> AAIJ 18Q LHCb  $p p$  at 7, 8 TeV

<sup>1</sup> AAIJ 18Q reports  $[\Gamma(\Lambda_b^0 \rightarrow p \pi^- \pi^+ \pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = (6.85 \pm 0.19 \pm 0.08 \pm 0.32) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.26 \pm 0.29) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(p K^- K^+ \pi^-)/\Gamma(\Lambda_c^+ \pi^-)$   $\Gamma_{72}/\Gamma_{32}$

VALUE (units $10^{-3}$ )	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	------	---------

**0.83±0.10±0.04** <sup>1</sup> AAIJ 18Q LHCb  $p p$  at 7, 8 TeV

<sup>1</sup> AAIJ 18Q reports  $[\Gamma(\Lambda_b^0 \rightarrow p K^- K^+ \pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)] / [B(\Lambda_c^+ \rightarrow p K^- \pi^+)] = (1.32 \pm 0.09 \pm 0.09 \pm 0.10) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_c^+ \rightarrow p K^- \pi^+) = (6.26 \pm 0.29) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(pK^-\pi^+\pi^-)/\Gamma(\Lambda_c^+\pi^-)$  $\Gamma_{73}/\Gamma_{32}$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>10.3 \pm 0.5 \pm 0.5</math></b>	<sup>1</sup> AAIJ	18Q LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup>AAIJ 18Q reports  $[\Gamma(\Lambda_b^0 \rightarrow pK^-\pi^+\pi^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)] / [B(\Lambda_c^+ \rightarrow pK^-\pi^+)] = (16.4 \pm 0.3 \pm 0.2 \pm 0.7) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (6.26 \pm 0.29) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

 $\Gamma(pK^-K^+K^-)/\Gamma(\Lambda_c^+\pi^-)$  $\Gamma_{74}/\Gamma_{32}$ 

<u>VALUE</u> (units $10^{-3}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.57 \pm 0.15 \pm 0.12</math></b>	<sup>1</sup> AAIJ	18Q LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup>AAIJ 18Q reports  $[\Gamma(\Lambda_b^0 \rightarrow pK^-K^+K^-)/\Gamma(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)] / [B(\Lambda_c^+ \rightarrow pK^-\pi^+)] = (4.11 \pm 0.12 \pm 0.06 \pm 0.19) \times 10^{-2}$  which we multiply by our best value  $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = (6.26 \pm 0.29) \times 10^{-2}$ . Our first error is their experiment's error and our second error is the systematic error from using our best value.

PARTIAL BRANCHING FRACTIONS IN  $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$  $B(\Lambda_b \rightarrow \Lambda\mu^+\mu^-) (q^2 < 2.0 \text{ GeV}^2/c^4)$ 

<u>VALUE</u> (units $10^{-7}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.71 \pm 0.27</math> OUR AVERAGE</b>			

$0.72^{+0.24}_{-0.22} \pm 0.14$  <sup>1</sup>AAIJ 15AE LHCb  $p\bar{p}$  at 7, 8 TeV

$0.15 \pm 2.01 \pm 0.05$  AALTONEN 11AI CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.56 \pm 0.76 \pm 0.80$  <sup>2</sup>AAIJ 13AJ LHCb Repl. by AAIJ 15AE

<sup>1</sup>AAIJ 15AE measurement covers  $0.1 < q^2 < 2.0 \text{ GeV}^2/c^4$ .

<sup>2</sup>Uses  $B(\Lambda_b^0 \rightarrow J/\psi\Lambda) = (6.2 \pm 1.4) \times 10^{-4}$ .

 $B(\Lambda_b \rightarrow \Lambda\mu^+\mu^-) (2.0 < q^2 < 4.3 \text{ GeV}^2/c^4)$ 

<u>VALUE</u> (units $10^{-7}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>0.28^{+0.28}_{-0.21}</math> OUR AVERAGE</b>			

$0.253^{+0.276}_{-0.207} \pm 0.046$  <sup>1</sup>AAIJ 15AE LHCb  $p\bar{p}$  at 7, 8 TeV

$1.8 \pm 1.7 \pm 0.6$  AALTONEN 11AI CDF  $p\bar{p}$  at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.71 \pm 0.60 \pm 0.23$  <sup>2</sup>AAIJ 13AJ LHCb Repl. by AAIJ 15AE

<sup>1</sup>AAIJ 15AE measurement covers  $2.0 < q^2 < 4.0 \text{ GeV}^2/c^4$ .

<sup>2</sup>Uses  $B(\Lambda_b^0 \rightarrow J/\psi\Lambda) = (6.2 \pm 1.4) \times 10^{-4}$ .

 $B(\Lambda_b \rightarrow \Lambda\mu^+\mu^-) (q^2 < 4.3 \text{ GeV}^2/c^4)$ 

<u>VALUE</u> (units $10^{-7}$ )	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b><math>2.7 \pm 2.5 \pm 0.9</math></b>	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV

### $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ( $4.0 < q^2 < 6.0$ GeV $^2/c^4$ )

VALUE (units 10 $^{-7}$ )	DOCUMENT ID	TECN	COMMENT
$0.04^{+0.18}_{-0.00} \pm 0.02$	AAIJ	15AE LHCb	$p\bar{p}$ at 7, 8 TeV

### $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ( $1.0 < q^2 < 6.0$ GeV $^2/c^4$ )

VALUE (units 10 $^{-7}$ )	DOCUMENT ID	TECN	COMMENT
$0.47^{+0.31}_{-0.27}$ OUR AVERAGE			

$0.45^{+0.30}_{-0.25} \pm 0.10$	<sup>1</sup> AAIJ	15AE LHCb	$p\bar{p}$ at 7 and 8 TeV
$1.3 \pm 2.1 \pm 0.4$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV
<sup>1</sup> AAIJ 15AE measurement covers $1.1 < q^2 < 6.0$ GeV $^2/c^4$ .			

### $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ( $6.0 < q^2 < 8.0$ GeV $^2/c^4$ )

VALUE (units 10 $^{-7}$ )	DOCUMENT ID	TECN	COMMENT
$0.50^{+0.24}_{-0.22} \pm 0.10$	AAIJ	15AE LHCb	$p\bar{p}$ at 7, 8 TeV

### $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ( $4.3 < q^2 < 8.68$ GeV $^2/c^4$ )

VALUE (units 10 $^{-7}$ )	DOCUMENT ID	TECN	COMMENT
$0.5 \pm 0.7$ OUR AVERAGE			
$0.66 \pm 0.74 \pm 0.18$	<sup>1</sup> AAIJ	13AJ LHCb	$p\bar{p}$ at 7 TeV
$-0.2 \pm 1.6 \pm 0.1$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV

<sup>1</sup> Uses  $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$ .

### $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ( $10.09 < q^2 < 12.86$ GeV $^2/c^4$ )

VALUE (units 10 $^{-7}$ )	DOCUMENT ID	TECN	COMMENT
$2.2 \pm 0.6$ OUR AVERAGE			
$2.08^{+0.42}_{-0.39} \pm 0.42$	<sup>1</sup> AAIJ	15AE LHCb	$p\bar{p}$ at 7, 8 TeV
$3.0 \pm 1.5 \pm 1.0$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.55 \pm 0.58 \pm 0.55$	<sup>2</sup> AAIJ	13AJ LHCb	Repl. by AAIJ 15AE
--------------------------	-------------------	-----------	--------------------

<sup>1</sup> AAIJ 15AE measurement covers  $11.0 < q^2 < 12.5$  GeV $^2/c^4$ .

<sup>2</sup> Uses  $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$ .

### $B(\Lambda_b \rightarrow \Lambda \mu^+ \mu^-)$ ( $14.18 < q^2 < 16.0$ GeV $^2/c^4$ )

VALUE (units 10 $^{-7}$ )	DOCUMENT ID	TECN	COMMENT
$1.7 \pm 0.5$ OUR AVERAGE	Error includes scale factor of 1.1.		
$2.04^{+0.35}_{-0.33} \pm 0.42$	<sup>1</sup> AAIJ	15AE LHCb	$p\bar{p}$ at 7, 8 TeV
$1.0 \pm 0.7 \pm 0.3$	AALTONEN	11AI CDF	$p\bar{p}$ at 1.96 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$1.44 \pm 0.44 \pm 0.42$	<sup>2</sup> AAIJ	13AJ LHCb	Repl. by AAIJ 15AE
--------------------------	-------------------	-----------	--------------------

<sup>1</sup> AAIJ 15AE measurement covers  $15.0 < q^2 < 16.0$  GeV $^2/c^4$ .

<sup>2</sup> Uses  $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$ .

### $B(\Lambda_b \rightarrow \Lambda\mu^+\mu^-)$ ( $16.0 < q^2 < 20.0$ GeV $^2/c^4$ )

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>7.0 ± 1.9 ± 2.2</b>	AALTENON	11AI CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.73 ± 0.77 ± 1.25	1,2 AAIJ	13AJ LHCb	Repl. by AAIJ 15AE
<sup>1</sup> Uses $B(\Lambda_b^0 \rightarrow J/\psi \Lambda) = (6.2 \pm 1.4) \times 10^{-4}$ .			
<sup>2</sup> Requires $16.00 < q^2 < 20.30$ GeV $^2/c^4$ .			

### $B(\Lambda_b \rightarrow \Lambda\mu^+\mu^-)$ ( $18.0 < q^2 < 20.0$ GeV $^2/c^4$ )

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.44 ± 0.28 ± 0.50</b>	AAIJ	15AE LHCb	$p\bar{p}$ at 7, 8 TeV

### $B(\Lambda_b \rightarrow \Lambda\mu^+\mu^-)$ ( $15.0 < q^2 < 20.0$ GeV $^2/c^4$ )

VALUE (units $10^{-7}$ )	DOCUMENT ID	TECN	COMMENT
<b>6.00 ± 0.45 ± 1.25</b>	AAIJ	15AE LHCb	$p\bar{p}$ at 7, 8 TeV

## CP VIOLATION

$A_{CP}$  is defined as

$$A_{CP} = \frac{B(\Lambda_b^0 \rightarrow f) - B(\bar{\Lambda}_b^0 \rightarrow \bar{f})}{B(\Lambda_b^0 \rightarrow f) + B(\bar{\Lambda}_b^0 \rightarrow \bar{f})},$$

the CP-violation asymmetry of exclusive  $\Lambda_b^0$  and  $\bar{\Lambda}_b^0$  decay.

### $A_{CP}(\Lambda_b \rightarrow p\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.025 ± 0.029 OUR AVERAGE</b>			Error includes scale factor of 1.2.
-0.035 ± 0.017 ± 0.020	AAIJ	18AX LHCb	$p\bar{p}$ at 7 and 8 TeV
0.06 ± 0.07 ± 0.03	AALTENON	14P CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.03 ± 0.17 ± 0.05	AALTENON	11N CDF	Repl. by AALTENON 14P

### $A_{CP}(\Lambda_b \rightarrow pK^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.025 ± 0.022 OUR AVERAGE</b>			
-0.020 ± 0.013 ± 0.019	AAIJ	18AX LHCb	$p\bar{p}$ at 7 and 8 TeV
-0.10 ± 0.08 ± 0.04	AALTENON	14P CDF	$p\bar{p}$ at 1.96 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.37 ± 0.17 ± 0.03	AALTENON	11N CDF	Repl. by AALTENON 14P

### $A_{CP}(\Lambda_b \rightarrow DpK^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.12 ± 0.09 ± 0.03</b>	<sup>1</sup> AAIJ	21AD LHCb	$p\bar{p}$ at 7, 8, 13 TeV

<sup>1</sup>  $A_{CP}$  is measured from  $(B(\Lambda_b^0 \rightarrow [K^+ \pi^-]_D p K^-) - B(\bar{\Lambda}_b^0 \rightarrow [K^- \pi^+]_D \bar{p} K^+)) / (B(\Lambda_b^0 \rightarrow [K^+ \pi^-]_D p K^-) + B(\bar{\Lambda}_b^0 \rightarrow [K^- \pi^+]_D \bar{p} K^+))$  in the full phase space.

### $\Delta A_{CP}(pK^-/\pi^-)$

$$\Delta A_{CP} \equiv A_{CP}(pK^-) - A_{CP}(p\pi^-)$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.014±0.022±0.010</b>	AAIJ	18AX LHCb	$p p$ at 7 and 8 TeV

### $A_{CP}(\Lambda_b \rightarrow p\bar{K}^0\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.22±0.13±0.03</b>	AAIJ	14Q LHCb	$p p$ at 7 TeV

### $\Delta A_{CP}(J/\psi p\pi^-/K^-)$

$$\Delta A_{CP} \equiv A_{CP}(J/\psi p\pi^-) - A_{CP}(J/\psi pK^-)$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.7±2.4±1.2</b>	AAIJ	14K LHCb	$p p$ at 7, 8 TeV

### $A_{CP}(\Lambda_b \rightarrow \Lambda K^+\pi^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.53±0.23±0.11</b>	<sup>1</sup> AAIJ	16W LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> Measured relative to  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$  decay.

### $A_{CP}(\Lambda_b \rightarrow \Lambda K^+ K^-)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.28±0.10±0.07</b>	<sup>1</sup> AAIJ	16W LHCb	$p p$ at 7, 8 TeV

<sup>1</sup> Measured relative to  $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$  decay.

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow pK^-\mu^+\mu^-)$

$$\Delta A_{CP} \equiv A_{CP}(pK^-\mu^+\mu^-) - A_{CP}(pK^- J/\psi)$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>-3.5±5.0±0.2</b>	AAIJ	17T LHCb	$p p$ at 7, 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-)$

$$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p\pi^-\pi^+)\pi^-)$$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.1±2.5±0.6</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Full phase space.

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow (p\pi^-\pi^+\pi^-)_{LBM})$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow (p\pi^-\pi^+\pi^-)_{LBM}) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p\pi^-\pi^+)\pi^-)$ . Two-body low invariant-mass region (LBM):  $m(p\pi^-) < 2000\text{MeV}$  and  $m(\pi^+\pi^-) < 1640\text{ MeV}$ .

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.7±4.1±0.5</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Measurement done with  $m(p\pi^-) < 2000\text{ MeV}/c^2$  and  $m(\pi^+\pi^-) < 1640\text{ MeV}/c^2$ .

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow p a_1(1260)^-)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow p a_1(1260)^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p\pi^- \pi^+) \pi^-)$ . 419  
 $< m(\pi^+ \pi^- \pi^+) < 1500$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>-1.5±4.2±0.6</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow N(1520)^0 \rho(770)^0)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow N(1520)^0 \rho(770)^0) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p\pi^- \pi^+) \pi^-)$ .  
 $1078 < m(p\pi^-) < 1800$  MeV and  $m(\pi^+ \pi^-) < 1100$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.0±4.9±0.4</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow \Delta(1232)^{++} \pi^- \pi^-)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow \Delta(1232)^{++} \pi^- \pi^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p\pi^- \pi^+) \pi^-)$ .  
 $1078 < m(p\pi^+) < 1432$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.1±3.2±0.6</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow p K^- \pi^+ \pi^-)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow p K^- \pi^+ \pi^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.2±1.1±0.6</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Full phase space.

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow (p K^- \pi^+ \pi^-)_{LBM})$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow (p K^- \pi^+ \pi^-)_{LBM}) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$ .  
Two-body low invariant-mass region (LBM):  $m(pK^-) < 2000$  MeV and  $m(\pi^+ \pi^-) < 1640$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>3.5±1.5±0.5</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Measurement done with  $m(pK^-) < 2000$  MeV/ $c^2$  and  $m(\pi^+ \pi^-) < 1640$  MeV/ $c^2$ .

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow N(1520)^0 K^*(892)^0)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow N(1520)^0 K^*(892)^0) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$ .  
 $1078 < m(p\pi^-) < 1800$  MeV and  $750 < m(\pi^+ K^-) < 1100$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>5.5±2.5±0.5</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow \Lambda(1520) \rho(770)^0)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow \Lambda(1520) \rho(770)^0) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$ .  
 $1460 < m(pK^-) < 1580$  MeV and  $m(\pi^+ \pi^-) < 1100$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.6±6.0±0.5</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow \Delta(1232)^{++} K^- \pi^-)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow \Delta(1232)^{++} K^- \pi^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$ .  $1078 < m(p\pi^+) < 1432$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.4±2.6±0.6</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow p K_1(1410)^-)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow p K_1(1410)^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$ .  $1200 < m(K^- \pi^+ \pi^-) < 1600$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.7±3.5±0.8</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow p K^- K^+ \pi^-)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow p K^- K^+ \pi^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p \pi^- \pi^+) \pi^-)$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>-6.9±4.9±0.8</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Full phase space.

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow p K^- K^+ K^-)$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow p K^- K^+ K^-) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>0.2±1.8±0.6</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Full phase space.

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow \Lambda(1520)\phi(1020))$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow \Lambda(1520)\phi(1020)) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$

$1460 < m(pK^-) < 1600$  MeV and  $1005 < m(K^+ K^-) < 1040$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>4.3±5.6±0.4</b>	AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow (p K^-)_{highmass} \phi(1020))$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow (p K^-)_{highmass} \phi(1020)) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$ .  $m(pK^-) > 1600$  MeV and  $1005 < m(K^+ K^-) < 1040$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>-0.7±3.3±0.7</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Measurement done with  $m(pK^-) > 1600$  MeV/c<sup>2</sup>.

### $\Delta A_{CP}(\Lambda_b^0 \rightarrow (p K^- K^+ K^-)_{LBM})$

$\Delta A_{CP} \equiv A_{CP}(\Lambda_b^0 \rightarrow (p K^- K^+ K^-)_{LBM}) - A_{CP}(\Lambda_b^0 \rightarrow (\Lambda_c^+ \rightarrow p K^- \pi^+) \pi^-)$ . Two-body low invariant-mass region (LBM):  $m(pK^-) < 2000$  MeV and  $m(K^+ K^-) < 1675$  MeV.

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>2.7±2.3±0.6</b>	<sup>1</sup> AAIJ	19AH LHCb	$p p$ at 7 and 8 TeV

<sup>1</sup> Measurement done with  $m(pK^-) < 2000$  MeV/c<sup>2</sup> and  $m(K^+ K^-) < 1675$  MeV/c<sup>2</sup>.

## CP AND T VIOLATION PARAMETERS

Measured values of the triple-product asymmetry parameters, odd under time-reversal, are defined as  $A_{c(s)}(\Lambda/\phi) = (N_{c(s)}^+ - N_{c(s)}^-) / (\text{sum})$

where  $N_{c(s)}^+$ ,  $N_{c(s)}^-$  are the number of  $\Lambda$  or  $\phi$  candidates for which the  $\cos(\phi)$  and  $\sin(\phi)$  observables are positive and negative, respectively. Angles  $\cos(\phi)$  and  $\sin(\phi)$  are defined as in LEITNER 07.

### $A_c(\Lambda)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.22±0.12±0.06</b>	AAIJ	16J	LHCb $p\bar{p}$ at 7, 8 TeV

### $A_s(\Lambda)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.13±0.12±0.05</b>	AAIJ	16J	LHCb $p\bar{p}$ at 7, 8 TeV

### $A_c(\phi)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.01±0.12±0.03</b>	AAIJ	16J	LHCb $p\bar{p}$ at 7, 8 TeV

### $A_s(\phi)$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.07±0.12±0.01</b>	AAIJ	16J	LHCb $p\bar{p}$ at 7, 8 TeV

### $a_{CP}(\Lambda_b^0 \rightarrow p\pi^-\pi^+\pi^-)$

Observable calculated as half of the difference between triple products for  $\Lambda_b^0$  and  $\bar{\Lambda}_b^0$ , which is sensitive to CP violation.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-0.7 ±0.7 ±0.2</b>	<sup>1</sup> AAIJ	20AB	LHCb $p\bar{p}$ at 7, 8, 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.15±1.45±0.32      <sup>2</sup> AAIJ      17H      LHCb      Repl. by AAIJ 20AB

<sup>1</sup> Used both triple product asymmetries and the unbinned energy test method.

<sup>2</sup> Measured over full phase space of the decay.

### $a_{CP}(\Lambda_b^0 \rightarrow pK^-\pi^+\pi^-)$

Observable calculated as half of the difference between triple products for  $\Lambda_b^0$  and  $\bar{\Lambda}_b^0$ , which is sensitive to CP violation.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-0.81±0.84±0.31</b>	<sup>1</sup> AAIJ	18AG	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

### $a_{CP}(\Lambda_b^0 \rightarrow pK^-\bar{K}^+\pi^-)$

Observable calculated as half of the difference between triple products for  $\Lambda_b^0$  and  $\bar{\Lambda}_b^0$ , which is sensitive to CP violation.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-0.93±4.54±0.42</b>	<sup>1</sup> AAIJ	17H	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

### $a_{CP}(\Lambda_b^0 \rightarrow p K^- K^+ K^-)$

Observable calculated as half of the difference between triple products for  $\Lambda_b^0$  and  $\bar{\Lambda}_b^0$ , which is sensitive to  $CP$  violation.

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>1.12±1.51±0.32</b>	<sup>1</sup> AAIJ	18AG LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

### $a_{CP}(\Lambda_b^0 \rightarrow p K^- \mu^+ \mu^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>1.2±5.0±0.7</b>	AAIJ	17T LHCb	$p\bar{p}$ at 7, 8 TeV

## P VIOLATION PARAMETERS

Observables calculated as average of the triple products for  $\Lambda_b^0$  and  $\bar{\Lambda}_b^0$ , which is sensitive to parity violation.

### $a_P(\Lambda_b^0 \rightarrow p \pi^- \pi^+ \pi^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-4.0 ±0.7 ±0.2</b>	<sup>1</sup> AAIJ	20AB LHCb	$p\bar{p}$ at 7, 8, 13 TeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$-3.71 \pm 1.45 \pm 0.32$	<sup>2</sup> AAIJ	17H LHCb	Repl. by AAIJ 20AB
---------------------------	-------------------	----------	--------------------

<sup>1</sup> Used both triple product asymmetries and the unbinned energy test method.

<sup>2</sup> Measured over full phase space of the decay.

### $a_P(\Lambda_b^0 \rightarrow p K^- \pi^+ \pi^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-0.60±0.84±0.31</b>	<sup>1</sup> AAIJ	18AG LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

### $a_P(\Lambda_b^0 \rightarrow p K^- K^+ \pi^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>3.62±4.54±0.42</b>	<sup>1</sup> AAIJ	17H LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

### $a_P(\Lambda_b^0 \rightarrow p K^- K^+ K^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-1.56±1.51±0.32</b>	<sup>1</sup> AAIJ	18AG LHCb	$p\bar{p}$ at 7, 8 TeV

<sup>1</sup> Measured over full phase space of the decay.

### $a_P(\Lambda_b^0 \rightarrow p K^- \mu^+ \mu^-)$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
<b>-4.8±5.0±0.7</b>	AAIJ	17T LHCb	$p\bar{p}$ at 7, 8 TeV

## $\Lambda_b^0$ DECAY PARAMETERS

See the note on “Baryon Decay Parameters” in the neutron Listings.

### $\alpha$ decay parameter for $\Lambda_b \rightarrow J/\psi \Lambda$

VALUE	DOCUMENT ID	TECN	COMMENT
<b><math>-0.017 \pm 0.026</math> OUR AVERAGE</b>			

$-0.022^{+0.027}_{-0.026}$	<sup>1</sup> AAIJ	200	LHCb	$p\bar{p}$ at 7, 8, 13 TeV
$-0.14 \pm 0.14 \pm 0.10$	<sup>2</sup> SIRUNYAN	18R	CMS	$p\bar{p}$ at 7, 8 TeV
$0.30 \pm 0.16 \pm 0.06$	<sup>3</sup> AAD	14L	ATLAS	$p\bar{p}$ at 7 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.05 \pm 0.17 \pm 0.07$	<sup>4</sup> AAIJ	13AG	LHCb	Repl. by AAIJ 200

<sup>1</sup> Extracted using a Bayesian analysis. The most probable value is given as  $-0.022$ , with a 68% credibility interval  $[-0.048, 0.005]$ . Transverse polarizations of  $\Lambda_b^0$  of  $-0.004$  (68% credibility interval  $[-0.064, 0.051]$ ),  $0.001$  (68% credibility interval  $[-0.035, 0.045]$ ), and  $0.032$  (68% credibility interval  $[-0.011, 0.065]$ ) are also reported at 7 TeV, 8 TeV and 13 TeV, respectively. Note that both statistical and systematic uncertainties are included.

<sup>2</sup> An angular analysis of  $\Lambda_b \rightarrow J/\psi \Lambda$  decay is performed. Note that the sign of  $\alpha$  in CMS definition is the opposite to that used by AAIJ 13AG and AAD 14L.  $\Lambda_b$  transverse production polarization of  $0.00 \pm 0.06 \pm 0.06$  is also reported, as well as squares of the helicity amplitudes.

<sup>3</sup> An angular analysis of  $\Lambda_b \rightarrow J/\psi \Lambda$  decay is performed and magnitudes of all helicity amplitudes are also reported.

<sup>4</sup> An angular analysis of  $\Lambda_b \rightarrow J/\psi \Lambda$  decay is performed and a  $\Lambda_b$  transverse production polarization of  $0.06 \pm 0.07 \pm 0.02$  is also reported.

### $\alpha_\gamma$ decay parameter for $\Lambda_b \rightarrow \Lambda \gamma$

Measures asymmetry between left- and right-handed photons in the decay.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.82^{+0.17+0.04}_{-0.26-0.13}$	<sup>1</sup> AAIJ	22M	LHCb $p\bar{p}$ at 13 TeV

<sup>1</sup> AAIJ 22M provides a combined measurement as well as measured  $\alpha_\gamma^- = 1.26 \pm 0.42 \pm 0.20$  and  $\alpha_\gamma^+ = 0.55 \pm 0.32 \pm 0.16$  for  $\Lambda_b^0$  and  $\bar{\Lambda}_b^0$  separately.

### $f_L(\mu\mu)$ longitudinal polarization fraction in $\Lambda_b \rightarrow \Lambda \mu^+ \mu^-$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.61^{+0.11}_{-0.14} \pm 0.03$	<sup>1</sup> AAIJ	15AE	LHCb $p\bar{p}$ at 7, 8 TeV

<sup>1</sup> AAIJ 15AE measurement covers  $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$ .

## FORWARD-BACKWARD ASYMMETRIES

The forward-backward assymmetry is defined as  $A_{FB}(\Lambda_b^0) = [N(F) - N(B)] / [N(F) + N(B)]$ , where the forward (F) direction corresponds to a particle ( $\Lambda_b^0$  or  $\bar{\Lambda}_b^0$ ) sharing valence quark flavors with a beam particle with the same sign of rapidity.

### $A_{FB}(\Lambda_b^0 \rightarrow J/\psi \Lambda)$

VALUE	DOCUMENT ID	TECN	COMMENT
$0.04 \pm 0.07 \pm 0.02$	<sup>1</sup> ABAZOV	15I	D0 $p\bar{p}$ at 1.96 TeV

<sup>1</sup> The measured asymmetry integrated over rapidity  $y$  in the range of  $0.1 < |y| < 2.0$ .

### $A_{FB}^\ell(\mu\mu)$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.39±0.04±0.01</b>	<sup>1</sup> AAIJ	18AP LHCb	$p p$ at 7, 8, 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.05±0.09±0.03	<sup>2</sup> AAIJ	15AE LHCb	Repl. by AAIJ 18AP.
<sup>1</sup> The measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$ .			
<sup>2</sup> AAIJ 15AE measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$ .			

### $\Delta(A_{FB}^\ell(\mu\mu))$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$

Difference of asymmetries $A_{FB}^\ell(\mu\mu)$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$ between $\Lambda_b$ and $\bar{\Lambda}_b$ decays	DOCUMENT ID	TECN	COMMENT
<b>-0.05±0.09±0.03</b>	AAIJ	18AO LHCb	$p p$ at 7, 8 TeV

### $A_{FB}^h(p\pi)$ in $\Lambda_b \rightarrow \Lambda(p\pi)\mu^+\mu^-$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>-0.30±0.05±0.02</b>	<sup>1</sup> AAIJ	18AP LHCb	$p p$ at 7, 8, 13 TeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.29±0.07±0.03	<sup>2</sup> AAIJ	15AE LHCb	Repl. by AAIJ 18AP.
<sup>1</sup> The measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$ .			
<sup>2</sup> AAIJ 15AE measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$ .			

### $A_{FB}^{th}$ in $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.25±0.04±0.01</b>	<sup>1</sup> AAIJ	18AP LHCb	$p p$ at 7, 8, 13 TeV
<sup>1</sup> The measurement covers $15.0 < q^2 < 20.0 \text{ GeV}^2/c^4$ .			

## $\Lambda_b^0 - \bar{\Lambda}_b^0$ Production Asymmetry

$$A_P(\Lambda_b^0) = [\sigma(\Lambda_b^0) - \sigma(\bar{\Lambda}_b^0)] / [\sigma(\Lambda_b^0) + \sigma(\bar{\Lambda}_b^0)]$$

### $A_P(\Lambda_b^0)$

VALUE (units $10^{-2}$ )	DOCUMENT ID	TECN	COMMENT
<b>1.4 ±0.4 OUR AVERAGE</b>	Error includes scale factor of 1.8.		
1.92±0.35	<sup>1</sup> AAIJ	21AJ LHCb	$p p$ at 7 TeV
1.09±0.29	<sup>1</sup> AAIJ	21AJ LHCb	$p p$ at 8 TeV
-0.11±2.53±1.08	<sup>2</sup> AAIJ	17BF LHCb	$p p$ at 7 TeV
3.44±1.61±0.76	<sup>2</sup> AAIJ	17BF LHCb	$p p$ at 8 TeV
<sup>1</sup> Integrated over the kinematic range $2 < p_T < 27 \text{ GeV}/c$ and $2.15 < y < 4.10$ .			
<sup>2</sup> Indirect determination in kinematic range $2 < p_T < 30 \text{ GeV}/c$ and $2.1 < \eta < 4.5$ from production asymmetries of $B^+$ , $B^0$ and $B_s^0$ .			

## $\Lambda_b^0$ REFERENCES

AAIJ	22K	PRL 128 191803	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	22M	PR D105 L051104	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	22R	JHEP 2203 153	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	21AD	PR D104 112008	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	21AJ	JHEP 2110 060	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	21B	PL B815 136172	R. Aaij <i>et al.</i>	(LHCb Collab.)

AAIJ	21R	JHEP 2105 095	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20AB	PR D102 051101	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20AK	PR D102 112012	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20M	JHEP 2005 040	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	20O	JHEP 2006 110	R. Aaij <i>et al.</i>	(LHCb Collab.)
SIRUNYAN	20H	PL B802 135203	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	19AH	EPJ C79 745	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19AN	JHEP 1909 028	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19F	JHEP 1903 126	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	19Z	PRL 123 031801	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AF	JHEP 1808 131	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AG	JHEP 1808 039	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AO	JHEP 1809 145 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AP	JHEP 1809 146	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AW	PL B784 101	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18AX	PL B787 124	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	18Q	JHEP 1802 098	R. Aaij <i>et al.</i>	(LHCb Collab.)
PDG	18	PR D98 030001	M. Tanabashi <i>et al.</i>	(PDG Collab.)
SIRUNYAN	18BY	EPJ C78 457	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
SIRUNYAN	18R	PR D97 072010	A.M. Sirunyan <i>et al.</i>	(CMS Collab.)
AAIJ	17AM	PRL 119 062001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17BF	PL B774 139	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17H	NATP 13 391	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17P	JHEP 1704 029	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17S	JHEP 1705 030	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	17T	JHEP 1706 108	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16	JHEP 1601 012	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16A	CP C40 011001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16J	PL B759 282	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16W	JHEP 1605 081	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	16Y	JHEP 1605 132	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAD	15CH	PL B751 63	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	15AE	JHEP 1506 115	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		JHEP 1809 145 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15AH	JHEP 1509 006	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	15BG	NATP 11 743	R. Aaij <i>et al.</i>	(LHCb Collab.)
ABAZOV	15I	PR D91 072008	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AAD	14L	PR D89 092009	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	14AA	PRL 112 202001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14E	JHEP 1404 114	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14H	PR D89 032001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14I	JHEP 1408 143	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14K	JHEP 1407 103	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14Q	JHEP 1404 087	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	14U	PL B734 122	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	14B	PR D89 072014	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	14P	PRL 113 242001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
PDG	14	CP C38 070001	K. Olive <i>et al.</i>	(PDG Collab.)
AAD	13U	PR D87 032002	G. Aad <i>et al.</i>	(ATLAS Collab.)
AAIJ	13AG	PL B724 27	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AJ	PL B725 25	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13AV	PRL 110 182001	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	13BB	PRL 111 102003	R. Aaij <i>et al.</i>	(LHCb Collab.)
CHATRCHYAN	13AC	JHEP 1307 163	S. Chatrchyan <i>et al.</i>	(CMS Collab.)
AAIJ	12AR	JHEP 1210 037	R. Aaij <i>et al.</i>	(LHCb Collab.)
AAIJ	12E	PL B708 241	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	12A	PR D85 032003	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	12U	PR D85 112003	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AAIJ	11E	PR D84 092001	R. Aaij <i>et al.</i>	(LHCb Collab.)
Also		PR D85 039904 (errat.)	R. Aaij <i>et al.</i>	(LHCb Collab.)
AALTONEN	11	PRL 106 121804	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11AI	PRL 107 201802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	11N	PRL 106 181802	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	11O	PR D84 031102	V.M. Abazov <i>et al.</i>	(D0 Collab.)
AALTONEN	10B	PRL 104 102002	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09C	PRL 103 031801	T. Aaltonen <i>et al.</i>	(CDF Collab.)
AALTONEN	09E	PR D79 032001	T. Aaltonen <i>et al.</i>	(CDF Collab.)
ABAZOV	07S	PRL 99 142001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABAZOV	07U	PRL 99 182001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ABULENCIA	07A	PRL 98 122001	A. Abulencia <i>et al.</i>	(FNAL CDF Collab.)
ABULENCIA	07B	PRL 98 122002	A. Abulencia <i>et al.</i>	(FNAL CDF Collab.)

LEITNER	07	NPBPS 174 169	O. Leitner, Z.J. Ajaltouni	
ACOSTA	06	PRL 96 202001	D. Acosta <i>et al.</i>	(CDF Collab.)
ABAZOV	05C	PRL 94 102001	V.M. Abazov <i>et al.</i>	(D0 Collab.)
ACOSTA	05O	PR D72 051104	D. Acosta <i>et al.</i>	(CDF Collab.)
ABDALLAH	04A	PL B585 63	J. Abdallah <i>et al.</i>	(DELPHI Collab.)
ACOSTA	02G	PR D66 112002	D. Acosta <i>et al.</i>	(CDF Collab.)
ABREU	99W	EPJ C10 185	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ACKERSTAFF	98G	PL B426 161	K. Ackerstaff <i>et al.</i>	(OPAL Collab.)
BARATE	98D	EPJ C2 197	R. Barate <i>et al.</i>	(ALEPH Collab.)
ABE	97B	PR D55 1142	F. Abe <i>et al.</i>	(CDF Collab.)
ABE	96M	PRL 77 1439	F. Abe <i>et al.</i>	(CDF Collab.)
ABREU	96D	ZPHY C71 199	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ABREU	96N	PL B374 351	P. Abreu <i>et al.</i>	(DELPHI Collab.)
ADAM	96D	ZPHY C72 207	W. Adam <i>et al.</i>	(DELPHI Collab.)
BUSKULIC	96L	PL B380 442	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
BUSKULIC	96V	PL B384 471	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
PDG	96	PR D54 1	R. M. Barnett <i>et al.</i>	(PDG Collab.)
ABREU	95S	ZPHY C68 375	P. Abreu <i>et al.</i>	(DELPHI Collab.)
AKERS	95K	PL B353 402	R. Akers <i>et al.</i>	(OPAL Collab.)
BUSKULIC	95L	PL B357 685	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ABE	93B	PR D47 2639	F. Abe <i>et al.</i>	(CDF Collab.)
BUSKULIC	92E	PL B294 145	D. Buskulic <i>et al.</i>	(ALEPH Collab.)
ALBAJAR	91E	PL B273 540	C. Albajar <i>et al.</i>	(UA1 Collab.)
BARI	91	NC 104A 1787	G. Bari <i>et al.</i>	(CERN R422 Collab.)
ARENTON	86	NP B274 707	M.W. Arenton <i>et al.</i>	(ARIZ, NDAM, VAND)
BASILE	81	LNC 31 97	M. Basile <i>et al.</i>	(CERN R415 Collab.)