

**$\omega_3(1670)$**  $I^G(J^{PC}) = 0^-(3^{--})$  **$\omega_3(1670)$  MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>1667 ± 4 OUR AVERAGE</b>				
1665.3 ± 5.2 ± 4.5	23400	AMELIN	96	VES $36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1685 ± 20	60	BAUBILLIER	79	HBC $8.2 K^- p$ backward
1673 ± 12	430	1,2 BALTAY	78E	HBC $15 \pi^+ p \rightarrow \Delta 3\pi$
1650 ± 12		CORDEN	78B	OMEG $8-12 \pi^- p \rightarrow N 3\pi$
1669 ± 11	600	2 WAGNER	75	HBC $7 \pi^+ p \rightarrow \Delta^{++} 3\pi$
1678 ± 14	500	DIAZ	74	DBC $6 \pi^+ n \rightarrow p 3\pi^0$
1660 ± 13	200	DIAZ	74	DBC $6 \pi^+ n \rightarrow p \omega \pi^0 \pi^0$
1679 ± 17	200	MATTHEWS	71D	DBC $7.0 \pi^+ n \rightarrow p 3\pi^0$
1670 ± 20		KENYON	69	DBC $8 \pi^+ n \rightarrow p 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1700	110	1 CERRADA	77B	HBC $4.2 K^- p \rightarrow \Lambda 3\pi$
1695 ± 20		BARNES	69B	HBC $4.6 K^- p \rightarrow \omega 2\pi X$
1636 ± 20		ARMENISE	68B	DBC $5.1 \pi^+ n \rightarrow p 3\pi^0$

<sup>1</sup> Phase rotation seen for  $J^P = 3^- \rho \pi$  wave.<sup>2</sup> From a fit to  $I(J^P) = 0(3^-) \rho \pi$  partial wave. **$\omega_3(1670)$  WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>168 ± 10 OUR AVERAGE</b>				
149 ± 19 ± 7	23400	AMELIN	96	VES $36 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
160 ± 80	60	3 BAUBILLIER	79	HBC $8.2 K^- p$ backward
173 ± 16	430	4,5 BALTAY	78E	HBC $15 \pi^+ p \rightarrow \Delta 3\pi$
253 ± 39		CORDEN	78B	OMEG $8-12 \pi^- p \rightarrow N 3\pi$
173 ± 28	600	3,5 WAGNER	75	HBC $7 \pi^+ p \rightarrow \Delta^{++} 3\pi$
167 ± 40	500	DIAZ	74	DBC $6 \pi^+ n \rightarrow p 3\pi^0$
122 ± 39	200	DIAZ	74	DBC $6 \pi^+ n \rightarrow p \omega \pi^0 \pi^0$
155 ± 40	200	3 MATTHEWS	71D	DBC $7.0 \pi^+ n \rightarrow p 3\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
90 ± 20		BARNES	69B	HBC $4.6 K^- p \rightarrow \omega 2\pi$
100 ± 40		KENYON	69	DBC $8 \pi^+ n \rightarrow p 3\pi^0$
112 ± 60		ARMENISE	68B	DBC $5.1 \pi^+ n \rightarrow p 3\pi^0$

<sup>3</sup> Width errors enlarged by us to  $4\Gamma/\sqrt{N}$ ; see the note with the  $K^*(892)$  mass.<sup>4</sup> Phase rotation seen for  $J^P = 3^- \rho \pi$  wave.<sup>5</sup> From a fit to  $I(J^P) = 0(3^-) \rho \pi$  partial wave.

## $\omega_3(1670)$ DECAY MODES

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \rho\pi$	seen
$\Gamma_2 \omega\pi\pi$	seen
$\Gamma_3 b_1(1235)\pi$	possibly seen

## $\omega_3(1670)$ BRANCHING RATIOS

$\Gamma(\omega\pi\pi)/\Gamma(\rho\pi)$	$\Gamma_2/\Gamma_1$
<u>VALUE</u>	<u>EVTS</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$0.71 \pm 0.27$	100    DIAZ    74    DBC $6\pi^+ n \rightarrow p5\pi^0$
$\Gamma(b_1(1235)\pi)/\Gamma(\rho\pi)$	$\Gamma_3/\Gamma_1$
<u>VALUE</u>	<u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
possibly seen	DIAZ    74    DBC $6\pi^+ n \rightarrow p5\pi^0$
$\Gamma(b_1(1235)\pi)/\Gamma(\omega\pi\pi)$	$\Gamma_3/\Gamma_2$
<u>VALUE</u>	<u>CL%</u> <u>DOCUMENT ID</u> <u>TECN</u> <u>COMMENT</u>
• • • We do not use the following data for averages, fits, limits, etc. • • •	
>0.75	68    BAUBILLIER    79    HBC $8.2 K^- p$ backward

## $\omega_3(1670)$ REFERENCES

AMELIN	96	ZPHY C70 71	D.V. Amelin <i>et al.</i>	(SERP, TBIL)
BAUBILLIER	79	PL 89B 131	M. Baubillier <i>et al.</i>	(BIRM, CERN, GLAS+)
BALTAY	78E	PRL 40 87	C. Baltay, C.V. Cautis, M. Kalelkar	(COLU) JP
CORDEN	78B	NP B138 235	M.J. Corden <i>et al.</i>	(BIRM, RHEL, TELA+)
CERRADA	77B	NP B126 241	M. Cerrada <i>et al.</i>	(AMST, CERN, NIJM+) JP
WAGNER	75	PL 58B 201	F. Wagner, M. Tabak, D.M. Chew	(LBL) JP
DIAZ	74	PRL 32 260	J. Diaz <i>et al.</i>	(CASE, CMU)
MATTHEWS	71D	PR D3 2561	J.A.J. Matthews <i>et al.</i>	(TNTO, WISC)
BARNES	69B	PRL 23 142	V.E. Barnes <i>et al.</i>	(BNL)
KENYON	69	PRL 23 146	I.R. Kenyon <i>et al.</i>	(BNL, UCND, ORNL)
ARMENISE	68B	PL 26B 336	N. Armenise <i>et al.</i>	(BARI, BGNA, FIRZ+)