

$\Xi_c(2790)$ $I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$ Status: ***

Seen in $\Xi_c' \pi$ decays. The simplest assignment, based on the mass, width, and decay mode, is that this belongs in the same SU(4) multiplet as the $\Lambda(1405)$ and the $\Lambda_c(2595)^+$, but the spin and parity have not been measured.

 $\Xi_c(2790)$ MASSES

The masses are obtained from the mass-difference measurements that follow.

 $\Xi_c(2790)^+$ MASS

| VALUE (MeV) | DOCUMENT ID |
|--|-------------|
| 2791.9 ± 0.5 OUR FIT | |

 $\Xi_c(2790)^0$ MASS

| VALUE (MeV) | DOCUMENT ID |
|--|-------------|
| 2793.9 ± 0.5 OUR FIT | |

 $\Xi_c(2790) - \Xi_c'$ MASS DIFFERENCES **$m_{\Xi_c(2790)^+} - m_{\Xi_c'}$**

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|-----------------------------------|
| 213.20 ± 0.22 OUR FIT | | | | |
| $213.2 \pm 0.2 \pm 0.1$ | | YELTON | 16 | BELL 2231 and 11,560 evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 211.2 $\pm 1.3 \pm 1.0$ | 18 | CSORNA | 01 | CLEO $e^+ e^- \approx \gamma(4S)$ |

 $m_{\Xi_c(2790)^0} - m_{\Xi_c'}$

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------|------|-----------------------------------|
| 215.70 ± 0.22 OUR FIT | | | | |
| $215.7 \pm 0.2 \pm 0.1$ | | YELTON | 16 | BELL 1241 and 7055 evts |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | |
| 216.2 $\pm 1.3 \pm 1.0$ | 14 | CSORNA | 01 | CLEO $e^+ e^- \approx \gamma(4S)$ |

 $\Xi_c(2790)^+ - \Xi_c(2790)^0$ MASS DIFFERENCE **$m_{\Xi_c(2790)^+} - m_{\Xi_c(2790)^0}$**

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|---|-------------|------|---------|
| -2.0 ± 0.7 OUR FIT | | | |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | |

$-3.3 \pm 0.4 \pm 0.5$ YELTON 16 BELL 2231 and 1241 evts

 $\Xi_c(2790)$ WIDTHS **$\Xi_c(2790)^+$ WIDTH**

| VALUE (MeV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|------|-----------------------------------|
| $8.9 \pm 0.6 \pm 0.8$ | | 2231 | YELTON | 16 | BELL $e^+ e^-$, γ regions |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<15 90 CSORNA 01 CLEO $e^+ e^- \approx \gamma(4S)$

$\Xi_c(2790)^0$ WIDTH

| VALUE (MeV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---------------------|-----|------|-------------|------|-----------------------------------|
| 10.0±0.7±0.8 | | 1241 | YELTON | 16 | BELL $e^+ e^-$, γ regions |

• • • We do not use the following data for averages, fits, limits, etc. • • •

<12 90 CSORNA 01 CLEO $e^+ e^- \approx \gamma(4S)$

$\Xi_c(2790)$ DECAY MODES

| Mode | Fraction (Γ_i/Γ) |
|---------------------------|--------------------------------|
| $\Gamma_1 \Xi_c' \pi$ | seen |
| $\Gamma_2 \Xi_c^0 \gamma$ | |
| $\Gamma_3 \Xi_c^+ \gamma$ | |

$\Xi_c(2790)$ BRANCHING RATIOS

$\Gamma(\Xi_c' \pi)/\Gamma_{\text{total}}$

| VALUE | DOCUMENT ID | TECN | COMMENT | Γ_1/Γ |
|-------------|-------------|------|-----------------------------------|-------------------|
| seen | YELTON | 16 | BELL $e^+ e^-$, γ regions | |
| seen | CSORNA | 01 | CLEO $e^+ e^- \approx \gamma(4S)$ | |

$\Gamma(\Xi_c^0 \gamma)/\Gamma(\Xi_c' \pi)$

| VALUE | EVTS | DOCUMENT ID | TECN | CHG | COMMENT | Γ_2/Γ_1 |
|-----------------------|------|---------------------|------|------|---------|---------------------------|
| 0.13±0.03±0.02 | 401 | ¹ YELTON | 20 | BELL | 0 | $e^+ e^-$ at $\gamma(4S)$ |

¹ Assumes $B(\Xi_c'^+ \rightarrow \Xi_c^+ \gamma) = 100\%$, noting no strong decay of the Ξ_c' is permitted in the available phase space. YELTON 20 measures $B(\Xi_c(2790)^0 \rightarrow \Xi_c^0 \gamma)/B(\Xi_c(2790)^0 \rightarrow \Xi_c'^+ \pi^- \rightarrow \Xi_c^+ \gamma \pi^-)$.

$\Gamma(\Xi_c^+ \gamma)/\Gamma(\Xi_c' \pi)$

| VALUE | CL% | DOCUMENT ID | TECN | CHG | COMMENT | Γ_3/Γ_1 |
|-----------------|-----|---------------------|------|------|---------|---------------------------|
| <0.06 | 90 | ¹ YELTON | 20 | BELL | + | $e^+ e^-$ at $\gamma(4S)$ |

¹ Assumes $B(\Xi_c'^0 \rightarrow \Xi_c^0 \gamma) = 100\%$, noting no strong decay of the Ξ_c' is permitted in the available phase space. YELTON 20 measures $B(\Xi_c(2790)^+ \rightarrow \Xi_c^+ \gamma)/B(\Xi_c(2790)^+ \rightarrow \Xi_c'^0 \pi^+ \rightarrow \Xi_c^0 \gamma \pi^+)$.

$\Xi_c(2790)$ REFERENCES

| | | |
|--------------------------|---------------------------|-----------------|
| YELTON 20 PR D102 071103 | J. Yelton <i>et al.</i> | (BELLE Collab.) |
| YELTON 16 PR D94 052011 | J. Yelton <i>et al.</i> | (BELLE Collab.) |
| CSORNA 01 PRL 86 4243 | S.E. Csorna <i>et al.</i> | (CLEO Collab.) |