

$f_J(2220)$

$I^G(J^{PC}) = 0^+(2^{++} \text{ or } 4^{++})$

OMITTED FROM SUMMARY TABLE

Needs confirmation. See our mini-review in the 2004 edition of this Review, PDG 04.

$f_J(2220)$ MASS

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2231.1 ± 3.5 OUR AVERAGE					
2235	± 4	± 6	74	BAI	96B BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
2230	± 6	± 16	46	BAI	96B BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
2232	± 8	± 15	23	BAI	96B BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
2235	± 4	± 5	32	BAI	96B BES $e^+ e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
2209	+ 17 - 15	± 10		ASTON	88F LASS 11 $K^- p \rightarrow K^+ K^- \Lambda$
2230	± 20		BOLONKIN	88 SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
2220	± 10	41	¹ ALDE	86B GA24	38–100 $\pi p \rightarrow n\eta\eta'$
2230	± 6	± 14	93	BALTRUSAIT..86D	MRK3 $e^+ e^- \rightarrow \gamma K^+ K^-$
2232	± 7	± 7	23	BALTRUSAIT..86D	MRK3 $e^+ e^- \rightarrow \gamma K_S^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2223.9 ± 2.5			² VLADIMIRSK...08	SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n + m\pi^0$
2246	± 36		BAI	98H BES	$J/\psi \rightarrow \gamma\pi^0\pi^0$

¹ ALDE 86B uses data from both the GAMS-2000 and GAMS-4000 detectors.

² $J^{PC} = 2^{++}$. Systematic uncertainties not evaluated

$f_J(2220)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
23 ± 8 OUR AVERAGE					
19 ± 13	± 12	74	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma\pi^+\pi^-$
20 ± 20	± 17	46	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K^+ K^-$
20 ± 25	± 14	23	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma K_S^0 K_S^0$
15 ± 12	± 9	32	BAI	96B BES	$e^+ e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}$
60 ± 107			ASTON	88F LASS	11 $K^- p \rightarrow K^+ K^- \Lambda$
80 ± 30			BOLONKIN	88 SPEC	40 $\pi^- p \rightarrow K_S^0 K_S^0 n$
26 ± 20	± 17	93	BALTRUSAIT..86D	MRK3	$e^+ e^- \rightarrow \gamma K^+ K^-$
18 ± 23	± 10	23	BALTRUSAIT..86D	MRK3	$e^+ e^- \rightarrow \gamma K_S^0 K_S^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.6 ± 2.5	${}^1 \text{VLADIMIRSK...08}$	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
			$+ m\pi^0$
< 80	90	ALDE	$87C \text{ GAM2} \quad 38 \pi^- p \rightarrow \eta' \eta n$

${}^1 JPC = 2^{++}$. Systematic uncertainties not evaluated

$f_J(2220)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi\pi$	not seen
$\Gamma_2 \pi^+ \pi^-$	not seen
$\Gamma_3 K\bar{K}$	not seen
$\Gamma_4 p\bar{p}$	not seen
$\Gamma_5 \gamma\gamma$	not seen
$\Gamma_6 \eta\eta'(958)$	seen
$\Gamma_7 \phi\phi$	not seen
$\Gamma_8 \eta\eta$	not seen

$f_J(2220) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_5/\Gamma$
VALUE (eV)	CL\%
< 1.4	95
${}^1 \text{ACCIARRI}$	$01H \text{ L3}$
$\gamma\gamma \rightarrow K_S^0 K_S^0, E_{\text{cm}}=91, 183-209 \text{ GeV}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
< 5.6	95
${}^1 \text{GODANG}$	97 CLE2
$\gamma\gamma \rightarrow K_S^0 K_S^0$	
< 86	95
${}^1 \text{ALBRECHT}$	$90G \text{ ARG}$
$\gamma\gamma \rightarrow K^+ K^-$	
< 1000	95
${}^2 \text{ALTHOFF}$	$85B \text{ TASS}$
$\gamma\gamma, K\bar{K}\pi$	

$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_1\Gamma_5/\Gamma$
VALUE (eV)	CL\%
< 2.5	95
${}^1 \text{ALAM}$	$98C \text{ CLE2}$
$\gamma\gamma \rightarrow \pi^+ \pi^-$	
1 Assuming $J^P = 2^+$.	
2 True for $J^P = 0^+$ and $J^P = 2^+$.	

$f_J(2220) \Gamma(i)\Gamma(p\bar{p})/\Gamma^2(\text{total})$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\pi\pi)/\Gamma_{\text{total}}$	$\Gamma_4/\Gamma \times \Gamma_1/\Gamma$
$\text{VALUE (units } 10^{-5} \text{)}$	CL\%
< 18	95
${}^1 \text{AMSLER}$	01 CBAR
$1.4-1.5 \text{ } p\bar{p} \rightarrow \pi^0 \pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
$< (11-42)$	99
${}^2 \text{HASAN}$	96 SPEC
$1.35-1.55 \text{ } p\bar{p} \rightarrow \pi^+ \pi^-$	

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\phi\phi)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma \times \Gamma_7/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6	95	3 EVANGELIS...	98	SPEC 1.1-2.0 $p\bar{p} \rightarrow \phi\phi$

$\Gamma(p\bar{p})/\Gamma_{\text{total}} \times \Gamma(\eta\eta)/\Gamma_{\text{total}}$

$\Gamma_4/\Gamma \times \Gamma_8/\Gamma$

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<4	95	1 AMSLER	01	CBAR 1.4-1.5 $p\bar{p} \rightarrow \eta\eta$

¹ For $J^P = 2^+$ in the mass range 2222–2240 MeV and the total width between 10 and 20 MeV.

² For $J^P = 2^+$ and $J^P = 4^+$ in the mass range 2220–2245 MeV and the total width of 15 MeV.

³ For $J^P = 2^+$, the mass of 2235 MeV and the total width of 15 MeV.

$f_J(2220)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$

Γ_1/Γ

VALUE	DOCUMENT ID	COMMENT
not seen	1 DOBBS	$J/\psi \rightarrow \gamma\pi\pi$
not seen	1 DOBBS	$\psi(2S) \rightarrow \gamma\pi\pi$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(K\bar{K})/\Gamma_{\text{total}}$

Γ_3/Γ

VALUE	DOCUMENT ID	COMMENT
not seen	1 DOBBS	$J/\psi \rightarrow \gamma K\bar{K}$
not seen	1 DOBBS	$\psi(2S) \rightarrow \gamma K\bar{K}$

¹ Using CLEO-c data but not authored by the CLEO Collaboration.

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$

Γ_1/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
1.0±0.5	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma 2\pi, K\bar{K}$

$\Gamma(p\bar{p})/\Gamma_{\text{total}}$

Γ_4/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen		1 AUBERT	07AV BABR	$B \rightarrow p\bar{p}K^{(*)}$
not seen		WANG	05A BELL	$B^+ \rightarrow \bar{p}pK^+$
<3.0	95	2 EVANGELIS...	97 SPEC	$1.96-2.40 \bar{p}p \rightarrow K_S^0 K_S^0$
<1.1	99.7	3 BARNES	93 SPEC	$1.3-1.57 \bar{p}p \rightarrow K_S^0 K_S^0$
<2.6	99.7	3 BARDIN	87 CNTR	$1.3-1.5 \bar{p}p \rightarrow K^+K^-$
<3.6	99.7	3 SCULLI	87 CNTR	$1.29-1.55 \bar{p}p \rightarrow K^+K^-$

¹ Assuming $\Gamma < 30$ MeV.

² Assuming $\Gamma \sim 20$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.

³ Assuming $\Gamma = 30-35$ MeV, $J^P = 2^+$ and $B(f_J(2220) \rightarrow K\bar{K}) = 100\%$.

$\Gamma(p\bar{p})/\Gamma(K\bar{K})$

Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.17±0.09	BAI	96B BES	$e^+e^- \rightarrow J/\psi \rightarrow \gamma p\bar{p}, K\bar{K}$

f_J(2220) REFERENCES

DOBBS	15	PR D91 052006	S. Dobbs <i>et al.</i>	(NWES)
VLADIMIRSK...	08	PAN 71 2129	V.V. Vladimirska <i>et al.</i>	(ITEP)
		Translated from YAF 71 2166.		
AUBERT	07AV	PR D76 092004	B. Aubert <i>et al.</i>	(BABAR Collab.)
WANG	05A	PL B617 141	M.-Z. Wang <i>et al.</i>	(BELLE Collab.)
PDG	04	PL B592 1	S. Eidelman <i>et al.</i>	(PDG Collab.)
ACCIARRI	01H	PL B501 173	M. Acciarri <i>et al.</i>	(L3 Collab.)
AMSLER	01	PL B520 175	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ALAM	98C	PRL 81 3328	M.S. Alam <i>et al.</i>	(CLEO Collab.)
BAI	98H	PRL 81 1179	J.Z. Bai <i>et al.</i>	(BES Collab.)
EVANGELIS...	98	PR D57 5370	C. Evangelista <i>et al.</i>	(JETSET Collab.)
EVANGELIS...	97	PR D56 3803	C. Evangelista <i>et al.</i>	(LEAR Collab.)
GODANG	97	PRL 79 3829	R. Godang <i>et al.</i>	(CLEO Collab.)
BAI	96B	PRL 76 3502	J.Z. Bai <i>et al.</i>	(BES Collab.)
HASAN	96	PL B388 376	A. Hasan, D.V. Bugg	(BRUN, LOQM)
BARNES	93	PL B309 469	P.D. Barnes <i>et al.</i>	(PS185 Collab.)
ALBRECHT	90G	ZPHY C48 183	H. Albrecht <i>et al.</i>	(ARGUS Collab.)
ASTON	88F	PL B215 199	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS) JP
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
ALDE	87C	SJNP 45 255	D. Alde <i>et al.</i>	
		Translated from YAF 45 405.		
BARDIN	87	PL B195 292	G. Bardin <i>et al.</i>	(SACL, FERR, CERN, PADO+)
SCULLI	87	PRL 58 1715	J. Sculli <i>et al.</i>	(NYU, BNL)
ALDE	86B	PL B177 120	D.M. Alde <i>et al.</i>	(SERP, BELG, LANL, LAPP)
BALTRUSAIT...	86D	PRL 56 107	R.M. Baltrusaitis	(CIT, UCSC, ILL, SLAC+)
ALTHOFF	85B	ZPHY C29 189	M. Althoff <i>et al.</i>	(TASSO Collab.)

OTHER RELATED PAPERS

DEL-AMO-SA... 10O PRL 105 172001 P. del Amo Sanchez *et al.* (BABAR Collab.)
