

# $f_2(1950)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

## $f_2(1950)$ T-MATRIX POLE $\sqrt{s}$

Note that  $\Gamma \approx 2 \text{Im}(\sqrt{s})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1830–2020) – i (110–220) OUR ESTIMATE</b>			
$(1955 \pm 75) - i (175 \pm 57)$	<sup>1</sup> RODAS	22	RVUE $J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K})$
$(1978.2 \pm 1.8^{+28.4}_{-16.9}) - i$ $(118.8 \pm 0.8^{+20.8}_{-7.8})$	<sup>2</sup> ALBRECHT	20	RVUE $0.9 \bar{p}p \rightarrow \pi^0\pi^0\eta, \pi^0\eta\eta,$ $\pi^0 K^+ K^-$
$(1867 \pm 46) - i (193 \pm 29)$	AMSLER	02	CBAR $0.9 \bar{p}p \rightarrow \pi^0\eta\eta, \pi^0\pi^0\pi^0$
<sup>1</sup> T-matrix pole from coupled channel K-matrix fit to data on $J/\psi \rightarrow \gamma\pi^0\pi^0$ (ABLIKIM 15AE) and $J/\psi \rightarrow \gamma K_S^0 K_S^0$ (ABLIKIM 18AA).			
<sup>2</sup> T-matrix pole, 4 poles, 4 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ).			

## $f_2(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1936 ± 12 OUR AVERAGE</b> Error includes scale factor of 1.3. See the ideogram below.			
1940 ± 50	BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
1980 ± 22	<sup>1</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
1940 ± 22	<sup>2</sup> BARBERIS	00C	450 $pp \rightarrow pp2\pi2\pi^0$
1960 ± 30	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+\pi^-)$
1918 ± 12	ANTINORI	95	OMEG 300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$2038^{+13+12}_{-11-73}$	<sup>3</sup> UEHARA	09	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1930 ± 25	<sup>4</sup> BINON	05	GAMS 33 $\pi^-p \rightarrow \eta\eta n$
1980 ± 2 ± 14	ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
2010 ± 25	ANISOVICH	00J	SPEC
1980 ± 50	ANISOVICH	99B	SPEC 1.35–1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
~ 1990	<sup>5</sup> OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
1950 ± 15	<sup>6</sup> ASTON	91	LASS 11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

<sup>1</sup> Decaying into  $\pi^+\pi^-\pi^0$ .


<sup>2</sup> Decaying into  $2(\pi^+\pi^-)$ .

<sup>3</sup> Taking into account  $f_4(2050)$ .

<sup>4</sup> First solution, PWA is ambiguous.

<sup>5</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

<sup>6</sup> Cannot determine spin to be 2.



### $f_2(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b><math>464 \pm 24</math> OUR AVERAGE</b>			
$380^{+120}_{-90}$	BAI	00A	BES $J/\psi \rightarrow \gamma(\pi^+\pi^-\pi^+\pi^-)$
$520 \pm 50$	<sup>1</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
$485 \pm 55$	<sup>2</sup> BARBERIS	00C	450 $pp \rightarrow pp4\pi$
$460 \pm 40$	BARBERIS	97B	OMEG 450 $pp \rightarrow pp2(\pi^+\pi^-)$
$390 \pm 60$	ANTINORI	95	OMEG 300,450 $pp \rightarrow pp2(\pi^+\pi^-)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
$441^{+27+28}_{-25-192}$	<sup>3</sup> UEHARA	09	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
$450 \pm 50$	<sup>4</sup> BINON	05	GAMS 33 $\pi^-p \rightarrow \eta\eta n$
$297 \pm 12 \pm 6$	ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
$495 \pm 35$	ANISOVICH	00J	SPEC
$500 \pm 100$	ANISOVICH	99B	SPEC 1.35–1.94 $p\bar{p} \rightarrow \eta\eta\pi^0$
$\sim 100$	<sup>5</sup> OAKDEN	94	RVUE 0.36–1.55 $\bar{p}p \rightarrow \pi\pi$
$250 \pm 50$	<sup>6</sup> ASTON	91	LASS 11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$

<sup>1</sup> Decaying into  $\pi^+\pi^-2\pi^0$ .

<sup>2</sup> Decaying into  $2(\pi^+\pi^-)$ .

<sup>3</sup> Taking into account  $f_4(2050)$ .

<sup>4</sup> First solution, PWA is ambiguous.

<sup>5</sup> From solution B of amplitude analysis of data on  $\bar{p}p \rightarrow \pi\pi$ . See however KLOET 96 who fit  $\pi^+\pi^-$  only and find waves only up to  $J = 3$  to be important but not significantly resonant.

<sup>6</sup> Cannot determine spin to be 2.

## $f_2(1950)$ DECAY MODES

	Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1$	$K^*(892)\bar{K}^*(892)$	seen
$\Gamma_2$	$\pi\pi$	
$\Gamma_3$	$\pi^+\pi^-$	seen
$\Gamma_4$	$\pi^0\pi^0$	seen
$\Gamma_5$	$4\pi$	seen
$\Gamma_6$	$\pi^+\pi^-\pi^+\pi^-$	
$\Gamma_7$	$a_2(1320)\pi$	
$\Gamma_8$	$f_2(1270)\pi\pi$	
$\Gamma_9$	$\eta\eta$	seen
$\Gamma_{10}$	$K\bar{K}$	seen
$\Gamma_{11}$	$\gamma\gamma$	seen
$\Gamma_{12}$	$p\bar{p}$	seen

### $f_2(1950)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \qquad \Gamma_{10}\Gamma_{11}/\Gamma$$

<u>VALUE (eV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
-------------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

122 ± 4 ± 26	<sup>1</sup> ABE	04	BELL 10.6 $e^+e^- \rightarrow e^+e^-K^+K^-$
--------------	------------------	----	---

<sup>1</sup> Assuming spin 2.

$$\Gamma(\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}} \qquad \Gamma_2\Gamma_{11}/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

162 <sup>+69</sup> <sub>-42</sub> + 1137 <sub>-204</sub>	<sup>1</sup> UEHARA	09	BELL 10.6 $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
--	---------------------	----	---

<sup>1</sup> Taking into account  $f_4(2050)$ .

### $f_2(1950)$ BRANCHING RATIOS

$$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}} \qquad \Gamma_1/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>CHG</u>	<u>COMMENT</u>
--------------	--------------------	-------------	------------	----------------

seen	ASTON	91	LASS	0 11 $K^-p \rightarrow \Lambda K\bar{K}\pi\pi$
------	-------	----	------	--

$$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}} \qquad \Gamma_7/\Gamma$$

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	--------------------	-------------	----------------

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen	BARBERIS	00B	450 $pp \rightarrow p_f\eta\pi^+\pi^-p_s$
not seen	BARBERIS	00C	450 $pp \rightarrow p_f4\pi p_s$
possibly seen	BARBERIS	97B	OMEG 450 $pp \rightarrow p\rho2(\pi^+\pi^-)$

$\Gamma(\eta\eta)/\Gamma(4\pi)$				$\Gamma_9/\Gamma_5$
VALUE	CL%	DOCUMENT ID	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<5.0 \times 10^{-3}$	90	BARBERIS	00E 450 $pp \rightarrow p_f \eta \eta p_s$	

$\Gamma(\eta\eta)/\Gamma(\pi^+\pi^-)$				$\Gamma_9/\Gamma_3$
VALUE	DOCUMENT ID	TECN	COMMENT	
<b>0.14 ± 0.05</b>	AMSLER	02	CBAR 0.9 $\bar{p}p \rightarrow \pi^0 \eta \eta, \pi^0 \pi^0 \pi^0$	

$\Gamma(p\bar{p})/\Gamma_{total}$				$\Gamma_{12}/\Gamma$
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
<b>seen</b>	111	ALEXANDER	10	CLEO $\psi(2S) \rightarrow \gamma p \bar{p}$

### $f_2(1950)$ REFERENCES

RODAS	22	EPJ C82 80	A. Rodas <i>et al.</i>	(JPAC Collab.)
ALBRECHT	20	EPJ C80 453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
ABLIKIM	18AA	PR D98 072003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ABLIKIM	15AE	PR D92 052003	M. Ablikim <i>et al.</i>	(BESIII Collab.)
ALEXANDER	10	PR D82 092002	J.P. Alexander <i>et al.</i>	(CLEO Collab.)
UEHARA	09	PR D79 052009	S. Uehara <i>et al.</i>	(BELLE Collab.)
BINON	05	PAN 68 960	F. Binon <i>et al.</i>	
		Translated from YAF 68 998.		
ABE	04	EPJ C32 323	K. Abe <i>et al.</i>	(BELLE Collab.)
AMSLER	02	EPJ C23 29	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ANISOVICH	00J	PL B491 47	A.V. Anisovich <i>et al.</i>	(RAL, LOQM, PNPI+)
BAI	00A	PL B472 207	J.Z. Bai <i>et al.</i>	(BES Collab.)
BARBERIS	00B	PL B471 435	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
ANISOVICH	99B	PL B449 154	A.V. Anisovich <i>et al.</i>	
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
KLOET	96	PR D53 6120	W.M. Kloet, F. Myhrer	(RUTG, NORD)
ANTINORI	95	PL B353 589	F. Antinori <i>et al.</i>	(ATHU, BARI, BIRM+) JP
OAKDEN	94	NP A574 731	M.N. Oakden, M.R. Pennington	(DURH)
ASTON	91	NPBPS B21 5	D. Aston <i>et al.</i>	(LASS Collab.)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)