

**$f_0(1370)$**

$I^G(J^{PC}) = 0^+(0^{++})$

See the review on "Spectroscopy of Light Meson Resonances" and  
a note on "Non- $q\bar{q}$  Candidates" in PDG 06, Journal of Physics **G33**  
1 (2006).

### **$f_0(1370)$ T-MATRIX POLE $\sqrt{s}$**

Note that  $\Gamma \approx 2 \operatorname{Im}(\sqrt{s}_{\text{pole}})$ .

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>(1250–1440) –<math>i</math> (60–300) OUR ESTIMATE</b>			
$(1370 \pm 40) - i(195 \pm 20)$	SARANTSEV 21	RVUE	$J/\psi(1S) \rightarrow \gamma (\pi\pi, K\bar{K}, \eta\eta, \omega\phi)$
$(1280.6 \pm 1.6 \pm 47.4) - i(205.2 \pm 1.7 \pm 20.7)$	<sup>1</sup> ALBRECHT 20	RVUE	$0.9 \bar{p}p \rightarrow \pi^0 \pi^0 \eta, \pi^0 \eta\eta, \pi^0 K^+ K^-$
$(1290 \pm 50) - i(170^{+20}_{-40})$	<sup>2</sup> ANISOVICH 09	RVUE	$0.0 \bar{p}p, \pi N$
$(1373 \pm 15) - i(137 \pm 10)$	<sup>3</sup> BARGIOTTI 03	OBLX	$\bar{p}p$
$(1302 \pm 17) - i(166 \pm 18)$	<sup>4</sup> BARBERIS 00C		$450 pp \rightarrow p_f 4\pi p_s$
$(1312 \pm 25 \pm 10) - i(109 \pm 22 \pm 15)$	BARBERIS 99D	OMEG	$450 pp \rightarrow K^+ K^-, \pi^+ \pi^-$
$(1406 \pm 19) - i(80 \pm 6)$	<sup>5</sup> KAMINSKI 99	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, \sigma\sigma$
$(1300 \pm 20) - i(120 \pm 20)$	ANISOVICH 98B	RVUE	Compilation
$(1290 \pm 15) - i(145 \pm 15)$	BARBERIS 97B	OMEG	$450 pp \rightarrow pp 2(\pi^+ \pi^-)$
$(1548 \pm 40) - i(560 \pm 40)$	BERTIN 97C	OBLX	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- \pi^0$
$(1380 \pm 40) - i(180 \pm 25)$	ABELE 96B	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 K_L^0 K_L^0$
$(1300 \pm 15) - i(115 \pm 8)$	BUGG 96	RVUE	
$(1330 \pm 50) - i(150 \pm 40)$	<sup>6</sup> AMSLER 95B	CBAR	$\bar{p}p \rightarrow 3\pi^0$
$(1360 \pm 35) - i(150–300)$	<sup>6</sup> AMSLER 95C	CBAR	$\bar{p}p \rightarrow \pi^0 \eta\eta$
$(1390 \pm 30) - i(190 \pm 40)$	<sup>7</sup> AMSLER 95D	CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta, \pi^0 \pi^0 \eta$
$1346 - i249$	<sup>8,9</sup> JANSEN 95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
$1214 - i168$	<sup>9,10</sup> TORNQVIST 95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
$1364 - i139$	AMSLER 94D	CBAR	$\bar{p}p \rightarrow \pi^0 \pi^0 \eta$
$(1365^{+20}_{-55}) - i(134 \pm 35)$	ANISOVICH 94	CBAR	$\bar{p}p \rightarrow 3\pi^0, \pi^0 \eta\eta$
$(1340 \pm 40) - i(127^{+30}_{-20})$	<sup>11</sup> BUGG 94	RVUE	$\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0, \eta\pi^0 \pi^0$
$(1430 \pm 5) - i(73 \pm 13)$	<sup>12</sup> KAMINSKI 94	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$
$1420 - i220$	<sup>13</sup> AU 87	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}$

<sup>1</sup> T-matrix pole, 5 poles, 5 channels, including scattering data from HYAMS 75 ( $\pi\pi$ ), LONGACRE 86 ( $K\bar{K}$ ), BINON 83 ( $\eta\eta$ ), and BINON 84C ( $\eta\eta'$ ).

<sup>2</sup> Another pole is found at  $(1510 \pm 130) - i(800^{+100}_{-150})$  MeV.

<sup>3</sup> Coupled channel analysis of  $\pi^+ \pi^- \pi^0$ ,  $K^+ K^- \pi^0$ , and  $K^\pm K_S^0 \pi^\mp$ .

<sup>4</sup> Average between  $\pi^+ \pi^- 2\pi^0$  and  $2(\pi^+ \pi^-)$ .

<sup>5</sup> T-matrix pole on sheet — — .

<sup>6</sup> Supersedes ANISOVICH 94.

- 7 Coupled-channel analysis of  $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ , and  $\pi^0\pi^0\eta$  on sheet IV. Demonstrates explicitly that  $f_0(500)$  and  $f_0(1370)$  are two different poles.
- 8 Analysis of data from FALVARD 88.
- 9 The pole is on Sheet III. Demonstrates explicitly that  $f_0(500)$  and  $f_0(1370)$  are two different poles.
- 10 Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.
- 11 Reanalysis of ANISOVICH 94 data.
- 12 T-matrix pole on sheet III.
- 13 Analysis of data from OCHS 73, GRAYER 74, BECKER 79, and CASON 83.

## $f_0(1370)$ BREIT-WIGNER MASS

VALUE (MeV)	DOCUMENT ID
<b>1200 to 1500 OUR ESTIMATE</b>	

### $\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1400 $\pm$ 40		1 AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm\pi^\pm\pi^\mp$
1470 $^{+6+72}_{-7-255}$		2 UEHARA	08A BELL	$10.6 e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
1259 $\pm$ 55	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^-\pi^+\pi^+$
1309 $\pm$ 1 $\pm$ 15		3 BUGG	07A RVUE	$0.0 p\bar{p} \rightarrow 3\pi^0$
1449 $\pm$ 13	4.3k	4 GARMASH	06 BELL	$B^+ \rightarrow K^+\pi^+\pi^-$
1350 $\pm$ 50		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi\pi^+\pi^-$
1265 $\pm$ 30 $^{+20}_{-35}$		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma\pi^+\pi^-K^+K^-$
1434 $\pm$ 18 $\pm$ 9	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^-\pi^+\pi^+$
1308 $\pm$ 10		BARBERIS	99B OMEG	$450 pp \rightarrow p_S p_f \pi^+\pi^-$
1315 $\pm$ 50		BELLAZZINI	99 GAM4	$450 pp \rightarrow pp\pi^0\pi^0$
1315 $\pm$ 30		ALDE	98 GAM4	$100 \pi^-p \rightarrow \pi^0\pi^0n$
1280 $\pm$ 55		BERTIN	98 OBLX	$0.05-0.405 \bar{n}p \rightarrow \pi^+\pi^+\pi^-$
1186	5,6 TORNQVIST	95	RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
1472 $\pm$ 12		ARMSTRONG	91 OMEG	$300 pp \rightarrow pp\pi\pi, ppK\bar{K}$
1275 $\pm$ 20		BREAKSTONE	90 SFM	$62 pp \rightarrow pp\pi^+\pi^-$
1420 $\pm$ 20		AKESSON	86 SPEC	$63 pp \rightarrow pp\pi^+\pi^-$
1256		FROGGATT	77 RVUE	$\pi^+\pi^-$ channel

<sup>1</sup> Breit-Wigner mass.

<sup>2</sup> Breit-Wigner mass. May also be the  $f_0(1500)$ .

<sup>3</sup> Reanalysis of ABELE 96C data.

<sup>4</sup> Also observed by GARMASH 07 in  $B^0 \rightarrow K_S^0\pi^+\pi^-$  decays. Supersedes GARMASH 05.

<sup>5</sup> Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

<sup>6</sup> Also observed by ASNER 00 in  $\tau^- \rightarrow \pi^-\pi^0\pi^0\nu_\tau$  decays

## **$K\bar{K}$ MODE**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
1422 $\pm 15 \pm 28$	<sup>1</sup> AAIJ	19H	LHCb	$p p \rightarrow D^\pm X$
1360 $\pm 31 \pm 28$	430 <sup>2,3</sup> DOBBS	15		$J/\psi \rightarrow \gamma K^+ K^-$
1350 $\pm 48 \pm 15$	168 <sup>2,3</sup> DOBBS	15		$\psi(2S) \rightarrow \gamma K^+ K^-$
1440 $\pm 6$		VLADIMIRSK...06	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1391 $\pm 10$		TIKHOMIROV 03	SPEC	$40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
1440 $\pm 50$		BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1463 $\pm 9$		ETKIN 82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$
1425 $\pm 15$		WICKLUND 80	SPEC	$6 \pi N \rightarrow K^+ K^- N$
$\sim 1300$		POLYCHRO... 79	STRC	$7 \pi^- p \rightarrow n2K_S^0$

<sup>1</sup> From the  $D^\pm \rightarrow K^\pm K^+ K^-$  Dalitz plot fit with the isobar model A.

<sup>2</sup> Using CLEO-c data but not authored by the CLEO Collaboration.

<sup>3</sup> From a fit to a Breit-Wigner line shape with fixed  $\Gamma = 346$  MeV.

## **$4\pi$ MODE $2(\pi\pi)_S + \rho\rho$**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
1395 $\pm 40$		ABELE 01	CBAR	$0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
1374 $\pm 38$		AMSLER 94	CBAR	$0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
1345 $\pm 12$		ADAMO 93	OBLX	$\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
1386 $\pm 30$		GASPERO 93	DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
$\sim 1410$	5751	<sup>1</sup> BETTINI 66	DBC	$0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$

<sup>1</sup>  $\rho\rho$  dominant.

## **$\eta\eta$ MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
1262 $^{+51}_{-78} {}^{+82}_{-103}$	<sup>1</sup> UEHARA 10A	BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
1430	AMSLER 92	CBAR	$0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
1220 $\pm 40$	ALDE 86D	GAM4	$100 \pi^- p \rightarrow n2\eta$

<sup>1</sup> Breit-Wigner mass. May also be the  $f_0(1500)$ .

## **COUPLED CHANNEL MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
1330.2 $^{+5.9}_{-6.5} \pm 5.1$	<sup>1</sup> AAIJ	19H	LHCb $p p \rightarrow D^\pm X$
1306 $\pm 20$	<sup>2</sup> ANISOVICH 03	RVUE	
<sup>1</sup> From the $D^\pm \rightarrow K^\pm K^+ K^-$ Dalitz plot fit with the Triple-M amplitude in the multi-meson model of AOUCHE 18.			
<sup>2</sup> K-matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$ , $\pi^- p \rightarrow K\bar{K} n$ , $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ , $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$ , $\pi^0 \eta\eta$ , $\pi^0 \pi^0 \eta$ , $\pi^+ \pi^- \pi^0$ , $K^+ K^- \pi^0$ , $K_S^0 K_S^0 \pi^0$ , $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$ , $K_S^0 K^- \pi^0$ , $K_S^0 K_S^0 \pi^-$ at rest.			

## **$f_0(1370)$ BREIT-WIGNER WIDTH**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>
<b>200 to 500 OUR ESTIMATE</b>	

### **$\pi\pi$ MODE**

<u>VALUE (MeV)</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
300 ± 80		<sup>1</sup> AUBERT	09L BABR	$B^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp$
90 + 2 + 50 1 - 22		<sup>2</sup> UEHARA	08A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$
298 ± 21	2.6k	BONVICINI	07 CLEO	$D^+ \rightarrow \pi^- \pi^+ \pi^+$
126 ± 25	4286	<sup>3</sup> GARMASH	06 BELL	$B^+ \rightarrow K^+ \pi^+ \pi^-$
265 ± 40		ABLIKIM	05 BES2	$J/\psi \rightarrow \phi \pi^+ \pi^-$
350 ± 100 + 105 - 60		ABLIKIM	05Q BES2	$\psi(2S) \rightarrow \gamma \pi^+ \pi^- K^+ K^-$
173 ± 32 ± 6	848	AITALA	01A E791	$D_s^+ \rightarrow \pi^- \pi^+ \pi^+$
222 ± 20		BARBERIS	99B OMEG	$450 pp \rightarrow p_S p_f \pi^+ \pi^-$
255 ± 60		BELLAZZINI	99 GAM4	$450 pp \rightarrow p p \pi^0 \pi^0$
190 ± 50		ALDE	98 GAM4	$100 \pi^- p \rightarrow \pi^0 \pi^0 n$
323 ± 13		BERTIN	98 OBLX	$0.05-0.405 \bar{n} p \rightarrow \pi^+ \pi^+ \pi^-$
350		<sup>4,5</sup> TORNQVIST	95 RVUE	$\pi\pi \rightarrow \pi\pi, K\bar{K}, K\pi, \eta\pi$
195 ± 33		ARMSTRONG	91 OMEG	$300 pp \rightarrow p p \pi\pi, p p K\bar{K}$
285 ± 60		BREAKSTONE	90 SFM	$62 pp \rightarrow p p \pi^+ \pi^-$
460 ± 50		AKESSON	86 SPEC	$63 pp \rightarrow p p \pi^+ \pi^-$
~ 400		<sup>6</sup> FROGGATT	77 RVUE	$\pi^+ \pi^-$ channel

<sup>1</sup> The systematic errors are not reported.

<sup>2</sup> Breit-Wigner width. May also be the  $f_0(1500)$ .

<sup>3</sup> Also observed by GARMASH 07 in  $B^0 \rightarrow K_S^0 \pi^+ \pi^-$  decays. Supersedes GARMASH 05.

<sup>4</sup> Uses data from BEIER 72B, OCHS 73, HYAMS 73, GRAYER 74, ROSSELET 77, CASON 83, ASTON 88, and ARMSTRONG 91B. Coupled channel analysis with flavor symmetry and all light two-pseudoscalars systems.

<sup>5</sup> Also observed by ASNER 00 in  $\tau^- \rightarrow \pi^- \pi^0 \pi^0 \nu_\tau$  decays

<sup>6</sup> Width defined as distance between 45 and 135° phase shift.

### **$K\bar{K}$ MODE**

<u>VALUE (MeV)</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
324 ± 38 ± 42		<sup>1</sup> AAIJ	19H LHCb $pp \rightarrow D^\pm X$
121 ± 15		VLADIMIRSK...06	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
55 ± 26		TIKHOMIROV 03	SPEC $40.0 \pi^- C \rightarrow K_S^0 K_S^0 K_L^0 X$
250 ± 80		BOLONKIN 88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$
118 + 138 - 16		ETKIN 82B MPS	$23 \pi^- p \rightarrow n 2 K_S^0$
160 ± 30		WICKLUND 80	SPEC $6 \pi N \rightarrow K^+ K^- N$
~ 150		POLYCHRO... 79 STRC	$7 \pi^- p \rightarrow n 2 K_S^0$

<sup>1</sup> From the  $D^\pm \rightarrow K^\pm K^+ K^-$  Dalitz plot fit with the isobar model A.

**$4\pi$  MODE  $2(\pi\pi)_S + \rho\rho$** 

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>				
275 $\pm$ 55		ABELE	01	CBAR $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$
375 $\pm$ 61		AMSLER	94	CBAR $0.0 \bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
398 $\pm$ 26		ADAMO	93	OBIX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
310 $\pm$ 50		GASPERO	93	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
$\sim 90$	5751	<sup>1</sup> BETTINI	66	DBC $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$
$^1 \rho\rho$ dominant.				

 **$\eta\eta$  MODE**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
$484^{+246}_{-170}{}^{+246}_{-263}$	<sup>1</sup> UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
250	AMSLER	92	CBAR $0.0 \bar{p}p \rightarrow \pi^0 \eta\eta$
320 $\pm$ 40	ALDE	86D GAM4	$100 \pi^- p \rightarrow n 2\eta$

<sup>1</sup> Breit-Wigner width. May also be the  $f_0(1500)$ .

**COUPLED CHANNEL MODE**

VALUE (MeV)	DOCUMENT ID	TECN	
<b>• • • We do not use the following data for averages, fits, limits, etc. • • •</b>			
$147^{+30}_{-50}$	<sup>1</sup> ANISOVICH	03 RVUE	
$^1 K$ -matrix pole from combined analysis of $\pi^- p \rightarrow \pi^0 \pi^0 n$ , $\pi^- p \rightarrow K\bar{K}n$ , $\pi^+ \pi^- \rightarrow \pi^+ \pi^-$ , $\bar{p}p \rightarrow \pi^0 \pi^0 \pi^0$ , $\pi^0 \eta\eta$ , $\pi^0 \pi^0 \eta$ , $\pi^+ \pi^- \pi^0$ , $K^+ K^- \pi^0$ , $K_S^0 K_S^0 \pi^0$ , $K^+ K_S^0 \pi^-$ at rest, $\bar{p}n \rightarrow \pi^- \pi^- \pi^+$ , $K_S^0 K^- \pi^0$ , $K_S^0 K_S^0 \pi^-$ at rest.			

 **$f_0(1370)$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \pi\pi$	seen
$\Gamma_2 4\pi$	seen
$\Gamma_3 4\pi^0$	seen
$\Gamma_4 2\pi^+ 2\pi^-$	seen
$\Gamma_5 \pi^+ \pi^- 2\pi^0$	seen
$\Gamma_6 \rho\rho$	seen
$\Gamma_7 2(\pi\pi)_S$ -wave	seen
$\Gamma_8 \pi(1300)\pi$	seen
$\Gamma_9 a_1(1260)\pi$	seen
$\Gamma_{10} \eta\eta$	seen
$\Gamma_{11} K\bar{K}$	seen
$\Gamma_{12} K\bar{K}n\pi$	not seen
$\Gamma_{13} 6\pi$	not seen
$\Gamma_{14} \omega\omega$	not seen

$\Gamma_{15}$	$\gamma\gamma$	seen
$\Gamma_{16}$	$e^+ e^-$	not seen

## $f_0(1370)$ PARTIAL WIDTHS

### $\Gamma(\gamma\gamma)$

See  $\gamma\gamma$  widths under  $f_0(500)$  and MORGAN 90.

$\Gamma_{15}$

### $\Gamma(e^+ e^-)$

$\Gamma_{16}$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<20	90	VOROBYEV	88	$e^+ e^- \rightarrow \pi^0 \pi^0$

## $f_0(1370) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

### $\Gamma(\eta\eta) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$

$\Gamma_{10}\Gamma_{15}/\Gamma$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$121^{+133+169}_{-53-106}$	<sup>1</sup> UEHARA	10A BELL	$10.6 e^+ e^- \rightarrow e^+ e^- \eta\eta$
----------------------------	---------------------	----------	---

<sup>1</sup> Including interference with the  $f'_2(1525)$  (parameters fixed to the values from the 2008 edition of this review, PDG 08) and  $f_2(1270)$ . May also be the  $f_0(1500)$ .

## $f_0(1370)$ BRANCHING RATIOS

### $\Gamma(\pi\pi)/\Gamma_{\text{total}}$

$\Gamma_1/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.10	95	OCHS	13	RVUE
$0.26 \pm 0.09$		BUGG	96	RVUE
<0.15	<sup>1</sup> AMSLER		94	CBAR $\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
<0.06		GASPERO	93	DBC $0.0 \bar{p}n \rightarrow$ hadrons

<sup>1</sup> Using AMSLER 95B ( $3\pi^0$ ).

### $\Gamma(4\pi)/\Gamma_{\text{total}}$

$\Gamma_2/\Gamma = (\Gamma_3 + \Gamma_4 + \Gamma_5)/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.72		GASPERO	93	DBC $0.0 \bar{p}n \rightarrow$ hadrons
-------	--	---------	----	--

### $\Gamma(4\pi^0)/\Gamma(4\pi)$

$\Gamma_3/\Gamma_2$

VALUE	DOCUMENT ID	TECN	COMMENT
-------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ABELE	96	CBAR $0.0 \bar{p}p \rightarrow 5\pi^0$
$0.068 \pm 0.005$	<sup>1</sup> GASPERO	93	DBC $0.0 \bar{p}n \rightarrow$ hadrons

<sup>1</sup> Model-dependent evaluation.

$\Gamma(2\pi^+ 2\pi^-)/\Gamma(4\pi)$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.420 \pm 0.014$  <sup>1</sup> GASPERO 93 DBC  $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$

<sup>1</sup> Model-dependent evaluation.

$\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3 + \Gamma_4 + \Gamma_5)$

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.420 \pm 0.014$  <sup>1</sup> GASPERO 93 DBC  $0.0 \bar{p}n \rightarrow 2\pi^+ 3\pi^-$

<sup>1</sup> Model-dependent evaluation.

$\Gamma(\pi^+ \pi^- 2\pi^0)/\Gamma(4\pi)$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.512 \pm 0.019$  <sup>1</sup> GASPERO 93 DBC  $0.0 \bar{p}n \rightarrow \text{hadrons}$

<sup>1</sup> Model-dependent evaluation.

$\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3 + \Gamma_4 + \Gamma_5)$

DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.512 \pm 0.019$  <sup>1</sup> GASPERO 93 DBC  $0.0 \bar{p}n \rightarrow \text{hadrons}$

$\Gamma(\rho\rho)/\Gamma(4\pi)$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.26 \pm 0.07$  ABELE 01B CBAR  $0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma_6/\Gamma_2$

$\Gamma(2(\pi\pi)_S\text{-wave})/\Gamma(\pi\pi)$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.6 \pm 2.6$  <sup>1</sup> ABELE 01 CBAR  $0.0 \bar{p}d \rightarrow \pi^- 4\pi^0 p$

<sup>1</sup> From the combined data of ABELE 96 and ABELE 96c.

$\Gamma_7/\Gamma_1$

$\Gamma(2(\pi\pi)_S\text{-wave})/\Gamma(4\pi)$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.51 \pm 0.09$  ABELE 01B CBAR  $0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma_7/\Gamma_2$

$\Gamma(\rho\rho)/\Gamma(2(\pi\pi)_S\text{-wave})$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

large	BARBERIS	00C	$450 pp \rightarrow p_f 4\pi p_s$
$1.6 \pm 0.2$	AMSLER	94	$\bar{p}p \rightarrow \pi^+ \pi^- 3\pi^0$
$\sim 0.65$	GASPERO	93	$0.0 \bar{p}n \rightarrow \text{hadrons}$

$\Gamma_6/\Gamma_7$

$\Gamma(\pi(1300)\pi)/\Gamma(4\pi)$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.17 \pm 0.06$  ABELE 01B CBAR  $0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma_8/\Gamma_2$

$\Gamma(a_1(1260)\pi)/\Gamma(4\pi)$

VALUE

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.06 \pm 0.02$  ABELE 01B CBAR  $0.0 \bar{p}d \rightarrow 5\pi p$

$\Gamma_9/\Gamma_2$

### $\Gamma(\eta\eta)/\Gamma(4\pi)$

VALUE

DOCUMENT ID

### $\Gamma_{10}/\Gamma_2 = \Gamma_{10}/(\Gamma_3 + \Gamma_4 + \Gamma_5)$

TECN

COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(28 \pm 11) \times 10^{-3}$   
 $(4.7 \pm 2.0) \times 10^{-3}$

<sup>1</sup> ANISOVICH 02D SPEC Combined fit  
BARBERIS 00E 450  $p\bar{p} \rightarrow p_f \eta\eta p_s$

<sup>1</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$ ,  $\pi^0\eta\eta$ ,  $\pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n$ ,  $\eta\eta n$ ,  $\eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

### $\Gamma(K\bar{K})/\Gamma_{\text{total}}$

VALUE

DOCUMENT ID

TECN

### $\Gamma_{11}/\Gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.35 \pm 0.13$

BUGG 96 RVUE

### $\Gamma(K\bar{K})/\Gamma(\pi\pi)$

VALUE

DOCUMENT ID

TECN

### $\Gamma_{11}/\Gamma_1$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.08 \pm 0.08$

ABLIKIM 05 BES2  $J/\psi \rightarrow \phi\pi^+\pi^-$ ,  $\phi K^+K^-$

$0.91 \pm 0.20$

<sup>1</sup> BARGIOTTI 03 OBLX  $\bar{p}p$

$0.12 \pm 0.06$

<sup>2</sup> ANISOVICH 02D SPEC Combined fit

$0.46 \pm 0.15 \pm 0.11$

BARBERIS 99D OMEG 450  $p\bar{p} \rightarrow K^+K^-$ ,  $\pi^+\pi^-$

<sup>1</sup> Coupled channel analysis of  $\pi^+\pi^-\pi^0$ ,  $K^+K^-\pi^0$ , and  $K^\pm K_S^0\pi^\mp$ .

<sup>2</sup> From a combined K-matrix analysis of Crystal Barrel (0.  $p\bar{p} \rightarrow \pi^0\pi^0\pi^0$ ,  $\pi^0\eta\eta$ ,  $\pi^0\pi^0\eta$ ), GAMS ( $\pi p \rightarrow \pi^0\pi^0 n$ ,  $\eta\eta n$ ,  $\eta\eta' n$ ), and BNL ( $\pi p \rightarrow K\bar{K} n$ ) data.

### $\Gamma(K\bar{K}\eta\pi)/\Gamma_{\text{total}}$

VALUE

DOCUMENT ID

TECN

### $\Gamma_{12}/\Gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.03$

GASPERO 93 DBC 0.0  $\bar{p}n \rightarrow$  hadrons

### $\Gamma(6\pi)/\Gamma_{\text{total}}$

VALUE

DOCUMENT ID

TECN

### $\Gamma_{13}/\Gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.22$

GASPERO 93 DBC 0.0  $\bar{p}n \rightarrow$  hadrons

### $\Gamma(\omega\omega)/\Gamma_{\text{total}}$

VALUE

DOCUMENT ID

TECN

### $\Gamma_{14}/\Gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<0.13$

GASPERO 93 DBC 0.0  $\bar{p}n \rightarrow$  hadrons

## $f_0(1370)$ REFERENCES

SARANTSEV	21	PL B816	136227	A.V. Sarantsev <i>et al.</i>	(BONN, PNPI)
ALBRECHT	20	EPJ C80	453	M. Albrecht <i>et al.</i>	(Crystal Barrel Collab.)
AAIJ	19H	JHEP	1904 063	R. Aaij <i>et al.</i>	(LHCb Collab.)
AOUDE	18	PR D98	056021	R.T. Aoude <i>et al.</i>	
DOBBS	15	PR D91	052006	S. Dobbs <i>et al.</i>	(NWES)
OCHS	13	JP G40	043001	W. Ochs	
UEHARA	10A	PR D82	114031	S. Uehara <i>et al.</i>	(BELLE Collab.)
ANISOVICH	09	IJMP A24	2481	V.V. Anisovich, A.V. Sarantsev	(PNPI)
AUBERT	09L	PR D79	072006	B. Aubert <i>et al.</i>	(BABAR Collab.)

PDG	08	PL B667 1	C. Amsler <i>et al.</i>	(PDG Collab.)
UEHARA	08A	PR D78 052004	S. Uehara <i>et al.</i>	(BELLE Collab.)
BONVICINI	07	PR D76 012001	G. Bonvicini <i>et al.</i>	(CLEO Collab.)
BUGG	07A	JP G34 151	D.V. Bugg <i>et al.</i>	
GARMASH	07	PR D75 012006	A. Garmash <i>et al.</i>	(BELLE Collab.)
GARMASH	06	PRL 96 251803	A. Garmash <i>et al.</i>	(BELLE Collab.)
PDG	06	JP G33 1	W.-M. Yao <i>et al.</i>	(PDG Collab.)
Vladimirsk...	06	PAN 69 493	V.V. Vladimirsy <i>et al.</i>	(ITEP, Moscow)
		Translated from YAF 69 515.		
ABLIKIM	05	PL B607 243	M. Ablikim <i>et al.</i>	(BES Collab.)
ABLIKIM	05Q	PR D72 092002	M. Ablikim <i>et al.</i>	(BES Collab.)
GARMASH	05	PR D71 092003	A. Garmash <i>et al.</i>	(BELLE Collab.)
ANISOVICH	03	EPJ A16 229	V.V. Anisovich <i>et al.</i>	
BARGIOTTI	03	EPJ C26 371	M. Bargiotti <i>et al.</i>	(OBELIX Collab.)
TIKHOMIROV	03	PAN 66 828	G.D. Tikhomirov <i>et al.</i>	
		Translated from YAF 66 860.		
ANISOVICH	02D	PAN 65 1545	V.V. Anisovich <i>et al.</i>	
		Translated from YAF 65 1583.		
ABELE	01	EPJ C19 667	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	01B	EPJ C21 261	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
AITALA	01A	PRL 86 765	E.M. Aitala <i>et al.</i>	(FNAL E791 Collab.)
ASNER	00	PR D61 012002	D.M. Asner <i>et al.</i>	(CLEO Collab.)
BARBERIS	00C	PL B471 440	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	00E	PL B479 59	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BARBERIS	99B	PL B453 316	D. Barberis <i>et al.</i>	(Omega Expt.)
BARBERIS	99D	PL B462 462	D. Barberis <i>et al.</i>	(Omega Expt.)
BELLAZZINI	99	PL B467 296	R. Bellazzini <i>et al.</i>	
KAMINSKI	99	EPJ C9 141	R. Kaminski, L. Lesniak, B. Loiseau	(CRAC, PARIN)
ALDE	98	EPJ A3 361	D. Alde <i>et al.</i>	(GAM4 Collab.)
Also		PAN 62 405	D. Alde <i>et al.</i>	(GAMS Collab.)
		Translated from YAF 62 446.		
ANISOVICH	98B	SPU 41 419	V.V. Anisovich <i>et al.</i>	
		Translated from UFN 168 481.		
BERTIN	98	PR D57 55	A. Bertin <i>et al.</i>	(OBELIX Collab.)
BARBERIS	97B	PL B413 217	D. Barberis <i>et al.</i>	(WA 102 Collab.)
BERTIN	97C	PL B408 476	A. Bertin <i>et al.</i>	(OBELIX Collab.)
ABELE	96	PL B380 453	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96B	PL B385 425	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
ABELE	96C	NP A609 562	A. Abele <i>et al.</i>	(Crystal Barrel Collab.)
BUGG	96	NP B471 59	D.V. Bugg, A.V. Sarantsev, B.S. Zou	(LOQM, PNPI)
AMSLER	95B	PL B342 433	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95C	PL B353 571	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
AMSLER	95D	PL B355 425	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
JANSSEN	95	PR D52 2690	G. Janssen <i>et al.</i>	(STON, ADLD, JULI)
TORNQVIST	95	ZPHY C68 647	N.A. Tornqvist	(HELS)
AMSLER	94	PL B322 431	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC
AMSLER	94D	PL B333 277	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.) JPC
ANISOVICH	94	PL B323 233	V.V. Anisovich <i>et al.</i>	(Crystal Barrel Collab.) JPC
BUGG	94	PR D50 4412	D.V. Bugg <i>et al.</i>	(LOQM)
KAMINSKI	94	PR D50 3145	R. Kaminski, L. Lesniak, J.P. Maillet	(CRAC+)
ADAMO	93	NP A558 13C	A. Adamo <i>et al.</i>	(OBELIX Collab.) JPC
GASPERO	93	NP A562 407	M. Gaspero	(ROMAI) JPC
AMSLER	92	PL B291 347	C. Amsler <i>et al.</i>	(Crystal Barrel Collab.)
ARMSTRONG	91	ZPHY C51 351	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
ARMSTRONG	91B	ZPHY C52 389	T.A. Armstrong <i>et al.</i>	(ATHU, BARI, BIRM+)
BREAKSTONE	90	ZPHY C48 569	A.M. Breakstone <i>et al.</i>	(ISU, BGNA, CERN+)
MORGAN	90	ZPHY C48 623	D. Morgan, M.R. Pennington	(RAL, DURH)
ASTON	88	NP B296 493	D. Aston <i>et al.</i>	(SLAC, NAGO, CINC, INUS)
BOLONKIN	88	NP B309 426	B.V. Bolonkin <i>et al.</i>	(ITEP, SERP)
FALVARD	88	PR D38 2706	A. Falvard <i>et al.</i>	(CLER, FRAS, LAZO+)
VOROBIEV	88	SJNP 48 273	P.V. Vorobiev <i>et al.</i>	(NOVO)
		Translated from YAF 48 436.		
AU	87	PR D35 1633	K.L. Au, D. Morgan, M.R. Pennington	(DURH, RAL)
AKESSON	86	NP B264 154	T. Akesson <i>et al.</i>	(Axial Field Spec. Collab.)
ALDE	86D	NP B269 485	D.M. Alde <i>et al.</i>	(BELG, LAPP, SERP, CERN+)
LONGACRE	86	PL B177 223	R.S. Longacre <i>et al.</i>	(BNL, BRAN, CUNY+)
BINON	84C	NC 80A 363	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
BINON	83	NC 78A 313	F.G. Binon <i>et al.</i>	(BELG, LAPP, SERP+)
CASON	83	PR D28 1586	N.M. Cason <i>et al.</i>	(NDAM, ANL)
ETKIN	82B	PR D25 1786	A. Etkin <i>et al.</i>	(BNL, CUNY, TUFTS, VAND)
WICKLUND	80	PRL 45 1469	A.B. Wicklund <i>et al.</i>	(ANL)

BECKER	79	NP B151 46	H. Becker <i>et al.</i>	(MPIM, CERN, ZEEM, CRAC)
POLYCHRO...	79	PR D19 1317	V.A. Polychronakos <i>et al.</i>	(NDAM, ANL)
FROGGATT	77	NP B129 89	C.D. Froggatt, J.L. Petersen	(GLAS, NORD)
ROSSELET	77	PR D15 574	L. Rosselet <i>et al.</i>	(GEVA, SACL)
HYAMS	75	NP B100 205	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
GRAYER	74	NP B75 189	G. Grayer <i>et al.</i>	(CERN, MPIM)
HYAMS	73	NP B64 134	B.D. Hyams <i>et al.</i>	(CERN, MPIM)
OCHS	73	Thesis	W. Ochs	(MPIM, MUNI)
BEIER	72B	PRL 29 511	E.W. Beier <i>et al.</i>	(PENN)
BETTINI	66	NC 42A 695	A. Bettini <i>et al.</i>	(PADO, PISA)

---