

$I(J^P) = 1(\frac{1}{2}^+)$ Status: *** *

We have omitted some results that have been superseded by later experiments. See our earlier editions.

Σ^+ MASS

The fit uses Σ^+ , Σ^0 , Σ^- , and Λ mass and mass-difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-------------	------	-------------	------	---------

1189.37±0.07 OUR FIT Error includes scale factor of 2.2.

1189.37±0.06 OUR AVERAGE Error includes scale factor of 1.8. See the ideogram below.


1189.33±0.04	607	¹ BOHM	72	EMUL
1189.16±0.12		HYMAN	67	HEBC
1189.61±0.08	4205	SCHMIDT	65	HBC See note with Λ mass
1189.48±0.22	58	² BHOWMIK	64	EMUL
1189.38±0.15	144	² BARKAS	63	EMUL

¹ BOHM 72 is updated with our 1973 K^- , π^- , and π^0 masses (Reviews of Modern Physics **45** S1 (1973)).

² These masses have been raised 30 keV to take into account a 46 keV increase in the proton mass and a 21 keV decrease in the π^0 mass (note added 1967 edition, Reviews of Modern Physics **39** 1 (1967)).

WEIGHTED AVERAGE

1189.37±0.06 (Error scaled by 1.8)



Σ^+ MEAN LIFE

Measurements with fewer than 1000 events have been omitted.

<u>VALUE</u> (10^{-10} s)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.8018±0.0026 OUR AVERAGE				
0.8038±0.0040±0.0014		BARBOSA 00	E761	hyperons, 375 GeV
0.8043±0.0080±0.0014	¹	BARBOSA 00	E761	hyperons, 375 GeV
0.798 ± 0.005	30k	MARRAFFINO 80	HBC	$K^- p$ 0.42–0.5 GeV/c
0.807 ± 0.013	5719	CONFORTO 76	HBC	$K^- p$ 1–1.4 GeV/c
0.795 ± 0.010	20k	EISELE 70	HBC	$K^- p$ at rest
0.803 ± 0.008	10664	BARLOUTAUD 69	HBC	$K^- p$ 0.4–1.2 GeV/c
0.83 ± 0.032	1300	² CHANG 66	HBC	

¹ This is a measurement of the $\bar{\Sigma}^-$ lifetime. Here we assume *CPT* invariance; see below for the fractional $\Sigma^+ - \bar{\Sigma}^-$ lifetime difference obtained by BARBOSA 00.

² We have increased the CHANG 66 error of 0.018; see our 1970 edition, *Reviews of Modern Physics* **42** 87 (1970).

$$(\tau_{\Sigma^+} - \tau_{\bar{\Sigma}^-}) / \tau_{\Sigma^+}$$

A test of *CPT* invariance.

<u>VALUE</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
$(-6 \pm 12) \times 10^{-4}$	BARBOSA 00	E761	hyperons, 375 GeV


Σ^+ MAGNETIC MOMENT

See the “Quark Model” review. Measurements with an error $\geq 0.1 \mu_N$ have been omitted.

<u>VALUE</u> (μ_N)	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.458 ±0.010 OUR AVERAGE Error includes scale factor of 2.1. See the ideogram below.				
2.4613±0.0034±0.0040	250k	MORELOS 93	SPEC	p Cu 800 GeV
2.428 ± 0.036 ± 0.007	12k	¹ MORELOS 93	SPEC	p Cu 800 GeV
2.479 ± 0.012 ± 0.022	137k	WILKINSON 87	SPEC	p Be 400 GeV
2.4040±0.0198	44k	² ANKENBRA... 83	CNTR	p Cu 400 GeV

¹ We assume *CPT* invariance: this is (minus) the $\bar{\Sigma}^-$ magnetic moment as measured by MORELOS 93. See below for the moment difference testing *CPT*.

² ANKENBRANDT 83 gives the value $2.38 \pm 0.02 \mu_N$. MORELOS 93 uses the same hyperon magnet and channel and claims to determine the field integral better, leading to the revised value given here.



$$(\mu_{\Sigma^+} + \mu_{\Sigma^-}) / \mu_{\Sigma^+}$$

A test of *CPT* invariance.

VALUE	DOCUMENT ID	TECN	COMMENT
0.014±0.015	¹ MORELOS 93	SPEC	p Cu 800 GeV

¹ This is our calculation from the MORELOS 93 measurements of the Σ^+ and $\bar{\Sigma}^-$ magnetic moments given above. The statistical error on $\mu_{\bar{\Sigma}^-}$ dominates the error here.

Σ^+ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 p\pi^0$	$(51.57 \pm 0.30) \%$	
$\Gamma_2 n\pi^+$	$(48.31 \pm 0.30) \%$	
$\Gamma_3 p\gamma$	$(1.23 \pm 0.05) \times 10^{-3}$	
$\Gamma_4 n\pi^+\gamma$	[a] $(4.5 \pm 0.5) \times 10^{-4}$	
$\Gamma_5 \Lambda e^+ \nu_e$	$(2.0 \pm 0.5) \times 10^{-5}$	

$\Delta S = \Delta Q$ (*SQ*) violating modes or
 $\Delta S = 1$ weak neutral current (*S1*) modes

$\Gamma_6 ne^+ \nu_e$	<i>SQ</i>	< 5	$\times 10^{-6}$	90%
$\Gamma_7 n\mu^+ \nu_\mu$	<i>SQ</i>	< 3.0	$\times 10^{-5}$	90%
$\Gamma_8 pe^+ e^-$	<i>S1</i>	< 7	$\times 10^{-6}$	
$\Gamma_9 p\mu^+ \mu^-$	<i>S1</i>	$(2.4 \pm 1.7) \times 10^{-8}$		

[a] See the Listings below for the pion momentum range used in this measurement.

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 14 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 7.7$ for 12 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \cdot \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_2 & -100 & \\ \hline x_3 & 12 & -14 \\ \hline & x_1 & x_2 \end{array}$$

Σ^+ BRANCHING RATIOS

$\Gamma(n\pi^+)/\Gamma(N\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	$\Gamma_2 / (\Gamma_1 + \Gamma_2)$
0.4836 ± 0.0030 OUR FIT				
0.4836 ± 0.0030 OUR AVERAGE				
0.4828 ± 0.0036	10k	¹ MARRAFFINO 80	HBC	$K^- p$ 0.42–0.5 GeV/c
0.488 ± 0.008	1861	NOWAK	78	HBC
0.484 ± 0.015	537	TOVEE	71	EMUL
0.488 ± 0.010	1331	BARLOUTAUD	69	HBC $K^- p$ 0.4–1.2 GeV/c
0.46 ± 0.02	534	CHANG	66	HBC
0.490 ± 0.024	308	HUMPHREY	62	HBC

¹ MARRAFFINO 80 actually gives $\Gamma(p\pi^0)/\Gamma(\text{total}) = 0.5172 \pm 0.0036$.

$\Gamma(p\gamma)/\Gamma(p\pi^0)$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	Γ_3 / Γ_1
2.38 ± 0.10 OUR FIT				
2.38 ± 0.10 OUR AVERAGE				
2.32 ± 0.11 ± 0.10	32k	TIMM	95	E761 Σ^+ 375 GeV
2.81 ± 0.39 ^{+0.21} _{-0.43}	408	HESSEY	89	CNTR $K^- p \rightarrow \Sigma^+ \pi^-$ at rest
2.52 ± 0.28	190	¹ KOBAYASHI	87	CNTR $\pi^+ p \rightarrow \Sigma^+ K^+$
2.46 ^{+0.30} _{-0.35}	155	BIAGI	85	CNTR CERN hyperon beam
2.11 ± 0.38	46	MANZ	80	HBC $K^- p \rightarrow \Sigma^+ \pi^-$
2.1 ± 0.3	45	ANG	69B	HBC $K^- p$ at rest
2.76 ± 0.51	31	GERSHWIN	69B	HBC $K^- p \rightarrow \Sigma^+ \pi^-$
3.7 ± 0.8	24	BAZIN	65	HBC $K^- p$ at rest

¹ KOBAYASHI 87 actually gives $\Gamma(p\gamma)/\Gamma(\text{total}) = (1.30 \pm 0.15) \times 10^{-3}$.

$\Gamma(n\pi^+\gamma)/\Gamma(n\pi^+)$ Γ_4/Γ_2

The π^+ momentum cuts differ, so we do not average the results but simply use the latest value in the Summary Table.

<u>VALUE</u> (units 10^{-3})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
0.93±0.10	180	EBENHOH	73	HBC $\pi^+ < 150 \text{ MeV}/c$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.27±0.05	29	ANG	69B	HBC $\pi^+ < 110 \text{ MeV}/c$
~1.8		BAZIN	65B	HBC $\pi^+ < 116 \text{ MeV}/c$

 $\Gamma(\Lambda e^+\nu_e)/\Gamma_{\text{total}}$ Γ_5/Γ

<u>VALUE</u> (units 10^{-5})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
2.0±0.5 OUR AVERAGE				
1.6±0.7	5	BALTAY	69	HBC $K^- p$ at rest
2.9±1.0	10	EISELE	69	HBC $K^- p$ at rest
2.0±0.8	6	BARASH	67	HBC $K^- p$ at rest

 $\Gamma(n e^+\nu_e)/\Gamma(n\pi^+)$ Γ_6/Γ_2

Test of $\Delta S = \Delta Q$ rule. Experiments with an effective denominator less than 100,000 have been omitted.

<u>EFFECTIVE DENOM.</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
< 1.1 × 10⁻⁵ OUR LIMIT				Our 90% CL limit = (2.3 events)/(effective denominator sum). [Number of events increased to 2.3 for a 90% confidence level.]
111000	0	¹ EBENHOH	74	HBC $K^- p$ at rest
105000	0	¹ SECHI-ZORN	73	HBC $K^- p$ at rest

¹ Effective denominator calculated by us.

 $\Gamma(n\mu^+\nu_\mu)/\Gamma(n\pi^+)$ Γ_7/Γ_2

Test of $\Delta S = \Delta Q$ rule.

<u>EFFECTIVE DENOM.</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
< 6.2 × 10⁻⁵ OUR LIMIT			Our 90% CL limit = (6.7 events)/(effective denominator sum). [Number of events increased to 6.7 for a 90% confidence level.]
33800	0	BAGGETT	69B HBC
62000	2	¹ EISELE	69B HBC
10150	0	² COURANT	64 HBC
1710	0	² NAUENBERG	64 HBC
120	1	GALTIERI	62 EMUL

¹ Effective denominator calculated by us.

² Effective denominator taken from EISELE 67.

 $\Gamma(pe^+e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

<u>VALUE</u> (units 10^{-6})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
<7		¹ ANG	69B	HBC $K^- p$ at rest

¹ ANG 69B found three pe^+e^- events in agreement with $\gamma \rightarrow e^+e^-$ conversion from $\Sigma^+ \rightarrow p\gamma$. The limit given here is for neutral currents.

$\Gamma(p\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_9/Γ

A test for a $\Delta S = 1$ weak neutral current, but also allowed by higher-order electroweak interactions.

<u>VALUE</u> (units 10^{-8})	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
---------------------------------	-------------	--------------------	-------------	----------------

2.4^{+1.7}_{-1.3} OUR AVERAGE

$2.2^{+0.9+1.5}_{-0.8-1.1}$	10.2	¹ AAIJ	18E	LHCb $p p$ at 7, 8 TeV
$8.6^{+6.6}_{-5.4}\pm 5.5$	3	² PARK	05	HYCP p Cu, 800 GeV

¹AAIJ 18E sees no structure in the dimuon mass distribution, contrary to PARK 05.

²The masses of the three dimuons of PARK 05 are within 1 MeV of one another, perhaps indicating the existence of a new state P^0 with mass 214.3 ± 0.5 MeV. In that case, the decay is $\Sigma^+ \rightarrow p P^0$, $P^0 \rightarrow \mu^+ \mu^-$, with a branching fraction of $(3.1^{+2.4}_{-1.9} \pm 1.5) \times 10^{-8}$.

 $\Gamma(\Sigma^+ \rightarrow n e^+ \nu_e)/\Gamma(\Sigma^- \rightarrow n e^- \bar{\nu}_e)$ $\Gamma_6/\Gamma_3^{\Sigma^-}$

<u>VALUE</u>	<u>CL%</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	------------	-------------	--------------------	-------------	----------------

<0.009 OUR LIMIT Our 90% CL limit, using $\Gamma(n e^+ \nu_e)/\Gamma(n \pi^+)$ above.

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.019	90	0	EBENHOH	74	HBC $K^- p$ at rest
<0.018	90	0	SECHI-ZORN	73	HBC $K^- p$ at rest
<0.12	95	0	COLE	71	HBC $K^- p$ at rest
<0.03	90	0	EISELE	69B	HBC See EBENHOH 74

 $\Gamma(\Sigma^+ \rightarrow n \mu^+ \nu_\mu)/\Gamma(\Sigma^- \rightarrow n \mu^- \bar{\nu}_\mu)$ $\Gamma_7/\Gamma_4^{\Sigma^-}$

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>	<u>COMMENT</u>
--------------	-------------	--------------------	-------------	----------------

<0.12 OUR LIMIT Our 90% CL limit, using $\Gamma(n \mu^+ \nu_\mu)/\Gamma(n \pi^+)$ above.

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.06^{+0.045}_{-0.03}$	2	EISELE	69B	HBC $K^- p$ at rest
-------------------------	---	--------	-----	---------------------

 $\Gamma(\Sigma^+ \rightarrow n \ell^+ \nu)/\Gamma(\Sigma^- \rightarrow n \ell^- \bar{\nu}_\ell)$ $(\Gamma_6 + \Gamma_7)/(\Gamma_3^{\Sigma^-} + \Gamma_4^{\Sigma^-})$

Test of $\Delta S = \Delta Q$ rule.

<u>VALUE</u>	<u>EVTS</u>	<u>DOCUMENT ID</u>	<u>TECN</u>
--------------	-------------	--------------------	-------------

<0.043 OUR LIMIT Our 90% CL limit, using $[\Gamma(\Sigma^+ \rightarrow n \mu^+ \nu_\mu) + \Gamma(\Sigma^+ \rightarrow n e^+ \nu_e)]/\Gamma(\Sigma^+ \rightarrow n \pi^+)$.

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.08	1	NORTON	69	HBC
<0.034	0	BAGGETT	67	HBC

Σ^+ DECAY PARAMETERS

See the “Note on Baryon Decay Parameters” in the neutron Listings. A few early results have been omitted.

α_0 FOR $\Sigma^+ \rightarrow p\pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.982 ± 0.014 OUR FIT				
$-0.982^{+0.016}_{-0.013}$ OUR AVERAGE				
–0.998±0.037±0.009	93k	¹ ABLIKIM	20X	BES3 $J/\psi/\psi(2S) \rightarrow \Sigma^+\bar{\Sigma}^-$
–0.945 ^{+0.055} _{–0.042}	1259	² LIPMAN	73	OSPK $\pi^+ p \rightarrow \Sigma^+$
–0.940±0.045	16k	BELLAMY	72	ASPK $\pi^+ p \rightarrow \Sigma^+ K^+$
–0.98 ^{+0.05} _{–0.02}	1335	³ HARRIS	70	OSPK $\pi^+ p \rightarrow \Sigma^+ K^+$
–0.999±0.022	32k	BANGERTER	69	HBC $K^- p$ 0.4 GeV/c

¹ ABLIKIM 20X uses production through $e^+e^- \rightarrow J/\psi \rightarrow \Sigma^+\bar{\Sigma}^-$ and $e^+e^- \rightarrow \psi(2S) \rightarrow \Sigma^+\bar{\Sigma}^-$, with 87,815 and 5,327 events, respectively. Note that the reported values of decay parameters α_0 of Σ^+ and $\bar{\alpha}_0$ of $\bar{\Sigma}^-$ are correlated.

² Decay protons scattered off aluminum.

³ Decay protons scattered off carbon.

$\bar{\alpha}_0$ FOR $\bar{\Sigma}^- \rightarrow \bar{p}\pi^0$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.990 \pm 0.037 \pm 0.011$	93k	¹ ABLIKIM	20X	BES3 $J/\psi/\psi(2S) \rightarrow \Sigma^+\bar{\Sigma}^-$

¹ ABLIKIM 20X uses production through $e^+e^- \rightarrow J/\psi \rightarrow \Sigma^+\bar{\Sigma}^-$ and $e^+e^- \rightarrow \psi(2S) \rightarrow \Sigma^+\bar{\Sigma}^-$, with 87,815 and 5,327 events, respectively. Note that the reported values of decay parameters α_0 of Σ^+ and $\bar{\alpha}_0$ of $\bar{\Sigma}^-$ are correlated.

$(\alpha_0 + \bar{\alpha}_0) / (\alpha_0 - \bar{\alpha}_0)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.004 \pm 0.037 \pm 0.010$	93k	¹ ABLIKIM	20X	BES3 $J/\psi/\psi(2S) \rightarrow \Sigma^+\bar{\Sigma}^-$

¹ ABLIKIM 20X uses production through $e^+e^- \rightarrow J/\psi \rightarrow \Sigma^+\bar{\Sigma}^-$ and $e^+e^- \rightarrow \psi(2S) \rightarrow \Sigma^+\bar{\Sigma}^-$, with 87,815 and 5,327 events, respectively. Note that the reported values of decay parameters α_0 of Σ^+ and $\bar{\alpha}_0$ of $\bar{\Sigma}^-$ are correlated.

ϕ_0 ANGLE FOR $\Sigma^+ \rightarrow p\pi^0$

$$(\tan \phi_0 = \beta/\gamma)$$

VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
36 ± 34 OUR AVERAGE				
38.1 ^{+35.7} _{–37.1}	1259	¹ LIPMAN	73	OSPK $\pi^+ p \rightarrow \Sigma^+ K^+$
22 ± 90		² HARRIS	70	OSPK $\pi^+ p \rightarrow \Sigma^+ K^+$

¹ Decay proton scattered off aluminum.

² Decay protons scattered off carbon.

α_+ / α_0

Older results have been omitted.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.069 ± 0.013 OUR FIT				
-0.073 ± 0.021	23k	MARRAFFINO	80	HBC $K^- p$ 0.42–0.5 GeV/c

α_+ FOR $\Sigma^+ \rightarrow n\pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.068±0.013 OUR FIT				
0.066±0.016 OUR AVERAGE				
0.037±0.049	4101	BERLEY	70B	HBC
0.069±0.017	35k	BANGERTER	69	HBC $K^- p$ 0.4 GeV/c

ϕ_+ ANGLE FOR $\Sigma^+ \rightarrow n\pi^+$

$$(\tan \phi_+ = \beta/\gamma)$$

VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
167±20 OUR AVERAGE Error includes scale factor of 1.1.				
184±24	1054	¹ BERLEY	70B	HBC
143±29	560	BANGERTER	69B	HBC $K^- p$ 0.4 GeV/c

¹ Changed from 176 to 184° to agree with our sign convention.

α_γ FOR $\Sigma^+ \rightarrow p\gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.76 ±0.08 OUR AVERAGE				
-0.720±0.086±0.045	35k	¹ FOUCHER	92	SPEC $\Sigma^+ 375$ GeV
-0.86 ±0.13 ±0.04	190	KOBAYASHI	87	CNTR $\pi^+ p \rightarrow \Sigma^+ K^+$
-0.53 +0.38 -0.36	46	MANZ	80	HBC $K^- p \rightarrow \Sigma^+ \pi^-$
-1.03 +0.52 -0.42	61	GERSHWIN	69B	HBC $K^- p \rightarrow \Sigma^+ \pi^-$

¹ See TIMM 95 for a detailed description of the analysis.

Σ^+ REFERENCES

We have omitted some papers that have been superseded by later experiments. See our earlier editions.

ABLIKIM	20X	PRL 125 052004	M. Ablikim <i>et al.</i>	(BESIII Collab.)
AAIJ	18E	PRL 120 221803	R. Aaij <i>et al.</i>	(LHCb Collab.)
PARK	05	PRL 94 021801	H.K. Park <i>et al.</i>	(FNAL HyperCP Collab.)
BARBOSA	00	PR D61 031101	R.F. Barbosa <i>et al.</i>	(FNAL E761 Collab.)
TIMM	95	PR D51 4638	S. Timm <i>et al.</i>	(FNAL E761 Collab.)
MORELOS	93	PRL 71 3417	A. Morelos <i>et al.</i>	(FNAL E761 Collab.)
FOUCHER	92	PRL 68 3004	M. Foucher <i>et al.</i>	(FNAL E761 Collab.)
HESSEY	89	ZPHY C42 175	N.P. Hessey <i>et al.</i>	(BNL-811 Collab.)
KOBAYASHI	87	PRL 59 868	M. Kobayashi <i>et al.</i>	(KYOT)
WILKINSON	87	PRL 58 855	C.A. Wilkinson <i>et al.</i>	(WISC, MICH, RUTG+)
BIAGI	85	ZPHY C28 495	S.F. Biagi <i>et al.</i>	(CERN WA62 Collab.)
ANKENBRA...	83	PRL 51 863	C.M. Ankenbrandt <i>et al.</i>	(FNAL, IOWA, ISU+)
MANZ	80	PL 96B 217	A. Manz <i>et al.</i>	(MPIM, VAND)
MARRAFFINO	80	PR D21 2501	J. Marraffino <i>et al.</i>	(VAND, MPIM)
NOWAK	78	NP B139 61	R.J. Nowak <i>et al.</i>	(LOUC, BELG, DURH+)
CONFORTO	76	NP B105 189	B. Conforto <i>et al.</i>	(RHEL, LOIC)
EBENHOH	74	ZPHY 266 367	H. Ebenhoh <i>et al.</i>	(HEIDT)
EBENHOH	73	ZPHY 264 413	W. Ebenhoh <i>et al.</i>	(HEIDT)
LIPMAN	73	PL 43B 89	N.H. Lipman <i>et al.</i>	(RHEL, SUSS, LOWC)
PDG	73	RMP 45 S1	T.A. Lasinski <i>et al.</i>	(LBL, BRAN, CERN+)
SECHI-ZORN	73	PR D8 12	B. Sechi-Zorn, G.A. Snow	(UMD)
BELLAMY	72	PL 39B 299	E.H. Bellamy <i>et al.</i>	(LOWC, RHEL, SUSS)
BOHM	72	NP B48 1	G. Bohm <i>et al.</i>	(BERL, KIDR, BRUX, IASD+)
Also		IIHE-73.2 Nov	G. Bohm	(BERL, KIDR, BRUX, IASD, DUUC+)
COLE	71	PR D4 631	J. Cole <i>et al.</i>	(STON, COLU)
TOVEE	71	NP B33 493	D.N. Tovee <i>et al.</i>	(LOUC, KIDR, BERL+)
BERLEY	70B	PR D1 2015	D. Berley <i>et al.</i>	(BNL, MASA, YALE)
EISELE	70	ZPHY 238 372	F. Eisele <i>et al.</i>	(HEID)

HARRIS	70	PRL 24 165	F. Harris <i>et al.</i>	(MICH, WISC)
PDG	70	RMP 42 87	A. Barbaro-Galtieri <i>et al.</i>	(LRL, BRAN+)
ANG	69B	ZPHY 228 151	G. Ang <i>et al.</i>	(HEID)
BAGGETT	69B	Thesis MDDP-TR-973	N.V. Baggett	(UMD)
BALTAY	69	PRL 22 615	C. Baltay <i>et al.</i>	(COLU, STON)
BANGERTER	69	Thesis UCRL 19244	R.O. Bangerter	(LRL)
BANGERTER	69B	PR 187 1821	R.O. Bangerter <i>et al.</i>	(LRL)
BARLOUTAUD	69	NP B14 153	R. Barloutaud <i>et al.</i>	(SACL, CERN, HEID)
EISELE	69	ZPHY 221 1	F. Eisele <i>et al.</i>	(HEID)
Also		PRL 13 291	W. Willis <i>et al.</i>	(BNL, CERN, HEID, UMD)
EISELE	69B	ZPHY 221 401	F. Eisele <i>et al.</i>	(HEID)
GERSHWIN	69B	PR 188 2077	L.K. Gershwin <i>et al.</i>	(LRL)
Also		Thesis UCRL 19246	L.K. Gershwin	(LRL)
NORTON	69	Thesis Nevis 175	H. Norton	(COLU)
BAGGETT	67	PRL 19 1458	N. Baggett <i>et al.</i>	(UMD)
Also		Vienna Abs. 374	N.V. Baggett, B. Kehoe	(UMD)
Also		Private Comm.	N.V. Baggett	(UMD)
BARASH	67	PRL 19 181	N. Barash <i>et al.</i>	(UMD)
EISELE	67	ZPHY 205 409	F. Eisele <i>et al.</i>	(HEID)
HYMAN	67	PL 25B 376	L.G. Hyman <i>et al.</i>	(ANL, CMU, NWES)
PDG	67	RMP 39 1	A.H. Rosenfeld <i>et al.</i>	(LRL, CERN, YALE)
CHANG	66	PR 151 1081	C.Y. Chang	(COLU)
Also		Thesis Nevis 145	C.Y. Chang	(COLU)
BAZIN	65	PRL 14 154	M. Bazin <i>et al.</i>	(PRIN, COLU)
BAZIN	65B	PR 140 B1358	M. Bazin <i>et al.</i>	(PRIN, RUTG, COLU)
SCHMIDT	65	PR 140 B1328	P. Schmidt	(COLU)
BHOWMIK	64	NP 53 22	B. Bhowmik <i>et al.</i>	(DELH)
COURANT	64	PR 136 B1791	H. Courant <i>et al.</i>	(CERN, HEID, UMD+)
NAUENBERG	64	PRL 12 679	U. Nauenberg <i>et al.</i>	(COLU, RUTG, PRIN)
BARKAS	63	PRL 11 26	W.H. Barkas, J.N. Dyer, H.H. Heckman	(LRL)
Also		Thesis UCRL 9450	J.N. Dyer	(LRL)
GALTIERI	62	PRL 9 26	A. Barbaro-Galtieri <i>et al.</i>	(LRL)
HUMPHREY	62	PR 127 1305	W.E. Humphrey, R.R. Ross	(LRL)
