

**$\Xi_b(5955)^-$**  $J^P = \frac{3}{2}^+$ 

Status: \*\*\*

 **$\Xi_b(5955)^-$  MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>5955.33±0.12±0.05</b>	<sup>1</sup> AAIJ	15H	LHCb $p p$ at 7, 8 TeV

<sup>1</sup> Not independent of the mass difference measurement below. Observed in  $\Xi_b^0 \pi^-$  channel with  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$ .

 **$m_{\Xi_b(5955)^-} - m_{\Xi_b^0} - m_{\pi^-}$** 

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>23.96±0.12±0.06</b>	<sup>1</sup> AAIJ	15H	LHCb $p p$ at 7, 8 TeV

<sup>1</sup> Observed in  $\Xi_b^0 \pi^-$  channel with  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$ .

 **$\Xi_b(5955)^-$  WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
<b>1.65±0.31±0.10</b>	<sup>1</sup> AAIJ	15H	LHCb $p p$ at 7, 8 TeV

<sup>1</sup> Observed in  $\Xi_b^0 \pi^-$  channel with  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$ .

 **$\Xi_b(5955)^-$  DECAY MODES**

Mode	Fraction ( $\Gamma_i/\Gamma$ )
$\Gamma_1 \quad \Xi_b^0 \pi^- \times B(\bar{b} \rightarrow \Xi_b^*(5955)^-) / B(\bar{b} \rightarrow \Xi_b^0)$	(20.7±3.5) %

 **$\Xi_b(5955)^-$  BRANCHING RATIOS**

$$\Gamma(\Xi_b^0 \pi^- \times B(\bar{b} \rightarrow \Xi_b^*(5955)^-) / B(\bar{b} \rightarrow \Xi_b^0)) / \Gamma_{\text{total}} \quad \Gamma_1 / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
<b>0.207±0.032±0.015</b>	<sup>1</sup> AAIJ	15H	LHCb $p p$ at 7, 8 TeV

<sup>1</sup> Observed in  $\Xi_b^0 \pi^-$  channel with  $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$  and  $\Xi_c^+ \rightarrow p K^- \pi^+$ .

 **$\Xi_b(5955)^-$  REFERENCES**

AAIJ

15H PRL 114 062004

R. Aaij *et al.*

(LHCb Collab.)