

$D_{s1}(2460)^\pm$

$$I(J^P) = 0(1^+)^\pm$$

See the review on "Heavy Non- $q\bar{q}$ Mesons."

$D_{s1}(2460)^\pm$ MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.


| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|------|-------------------------------------|----------|--|
| 2459.5±0.6 OUR FIT | | Error includes scale factor of 1.1. | | |
| 2459.6±0.9 OUR AVERAGE | | Error includes scale factor of 1.3. | | |
| 2460.1±0.2±0.8 | | ¹ AUBERT | 06P BABR | 10.6 e^+e^- |
| 2458.0±1.0±1.0 | 195 | AUBERT | 04E BABR | 10.6 e^+e^- |
| ● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ● | | | | |
| 2459.5±1.2±3.7 | 920 | AUBERT | 06P BABR | 10.6 $e^+e^- \rightarrow D_s^+ \gamma X$ |
| 2458.6±1.0±2.5 | 560 | AUBERT | 06P BABR | 10.6 $e^+e^- \rightarrow D_s^+ \pi^0 \gamma X$ |
| 2460.2±0.2±0.8 | 123 | AUBERT | 06P BABR | 10.6 $e^+e^- \rightarrow D_s^+ \pi^+ \pi^- X$ |
| 2458.9±1.5 | 112 | ² AUBERT,B | 04S BABR | $B \rightarrow D_{s1}(2460)^+ \bar{D}^{*0}$ |
| 2461.1±1.6 | 139 | ³ AUBERT,B | 04S BABR | $B \rightarrow D_{s1}(2460)^+ \bar{D}^{*0}$ |
| 2456.5±1.3±1.3 | 126 | ^{4,5} MIKAMI | 04 BELL | 10.6 e^+e^- |
| 2459.5±1.3±2.0 | 152 | ^{6,7} MIKAMI | 04 BELL | 10.6 e^+e^- |
| 2459.9±0.9±1.6 | 60 | ^{6,7} MIKAMI | 04 BELL | 10.6 e^+e^- |
| 2459.2±1.6±2.0 | 57 | KROKOVNY | 03B BELL | 10.6 e^+e^- |
| ¹ The average of the values obtained from the $D_s^+ \gamma$, $D_s^+ \pi^0 \gamma$, $D_s^+ \pi^+ \pi^-$ final state. | | | | |
| ² Systematic errors not evaluated. From the decay to $D_s^{*+} \pi^0$. | | | | |
| ³ Systematic errors not evaluated. From the decay to $D_s^+ \gamma$. | | | | |
| ⁴ Not independent of the corresponding $m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$. | | | | |
| ⁵ Using $m_{D_s^{*+}} = 2112.4 \pm 0.7$ MeV. | | | | |
| ⁶ Not independent of the corresponding $m_{D_{s1}(2460)^\pm} - m_{D_s^\pm}$. | | | | |
| ⁷ Using $m_{D_s^+} = 1968.5 \pm 0.6$ MeV. | | | | |

$m_{D_{s1}(2460)^\pm} - m_{D_s^{*\pm}}$

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^\pm$ mass and mass difference measurements.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|------|---|----------|---------------|
| 347.3±0.7 OUR FIT | | Error includes scale factor of 1.2. | | |
| 347.1±2.2 OUR AVERAGE | | Error includes scale factor of 1.9. See the ideogram below. | | |
| 344.1±1.3±1.1 | 126 | MIKAMI | 04 BELL | 10.6 e^+e^- |
| 351.2±1.7±1.0 | 41 | BESSON | 03 CLE2 | 10.6 e^+e^- |
| 346.8±1.6±1.9 | 57 | ⁸ KROKOVNY | 03B BELL | 10.6 e^+e^- |

⁸ Recalculated by us using $m_{D_s^{*+}} = 2112.4 \pm 0.7$ MeV.



$m_{D_{s1}(2460)^{\pm}} - m_{D_s^{\pm}}$

The fit includes D^{\pm} , D^0 , D_s^{\pm} , D_s^{*0} , $D_s^{*\pm}$, $D_1(2420)^0$, $D_2^*(2460)^0$, and $D_{s1}(2536)^{\pm}$ mass and mass difference measurements.

| VALUE (MeV) | EVTS | DOCUMENT ID | TECN | COMMENT |
|------------------------------|-------------------------------------|----------------------|---------|---------------|
| 491.1±0.6 OUR FIT | Error includes scale factor of 1.1. | | | |
| 491.3±1.4 OUR AVERAGE | | | | |
| 491.0±1.3±1.9 | 152 | ⁹ MIKAMI | 04 BELL | 10.6 e^+e^- |
| 491.4±0.9±1.5 | 60 | ¹⁰ MIKAMI | 04 BELL | 10.6 e^+e^- |

⁹ From the decay to $D_s^{\pm}\gamma$.

¹⁰ From the decay to $D_s^{\pm}\pi^+\pi^-$.

$D_{s1}(2460)^{\pm}$ WIDTH

| VALUE (MeV) | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
|---|-----|------|-------------|----------|--|
| < 3.5 | 95 | 123 | AUBERT | 06P BABR | 10.6 $e^+e^- \rightarrow D_s^+\pi^+\pi^-X$ |
| ●●● We do not use the following data for averages, fits, limits, etc. ●●● | | | | | |
| < 6.3 | 95 | 560 | AUBERT | 06P BABR | 10.6 $e^+e^- \rightarrow D_s^+\pi^0\gamma X$ |
| < 10 | | 195 | AUBERT | 04E BABR | 10.6 e^+e^- |

| | | | | | | |
|-------|----|-----|---------|----|------|------------------------------------|
| < 5.5 | 90 | 126 | MIKAMI | 04 | BELL | 10.6 e ⁺ e ⁻ |
| < 7 | 90 | 41 | BESSION | 03 | CLE2 | 10.6 e ⁺ e ⁻ |

$D_{s1}(2460)^+$ DECAY MODES

$D_{s1}(2460)^-$ modes are charge conjugates of the modes below.

| Mode | Fraction (Γ_i/Γ) | Scale factor/ Confidence level |
|--------------------------------------|--|-----------------------------------|
| Γ_1 $D_s^{*+} \pi^0$ | (48 ± 11) % | |
| Γ_2 $D_s^+ \gamma$ | (18 ± 4) % | |
| Γ_3 $D_s^+ \pi^+ \pi^-$ | (4.3 ± 1.3) % | S=1.1 |
| Γ_4 $D_s^{*+} \gamma$ | < 8 % | CL=90% |
| Γ_5 $D_{s0}^*(2317)^+ \gamma$ | (3.7 ^{+5.0} _{-2.4}) % | |
| Γ_6 $D_s^+ \pi^0$ | | |
| Γ_7 $D_s^+ \pi^0 \pi^0$ | | |
| Γ_8 $D_s^+ \gamma \gamma$ | | |

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 8 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 3.4$ for 4 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

| | | | |
|-------|-------|-------|-------|
| x_2 | 80 | | |
| x_3 | 68 | 62 | |
| x_5 | -3 | 25 | 26 |
| | x_1 | x_2 | x_3 |

$D_{s1}(2460)^\pm$ BRANCHING RATIOS

| $\Gamma(D_s^{*+} \pi^0)/\Gamma_{\text{total}}$ | | | | | Γ_1/Γ |
|--|------|-------------|------|---------|-------------------|
| VALUE | EVTS | DOCUMENT ID | TECN | COMMENT | |

0.48 ± 0.11 OUR FIT

0.56 ± 0.13 ± 0.09

¹¹ AUBERT 06N BABR $B \rightarrow D_{s1}(2460)^- \bar{D}^{(*)}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 41 BESSION 03 CLE2 10.6 e⁺e⁻

¹¹ Evaluated in AUBERT 06N including measurements from AUBERT,B 04s.

| $\Gamma(D_s^+ \gamma)/\Gamma_{\text{total}}$ | | | | Γ_2/Γ | |
|---|-----|------|----------------------|-------------------|---|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.18±0.04 OUR FIT | | | | | |
| 0.16±0.04±0.03 | | | ¹² AUBERT | 06N | BABR $B \rightarrow D_{s1}(2460)^- \bar{D}^{(*)}$ |
| ¹² Evaluated in AUBERT 06N including measurements from AUBERT,B 04s. | | | | | |

| $\Gamma(D_s^+ \gamma)/\Gamma(D_s^{*+} \pi^0)$ | | | | Γ_2/Γ_1 | |
|--|-----|------|------------------------|---------------------|---|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.38 ±0.05 OUR FIT | | | | | |
| 0.44 ±0.09 OUR AVERAGE | | | | | |
| 0.55 ±0.13 ±0.08 | | 152 | MIKAMI | 04 | BELL 10.6 $e^+ e^-$ |
| 0.38 ±0.11 ±0.04 | | 38 | KROKOVNY | 03B | BELL 10.6 $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| 0.274±0.045±0.020 | | 251 | ¹³ AUBERT,B | 04s | BABR $B \rightarrow D_{s1}(2460)^+ \bar{D}^{(*)}$ |
| < 0.49 | | 90 | BESSION | 03 | CLE2 10.6 $e^+ e^-$ |
| ¹³ Used by AUBERT 06N in their measurement of $B(D_s^{*-} \pi^0)$ and $B(D_s^- \gamma)$. | | | | | |

| $\Gamma(D_s^+ \pi^+ \pi^-)/\Gamma(D_s^{*+} \pi^0)$ | | | | Γ_3/Γ_1 | |
|---|-----|------|-------------|---------------------|-------------------------------------|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.090±0.020 OUR FIT | | | | | Error includes scale factor of 1.2. |
| 0.14 ±0.04 ±0.02 | | 60 | MIKAMI | 04 | BELL 10.6 $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.08 | | 90 | BESSION | 03 | CLE2 10.6 $e^+ e^-$ |

| $\Gamma(D_s^{*+} \gamma)/\Gamma(D_s^{*+} \pi^0)$ | | | | Γ_4/Γ_1 | |
|---|-----|------|-------------|---------------------|---------------------|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| <0.16 | | 90 | BESSION | 03 | CLE2 10.6 $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.31 | | 90 | MIKAMI | 04 | BELL 10.6 $e^+ e^-$ |

| $\Gamma(D_{s0}^*(2317)^+ \gamma)/\Gamma(D_s^{*+} \pi^0)$ | | | | Γ_5/Γ_1 | |
|---|-----|------|-------------|---------------------|---------------------|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| <0.22 | | 95 | AUBERT | 04E | BABR 10.6 $e^+ e^-$ |
| • • • We do not use the following data for averages, fits, limits, etc. • • • | | | | | |
| <0.58 | | 90 | BESSION | 03 | CLE2 10.6 $e^+ e^-$ |

| $\Gamma(D_s^{*+} \pi^0)/[\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | | | $\Gamma_1/(\Gamma_1+\Gamma_5)$ | |
|---|-----|------|-------------|--------------------------------|---------------------|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.93±0.09 OUR FIT | | | | | |
| 0.97±0.09±0.05 | | | AUBERT | 06P | BABR 10.6 $e^+ e^-$ |

| $\Gamma(D_s^+ \gamma)/[\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | | | $\Gamma_2/(\Gamma_1+\Gamma_5)$ | |
|---|-----|------|-------------|--------------------------------|---------------------|
| VALUE | CL% | EVTS | DOCUMENT ID | TECN | COMMENT |
| 0.35 ±0.04 OUR FIT | | | | | |
| 0.337±0.036±0.038 | | | AUBERT | 06P | BABR 10.6 $e^+ e^-$ |

| $\Gamma(D_s^+ \pi^+ \pi^-) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | $\Gamma_3 / (\Gamma_1 + \Gamma_5)$ | | |
|--|-----|------------------------------------|------|---|
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| 0.083 ± 0.017 OUR FIT | | | | Error includes scale factor of 1.2. |
| 0.077 ± 0.013 ± 0.008 | | AUBERT | 06P | BABR 10.6 e ⁺ e ⁻ |
| $\Gamma(D_s^{*+} \gamma) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | $\Gamma_4 / (\Gamma_1 + \Gamma_5)$ | | |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.24 | 95 | AUBERT | 06P | BABR 10.6 e ⁺ e ⁻ |
| $\Gamma(D_{s0}^*(2317)^+ \gamma) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | $\Gamma_5 / (\Gamma_1 + \Gamma_5)$ | | |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.25 | 95 | AUBERT | 06P | BABR 10.6 e ⁺ e ⁻ |
| $\Gamma(D_s^+ \pi^0) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | $\Gamma_6 / (\Gamma_1 + \Gamma_5)$ | | |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.042 | 95 | AUBERT | 06P | BABR 10.6 e ⁺ e ⁻ |
| $\Gamma(D_s^+ \pi^0 \pi^0) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | $\Gamma_7 / (\Gamma_1 + \Gamma_5)$ | | |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.68 | 95 | AUBERT | 06P | BABR 10.6 e ⁺ e ⁻ |
| $\Gamma(D_s^+ \gamma \gamma) / [\Gamma(D_s^{*+} \pi^0) + \Gamma(D_{s0}^*(2317)^+ \gamma)]$ | | $\Gamma_8 / (\Gamma_1 + \Gamma_5)$ | | |
| VALUE | CL% | DOCUMENT ID | TECN | COMMENT |
| <0.33 | 95 | AUBERT | 06P | BABR 10.6 e ⁺ e ⁻ |

$D_{s1}(2460)^\pm$ REFERENCES

| | | | | |
|----------|-----|---------------|---------------------------|-----------------|
| AUBERT | 06N | PR D74 031103 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 06P | PR D74 032007 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT | 04E | PR D69 031101 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| AUBERT,B | 04S | PRL 93 181801 | B. Aubert <i>et al.</i> | (BABAR Collab.) |
| MIKAMI | 04 | PRL 92 012002 | Y. Mikami <i>et al.</i> | (BELLE Collab.) |
| BESSON | 03 | PR D68 032002 | D. Besson <i>et al.</i> | (CLEO Collab.) |
| KROKOVNY | 03B | PRL 91 262002 | P. Krokovny <i>et al.</i> | (BELLE Collab.) |