

# GAUGE AND HIGGS BOSONS

**$\gamma$  (photon)**

$$I(J^{PC}) = 0,1(1^{--})$$

Mass  $m < 1 \times 10^{-18}$  eV

Charge  $q < 1 \times 10^{-46}$  e (mixed charge)

Charge  $q < 1 \times 10^{-35}$  e (single charge)

Mean life  $\tau =$  Stable

**$g$  (gluon)**

$$I(J^P) = 0(1^-)$$

Mass  $m = 0$  [a]

SU(3) color octet

**graviton**

$$J = 2$$

Mass  $m < 1.76 \times 10^{-23}$  eV

**$W$**

$$J = 1$$

Charge =  $\pm 1$  e

Mass  $m = 80.3692 \pm 0.0133$  GeV [b]

$W/Z$  mass ratio =  $0.88136 \pm 0.00015$

$m_Z - m_W = 10.818 \pm 0.013$  GeV

$m_{W^+} - m_{W^-} = -0.029 \pm 0.028$  GeV

Full width  $\Gamma = 2.085 \pm 0.042$  GeV

$\langle N_{\pi^\pm} \rangle = 15.70 \pm 0.35$

$\langle N_{K^\pm} \rangle = 2.20 \pm 0.19$

$\langle N_p \rangle = 0.92 \pm 0.14$

$\langle N_{\text{charged}} \rangle = 19.39 \pm 0.08$

$W^-$  modes are charge conjugates of the modes below.

| <b><math>W^+</math> DECAY MODES</b> | Fraction ( $\Gamma_i/\Gamma$ ) | Confidence level | $p$<br>(MeV/c) |
|-------------------------------------|--------------------------------|------------------|----------------|
| $\ell^+ \nu$                        | [c] $(10.86 \pm 0.09)$ %       |                  | –              |
| $e^+ \nu$                           | $(10.71 \pm 0.16)$ %           |                  | 40185          |
| $\mu^+ \nu$                         | $(10.63 \pm 0.15)$ %           |                  | 40185          |
| $\tau^+ \nu$                        | $(11.38 \pm 0.21)$ %           |                  | 40165          |
| hadrons                             | $(67.41 \pm 0.27)$ %           |                  | –              |
| $\pi^+ \gamma$                      | $< 7$                          | $\times 10^{-6}$ | 95% 40184      |

|                     |                       |                    |     |       |
|---------------------|-----------------------|--------------------|-----|-------|
| $D_s^+ \gamma$      | $< 6$                 | $\times 10^{-4}$   | 95% | 40160 |
| $cX$                | $(33.3 \pm 2.6) \%$   |                    |     | —     |
| $c\bar{s}$          | $(31^{+13}_{-11}) \%$ |                    |     | —     |
| invisible           | [d]                   | $(1.4 \pm 2.9) \%$ |     | —     |
| $\pi^+ \pi^+ \pi^-$ | $< 1.01$              | $\times 10^{-6}$   | 95% | 40184 |

**Z** $J = 1$ 

Charge = 0

Mass  $m = 91.1880 \pm 0.0020$  GeV [e]Full width  $\Gamma = 2.4955 \pm 0.0023$  GeV $\Gamma(\ell^+ \ell^-) = 83.984 \pm 0.086$  MeV [c] $\Gamma(\text{invisible}) = 499.2 \pm 1.5$  MeV [f] $\Gamma(\text{hadrons}) = 1744.4 \pm 2.0$  MeV $\Gamma(\mu^+ \mu^-) / \Gamma(e^+ e^-) = 1.0001 \pm 0.0024$  $\Gamma(\tau^+ \tau^-) / \Gamma(e^+ e^-) = 1.0020 \pm 0.0032$  [g]**Average charged multiplicity**

$$\langle N_{\text{charged}} \rangle = 20.76 \pm 0.16 \quad (S = 2.1)$$

**Couplings to quarks and leptons**

$$g_V^\ell = -0.03783 \pm 0.00041$$

$$g_V^u = 0.266 \pm 0.034$$

$$g_V^d = -0.38^{+0.04}_{-0.05}$$

$$g_A^\ell = -0.50123 \pm 0.00026$$

$$g_A^u = 0.519^{+0.028}_{-0.033}$$

$$g_A^d = -0.527^{+0.040}_{-0.028}$$

$$g^{\nu\ell} = 0.5008 \pm 0.0008$$

$$g^{\nu e} = 0.53 \pm 0.09$$

$$g^{\nu\mu} = 0.502 \pm 0.017$$

**Asymmetry parameters [h]**

$$A_e = 0.1515 \pm 0.0019$$

$$A_\mu = 0.142 \pm 0.015$$

$$A_\tau = 0.143 \pm 0.004$$

$$A_s = 0.90 \pm 0.09$$

$$A_c = 0.670 \pm 0.027$$

$$A_b = 0.923 \pm 0.020$$

**Charge asymmetry (%) at Z pole**

$$A_{FB}^{(0\ell)} = 1.71 \pm 0.10$$

$$A_{FB}^{(0u)} = 4 \pm 7$$

$$A_{FB}^{(0s)} = 9.8 \pm 1.1$$

$$A_{FB}^{(0c)} = 7.07 \pm 0.35$$

$$A_{FB}^{(0b)} = 9.92 \pm 0.16$$

| Z DECAY MODES   | Fraction ( $\Gamma_i/\Gamma$ )                                  | Scale factor/<br>Confidence level | $p$<br>(MeV/c) |
|---|---|-----------------------------------|----------------|
| $e^+ e^-$   | ( 3.3632±0.0042 ) %   |                                   | 45594          |
| $\mu^+ \mu^-$   | ( 3.3662±0.0066 ) %   |                                   | 45594          |
| $\tau^+ \tau^-$   | ( 3.3696±0.0083 ) %   |                                   | 45559          |
| $\ell^+ \ell^-$   | [c] ( 3.3658±0.0023 ) %   |                                   | —              |
| $\ell^+ \ell^- \ell^+ \ell^-$                           | [j] ( 4.55 ±0.17 ) × 10 <sup>-6</sup>                           |                                   | 45594          |
| invisible   | (20.000 ±0.055 ) %  |                                   | —              |
| hadrons   | (69.911 ±0.056 ) %  |                                   | —              |
| $(u\bar{u} + c\bar{c})/2$                               | (11.6 ±0.6 ) %  |                                   | —              |
| $(d\bar{d} + s\bar{s} + b\bar{b})/3$                    | (15.6 ±0.4 ) %  |                                   | —              |
| $c\bar{c}$  | (12.03 ±0.21 ) %  |                                   | —              |
| $b\bar{b}$  | (15.12 ±0.05 ) %  |                                   | —              |
| $b\bar{b}b\bar{b}$                                      | ( 3.6 ±1.3 ) × 10 <sup>-4</sup>                                 |                                   | —              |
| $g g g$   | < 1.1   | % CL=95%                          | —              |
| $\pi^0 \gamma$  | < 2.01  | × 10 <sup>-5</sup> CL=95%         | 45594          |
| $\eta \gamma$   | < 5.1   | × 10 <sup>-5</sup> CL=95%         | 45592          |
| $\rho^0 \gamma$   | < 4.0   | × 10 <sup>-6</sup> CL=95%         | 45591          |
| $\omega \gamma$   | < 3.9   | × 10 <sup>-6</sup> CL=95%         | 45591          |
| $\eta'(958) \gamma$                                     | < 4.2   | × 10 <sup>-5</sup> CL=95%         | 45589          |
| $\phi \gamma$   | < 7   | × 10 <sup>-7</sup> CL=95%         | 45588          |
| $\gamma \gamma$   | < 1.46  | × 10 <sup>-5</sup> CL=95%         | 45594          |
| $\pi^0 \pi^0$   | < 1.52  | × 10 <sup>-5</sup> CL=95%         | 45594          |
| $\gamma \gamma \gamma$                                  | < 2.2   | × 10 <sup>-6</sup> CL=95%         | 45594          |
| $\pi^\pm W^\mp$   | [j] < 7   | × 10 <sup>-5</sup> CL=95%         | 10176          |
| $\rho^\pm W^\mp$  | [j] < 8.3   | × 10 <sup>-5</sup> CL=95%         | 10151          |
| $J/\psi(1S) X$  | ( 3.51 <sup>+0.23</sup> / <sub>-0.25</sub> ) × 10 <sup>-3</sup> | S=1.1                             | —              |
| $J/\psi(1S) \gamma$                                     | < 1.2   | × 10 <sup>-6</sup> CL=95%         | 45541          |
| $\psi(2S) X$  | ( 1.60 ±0.29 ) × 10 <sup>-3</sup>                               |                                   | —              |
| $\psi(2S) \gamma$                                       | < 2.4   | × 10 <sup>-6</sup> CL=95%         | 45519          |
| $J/\psi(1S) J/\psi(1S)$                                 | < 2.2   | × 10 <sup>-6</sup> CL=95%         | 45489          |
| $\chi_{c1}(1P) X$                                       | ( 2.9 ±0.7 ) × 10 <sup>-3</sup>                                 |                                   | —              |
| $\chi_{c2}(1P) X$                                       | < 3.2   | × 10 <sup>-3</sup> CL=90%         | —              |
| $\Upsilon(1S) X + \Upsilon(2S) X$<br>+ $\Upsilon(3S) X$ | ( 1.0 ±0.5 ) × 10 <sup>-4</sup>                                 |                                   | —              |
| $\Upsilon(1S) X$  | < 4.4   | × 10 <sup>-5</sup> CL=95%         | —              |
| $\Upsilon(1S) \gamma$                                   | < 1.1   | × 10 <sup>-6</sup> CL=95%         | 45103          |
| $\Upsilon(2S) X$  | < 1.39  | × 10 <sup>-4</sup> CL=95%         | —              |
| $\Upsilon(2S) \gamma$                                   | < 1.3   | × 10 <sup>-6</sup> CL=95%         | 45043          |

|  |                                    |                  |        |       |
|--|------------------------------------|------------------|--------|-------|
| $\Upsilon(3S)X$                        | $< 9.4$                            | $\times 10^{-5}$ | CL=95% | —     |
| $\Upsilon(3S)\gamma$                   | $< 2.4$                            | $\times 10^{-6}$ | CL=95% | 45006 |
| $\Upsilon(1, 2, 3S)\Upsilon(1, 2, 3S)$ | $< 1.5$                            | $\times 10^{-6}$ | CL=95% | —     |
| $D^0\gamma$                            | $< 2.2$                            | $\times 10^{-3}$ | CL=95% | 45575 |
| $(D^0/\bar{D}^0)X$                     | (20.7 $\pm$ 2.0 ) %                |                  |        | —     |
| $D^\pm X$                              | (12.2 $\pm$ 1.7 ) %                |                  |        | —     |
| $D^*(2010)^\pm X$                      | [ <i>j</i> ] (11.4 $\pm$ 1.3 ) %   |                  |        | —     |
| $D_{s1}(2536)^\pm X$                   | ( 3.6 $\pm$ 0.8 ) $\times 10^{-3}$ |                  |        | —     |
| $D_{sJ}(2573)^\pm X$                   | ( 5.8 $\pm$ 2.2 ) $\times 10^{-3}$ |                  |        | —     |
| $D^{*l}(2629)^\pm X$                   | searched for                       |                  |        | —     |
| $B^+X$                                 | [ <i>k</i> ] ( 6.08 $\pm$ 0.13 ) % |                  |        | —     |
| $B_s^0X$                               | [ <i>k</i> ] ( 1.59 $\pm$ 0.13 ) % |                  |        | —     |
| $B_c^+X$                               | searched for                       |                  |        | —     |
| $\Lambda_c^+X$                         | ( 1.54 $\pm$ 0.33 ) %              |                  |        | —     |
| $\Xi_c^0X$                             | seen                               |                  |        | —     |
| $\Xi_b^0X$                             | seen                               |                  |        | —     |
| <i>b</i> -baryon $X$                   | [ <i>k</i> ] ( 1.38 $\pm$ 0.22 ) % |                  |        | —     |
| anomalous $\gamma$ + hadrons           | [ <i>l</i> ] $< 3.2$               | $\times 10^{-3}$ | CL=95% | —     |
| $e^+e^-\gamma$                         | [ <i>l</i> ] $< 5.2$               | $\times 10^{-4}$ | CL=95% | 45594 |
| $\mu^+\mu^-\gamma$                     | [ <i>l</i> ] $< 5.6$               | $\times 10^{-4}$ | CL=95% | 45594 |
| $\tau^+\tau^-\gamma$                   | [ <i>l</i> ] $< 7.3$               | $\times 10^{-4}$ | CL=95% | 45559 |
| $\ell^+\ell^-\gamma\gamma$             | [ <i>n</i> ] $< 6.8$               | $\times 10^{-6}$ | CL=95% | —     |
| $q\bar{q}\gamma\gamma$                 | [ <i>n</i> ] $< 5.5$               | $\times 10^{-6}$ | CL=95% | —     |
| $\nu\bar{\nu}\gamma\gamma$             | [ <i>n</i> ] $< 3.1$               | $\times 10^{-6}$ | CL=95% | 45594 |
| $e^\pm\mu^\mp$                         | LF [ <i>j</i> ] $< 2.62$           | $\times 10^{-7}$ | CL=95% | 45594 |
| $e^\pm\tau^\mp$                        | LF [ <i>j</i> ] $< 5.0$            | $\times 10^{-6}$ | CL=95% | 45577 |
| $\mu^\pm\tau^\mp$                      | LF [ <i>j</i> ] $< 6.5$            | $\times 10^{-6}$ | CL=95% | 45577 |
| $pe$                                   | L,B $< 1.8$                        | $\times 10^{-6}$ | CL=95% | 45589 |
| $p\mu$                                 | L,B $< 1.8$                        | $\times 10^{-6}$ | CL=95% | 45589 |

**H**

$$J = 0$$

was  $H^0$

$$\text{Mass } m = 125.20 \pm 0.11 \text{ GeV } (S = 1.4)$$

$$\text{Full width } \Gamma = 3.7_{-1.4}^{+1.9} \text{ MeV } (\text{assumes equal on-shell and off-shell effective couplings})$$

### **H Signal Strengths in Different Channels**

$$\text{Combined Final States} = 1.03 \pm 0.04$$

$$WW^* = 1.00 \pm 0.08$$

$$ZZ^* = 1.02 \pm 0.08$$

$$\gamma\gamma = 1.10 \pm 0.06$$

$$c\bar{c} \text{ Final State} < 14, \text{ CL} = 95\%$$

$$b\bar{b} = 0.99 \pm 0.12$$

$$\mu^+ \mu^- = 1.21 \pm 0.35$$

$$\tau^+ \tau^- = 0.91 \pm 0.09$$

$$Z\gamma = 2.2 \pm 0.7$$

$$\gamma^* \gamma \text{ Final State} = 1.5 \pm 0.5$$

$$\text{Fermion coupling } (\kappa_F) = 0.94 \pm 0.05$$

$$\text{Gauge boson coupling } (\kappa_V) = 1.023 \pm 0.026$$

$$t\bar{t}H \text{ Production} = 1.10 \pm 0.18$$

$$HH \text{ Production Cross Section in } pp \text{ Collisions} < 2.4, \text{ CL} = 95\%$$

$$tH \text{ production} = 6 \pm 4$$

$$H \text{ Production Cross Section in } pp \text{ Collisions at } \sqrt{s} = 13 \text{ TeV} = 56.8 \pm 3.4 \text{ pb}$$

| <b>H DECAY MODES</b>       | Fraction ( $\Gamma_i/\Gamma$ )      | Confidence level | $\rho$<br>(MeV/c) |
|----------------------------|-------------------------------------|------------------|-------------------|
| $W W^*$                    | (25.7 $\pm$ 2.5 ) %                 |                  | —                 |
| $Z Z^*$                    | ( 2.80 $\pm$ 0.30) %                |                  | —                 |
| $\gamma\gamma$             | ( 2.50 $\pm$ 0.20) $\times 10^{-3}$ |                  | 62600             |
| $b\bar{b}$                 | (53 $\pm$ 8 ) %                     |                  | —                 |
| $e^+ e^-$                  | < 3.0 $\times 10^{-4}$              | 95%              | 62600             |
| $\mu^+ \mu^-$              | ( 2.6 $\pm$ 1.3 ) $\times 10^{-4}$  |                  | 62600             |
| $\tau^+ \tau^-$            | ( 6.0 $^{+0.8}_{-0.7}$ ) %          |                  | 62575             |
| $Z\gamma$                  | ( 3.4 $\pm$ 1.1 ) $\times 10^{-3}$  |                  | 29392             |
| $Z\rho(770)$               | < 1.21 %                            | 95%              | 29384             |
| $Z\phi(1020)$              | < 3.6 $\times 10^{-3}$              | 95%              | 29378             |
| $ZJ/\psi$                  | < 1.9 $\times 10^{-3}$              | 95%              | 29267             |
| $Z\psi(2S)$                | < 6.6 $\times 10^{-3}$              | 95%              | 29214             |
| $J/\psi\gamma$             | < 2.0 $\times 10^{-4}$              | 95%              | 62561             |
| $J/\psi J/\psi$            | < 3.8 $\times 10^{-4}$              | 95%              | 62523             |
| $\psi(2S)\gamma$           | < 1.05 $\times 10^{-3}$             | 95%              | 62546             |
| $\psi(2S)J/\psi$           | < 2.1 $\times 10^{-3}$              | 95%              | 62507             |
| $\psi(2S)\psi(2S)$         | < 3.0 $\times 10^{-3}$              | 95%              | 62491             |
| $\Upsilon(1S)\gamma$       | < 2.5 $\times 10^{-4}$              | 95%              | 62242             |
| $\Upsilon(1S)\Upsilon(1S)$ | < 1.7 $\times 10^{-3}$              | 95%              | 61881             |
| $\Upsilon(2S)\gamma$       | < 4.2 $\times 10^{-4}$              | 95%              | 62199             |
| $\Upsilon(3S)\gamma$       | < 3.4 $\times 10^{-4}$              | 95%              | 62172             |
| $\Upsilon(nS)\Upsilon(mS)$ | < 3.5 $\times 10^{-4}$              | 95%              | —                 |
| $\rho(770)\gamma$          | < 1.04 $\times 10^{-3}$             | 95%              | 62597             |
| $\omega(782)\gamma$        | < 5.5 $\times 10^{-4}$              | 95%              | 62597             |
| $K^*(892)\gamma$           | < 2.2 $\times 10^{-4}$              | 95%              | 62597             |
| $\phi(1020)\gamma$         | < 5 $\times 10^{-4}$                | 95%              | 62596             |
| $e\mu$                     | LF < 4.4 $\times 10^{-5}$           | 95%              | 62600             |

|                    |      |          |                  |     |       |
|--------------------|------|----------|------------------|-----|-------|
| $e\tau$            | $LF$ | $< 2.0$  | $\times 10^{-3}$ | 95% | 62587 |
| $\mu\tau$          | $LF$ | $< 1.5$  | $\times 10^{-3}$ | 95% | 62587 |
| invisible          |      | $< 10.7$ | %                | 95% | —     |
| $\gamma$ invisible |      | $< 2.9$  | %                | 95% | —     |

## Neutral Higgs Bosons, Searches for

### Mass limits for heavy neutral Higgs bosons ( $H_2^0, A^0$ ) in the MSSM

|                          |                      |
|--------------------------|----------------------|
| $m > 1121$ GeV, CL = 95% | ( $\tan\beta = 10$ ) |
| $m > 1475$ GeV, CL = 95% | ( $\tan\beta = 20$ ) |
| $m > 1677$ GeV, CL = 95% | ( $\tan\beta = 30$ ) |
| $m > 1826$ GeV, CL = 95% | ( $\tan\beta = 40$ ) |
| $m > 1950$ GeV, CL = 95% | ( $\tan\beta = 50$ ) |
| $m > 2062$ GeV, CL = 95% | ( $\tan\beta = 60$ ) |

## Charged Higgs Bosons ( $H^\pm$ and $H^{\pm\pm}$ ), Searches for

### Mass limits for $m_{H^\pm} < m(\text{top})$ in the MSSM

|                         |
|-------------------------|
| $m > 155$ GeV, CL = 95% |
|-------------------------|

### Mass limits for $m_{H^\pm} > m(\text{top})$ in the MSSM

|                          |                      |
|--------------------------|----------------------|
| $m > 181$ GeV, CL = 95%  | ( $\tan\beta = 10$ ) |
| $m > 249$ GeV, CL = 95%  | ( $\tan\beta = 20$ ) |
| $m > 390$ GeV, CL = 95%  | ( $\tan\beta = 30$ ) |
| $m > 894$ GeV, CL = 95%  | ( $\tan\beta = 40$ ) |
| $m > 1017$ GeV, CL = 95% | ( $\tan\beta = 50$ ) |
| $m > 1103$ GeV, CL = 95% | ( $\tan\beta = 60$ ) |

## New Heavy Bosons ( $W', Z', \text{leptoquarks, etc.}$ ), Searches for

### Additional $W$ Bosons

|                                |                       |
|--------------------------------|-----------------------|
| $W'$ with standard couplings   |                       |
| Mass $m > 6000$ GeV, CL = 95%  | ( $pp$ direct search) |
| $W_R$ (Right-handed $W$ Boson) |                       |
| Mass $m > 715$ GeV, CL = 90%   | (electroweak fit)     |

### Additional Z Bosons

$Z'_{SM}$  with standard couplings

Mass  $m > 5150$  GeV, CL = 95% ( $pp$  direct search)

$Z_{LR}$  of  $SU(2)_L \times SU(2)_R \times U(1)$  (with  $g_L = g_R$ )

Mass  $m > 630$  GeV, CL = 95% ( $p\bar{p}$  direct search)

Mass  $m > 1162$  GeV, CL = 95% (electroweak fit)

$Z'_\chi$  of  $SO(10) \rightarrow SU(5) \times U(1)_\chi$  (with  $g_\chi = e/\cos\theta_W$ )

Mass  $m > 4800$  GeV, CL = 95% ( $pp$  direct search)

$Z'_\psi$  of  $E_6 \rightarrow SO(10) \times U(1)_\psi$  (with  $g_\psi = e/\cos\theta_W$ )

Mass  $m > 4560$  GeV, CL = 95% ( $pp$  direct search)

$Z'_\eta$  of  $E_6 \rightarrow SU(3) \times SU(2) \times U(1) \times U(1)_\eta$  (with  $g_\eta = e/\cos\theta_W$ )

Mass  $m > 3.900 \times 10^3$  GeV, CL = 95% ( $pp$  direct search)

### Scalar Leptoquarks

$m > 1800$  GeV, CL = 95% (1st gen., pair prod.,  $B(eq)=1$ )

$m > 1755$  GeV, CL = 95% (1st gen., single prod.,  $B(eq)=1$ )

$m > 1700$  GeV, CL = 95% (2nd gen., pair prod.,  $B(\mu q)=1$ )

$m > 660$  GeV, CL = 95% (2nd gen., single prod.,  $B(\mu q)=1$ )

$m > 1460$  GeV, CL = 95% (3rd gen., pair prod.,  $B(\tau b)=1$ )

$m > 1280$  GeV, CL = 95% (3rd gen., single prod.,  $B(\tau b)=1$ )

(See the Particle Listings for assumptions on leptoquark quantum numbers and branching fractions.)

### Diquarks

Mass  $m > 7200$  GeV, CL = 95% ( $E_6$  diquark)

### Axigluon

Mass  $m > 6600$  GeV, CL = 95%

## Axions ( $A^0$ ) and Other Very Light Bosons, Searches for

See the review on "Axions and other similar particles."

The best limit for the half-life of neutrinoless double beta decay with Majoron emission is  $> 7.2 \times 10^{24}$  years (CL = 90%).

## NOTES

- [a] Theoretical value. A mass as large as a few MeV may not be precluded.
- [b] This value does not include the AALTONEN 22 measurement by CDF. See the  $W$  mass section in the listings for details.
- [c]  $\ell$  indicates each type of lepton ( $e$ ,  $\mu$ , and  $\tau$ ), not sum over them.
- [d] This represents the width for the decay of the  $W$  boson into a charged particle with momentum below detectability,  $p < 200$  MeV.
- [e] The  $Z$ -boson mass listed here corresponds to a Breit-Wigner resonance parameter. It lies approximately 34 MeV above the real part of the position of the pole (in the energy-squared plane) in the  $Z$ -boson propagator.
- [f] This partial width takes into account  $Z$  decays into  $\nu\bar{\nu}$  and any other possible undetected modes.
- [g] This ratio has not been corrected for the  $\tau$  mass.
- [h] Here  $A \equiv 2g_V g_A / (g_V^2 + g_A^2)$ .
- [i] Here  $\ell$  indicates  $e$  or  $\mu$ .
- [j] The value is for the sum of the charge states or particle/antiparticle states indicated.
- [k] This value is updated using the product of (i) the  $Z \rightarrow b\bar{b}$  fraction from this listing and (ii) the  $b$ -hadron fraction in an unbiased sample of weakly decaying  $b$ -hadrons produced in  $Z$ -decays provided by the Heavy Flavor Averaging Group (HFLAV, <http://www.slac.stanford.edu/xorg/hflav/osc/PDG.2009/#FRACZ>).
- [l] See the  $Z$  Particle Listings for the  $\gamma$  energy range used in this measurement.
- [n] For  $m_{\gamma\gamma} = (60 \pm 5)$  GeV.